

**CYTOCHROME P450 MONOOXYGENASE AND
NADPH CYTOCHROME P450 OXIDOREDUCTASE GENES AND
PROTEINS RELATED TO THE OMEGA HYDROXYLASE COMPLEX OF
CANDIDA TROPICALIS AND METHODS RELATING THERETO**

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application Serial No. 60/123,555 filed March 10, 1999, U.S. Provisional Application Serial No. 60/103,099 filed October 5, 1998, and U.S. Provisional Application Serial No. 60/083,798 filed May 1, 1998.

BACKGROUND

1. Field of the Invention

The present invention relates to novel genes which encode enzymes of the ω -hydroxylase complex in yeast *Candida tropicalis* strains. In particular, the invention relates to novel genes encoding the cytochrome P450 and NADPH reductase enzymes of the ω -hydroxylase complex in yeast *Candida tropicalis*, and to a method of quantitating the expression of genes.

2. Description of the Related Art

Aliphatic dioic acids are versatile chemical intermediates useful as raw materials for the preparation of perfumes, polymers, adhesives and macrolid antibiotics. While several chemical routes to the synthesis of long-chain alpha, ω -dicarboxylic acids are available, the synthesis is not easy and most methods result in mixtures containing shorter chain lengths. As a result, extensive purification steps are necessary. While it is known that long-chain dioic acids can also be produced by microbial transformation of alkanes, fatty acids or esters thereof, chemical synthesis has remained the most commercially viable route, due to limitations with the current biological approaches.

Several strains of yeast are known to excrete alpha, ω -dicarboxylic acids as a byproduct when cultured on alkanes or fatty acids as the carbon source. In particular, yeast belonging to the Genus *Candida*, such as *C. albicans*, *C. cloacae*, *C. guilliermondii*, *C. intermedia*, *C. lipolytica*, *C. maltosa*, *C. parapsilosis* and *C. zeylenoides* are known to produce such dicarboxylic acids (*Agr. Biol. Chem.* 35: 2033-2042 (1971)). Also, various strains of *C. tropicalis* are known to produce dicarboxylic acids ranging in chain lengths from C₁₁ through C₁₈ (Okino et al., BM Lawrence, BD Mookherjee and BJ Willis (eds), in *Flavors and Fragrances: A World Perspective*. Proceedings of the 10th International Conference of Essential Oils, Flavors and Fragrances, Elsevier Science Publishers BV Amsterdam (1988)), and are the basis of several patents as reviewed by Bühler and Schindler, in *Aliphatic Hydrocarbons in Biotechnology*, H. J. Rehm and G. Reed (eds), Vol. 169, Verlag Chemie, Weinheim (1984).

Studies of the biochemical processes by which yeasts metabolize alkanes and fatty acids have revealed three types of oxidation reactions: α -oxidation of alkanes to alcohols, ω -oxidation of fatty acids to alpha, ω -dicarboxylic acids and the degradative β -oxidation of fatty acids to CO₂ and water. The first two types of oxidations are catalyzed by microsomal enzymes while the last type takes place in the peroxisomes. In *C. tropicalis*, the first step in the ω -oxidation pathway is catalyzed by a membrane-bound enzyme complex (ω -hydroxylase complex) including a cytochrome P450 monooxygenase and a NADPH cytochrome reductase. This hydroxylase complex is responsible for the primary oxidation of the terminal methyl group in alkanes and fatty acids (Gilewicz et al., *Can. J. Microbiol.* 25:201 (1979)). The genes which encode the cytochrome P450 and NADPH reductase components of the complex have previously been identified as P450ALK and P450RED respectively, and have also been cloned and sequenced (Sanglard et al., *Gene* 76:121-136 (1989)). P450ALK has also been designated P450ALK1. More recently, ALK genes have been designated by the symbol *CYP* and RED genes have been designated by the symbol *CPR*. See, e.g., Nelson, *Pharmacogenetics* 6(1):1-42 (1996), which is incorporated herein by reference. See also Ohkuma et al., *DNA and Cell Biology* 14:163-173 (1995), Seghezzi et al., *DNA and Cell Biology*, 11:767-780 (1992) and Kargel et al., *Yeast* 12:333-348 (1996), each incorporated herein by reference. For example, P450ALK

is also designated *CYP52* according to the nomenclature of Nelson, *supra*. Fatty acids are ultimately formed from alkanes after two additional oxidation steps, catalyzed by alcohol oxidase (Kemp et al., *Appl. Microbiol. and Biotechnol.* 28: 370-374 (1988)) and aldehyde dehydrogenase. The fatty acids can be further oxidized through the same or similar pathway to the corresponding dicarboxylic acid. The ω -oxidation of fatty acids proceeds via the ω -hydroxy fatty acid and its aldehyde derivative, to the corresponding dicarboxylic acid without the requirement for CoA activation. However, both fatty acids and dicarboxylic acids can be degraded, after activation to the corresponding acyl-CoA ester through the β -oxidation pathway in the peroxisomes, leading to chain shortening. In mammalian systems, both fatty acid and dicarboxylic acid products of ω -oxidation are activated to their CoA-esters at equal rates and are substrates for both mitochondrial and peroxisomal β -oxidation (*J. Biochem.*, 102:225-234 (1987)). In yeast, β -oxidation takes place solely in the peroxisomes (*Agr.Biol.Chem.* 49:1821-1828 (1985)).

It has recently been determined that certain eukaryotes, e.g., certain yeast, do not adhere, in some respects, to the "universal" genetic code which provides that particular codons (triplets of nucleic acids) code for specific amino acids. Indeed, the genetic code is "universal" because it is virtually the same in all living organisms. Certain *Candida* sp. are now known to translate the CTG codon (which, according to the "universal" code designates leucine) as serine. See, e.g., Ueda et al., *Biochemie* (1994) 76, 1217-1222, where *C. tropicalis*, *C. cylindracea*, *C. guilliermondii* and *C. lusitaniae* are shown to adhere to the "non-universal" code with respect to the CTG codon. Accordingly, nucleic acid sequences may code for one amino acid sequence in "universal" code organisms and a variant of that amino acid sequence in "non-universal" code organisms depending on the number of CTG codons present in the nucleic acid coding sequence. The difference may become evident when, in the course of genetic engineering, nucleic acid encoding a protein is transferred from a "non-universal" code organism to a "universal" code organism or *vice versa*. Obviously, there will be a different amino acid sequence depending on which organism is used to express the protein.

The production of dicarboxylic acids by fermentation of unsaturated C₁₄-C₁₆ monocarboxylic acids using a strain of the species *C. tropicalis* is disclosed in U.S. Patent


4,474,882. The unsaturated dicarboxylic acids correspond to the starting materials in the number and position of the double bonds. Similar processes in which other special microorganisms are used are described in U.S. Patents 3,975,234 and 4,339,536, in British Patent Specification 1,405,026 and in German Patent Publications 21 64 626, 28 53 847, 29 37 292, 29 51 177, and 21 40 133.

Cytochromes P450 (P450s) are terminal monooxidases of a multicomponent enzyme system as described above. They comprise a superfamily of proteins which exist widely in nature having been isolated from a variety of organisms as described e.g., in Nelson, *supra*. These organisms include various mammals, fish, invertebrates, plants, mollusk, crustaceans, lower eukaryotes and bacteria (Nelson, *supra*). First discovered in rodent liver microsomes as a carbon-monoxide binding pigment as described, e.g., in Garfinkel, *Arch. Biochem. Biophys.* 77:493-509 (1958), which is incorporated herein by reference, P450s were later named based on their absorption at 450 nm in a reduced-CO coupled difference spectrum as described, e.g., in Omura et al., *J. Biol. Chem.* 239:2370-2378 (1964), which is incorporated herein by reference.

P450s catalyze the metabolism of a variety of endogenous and exogenous compounds (Nelson, *supra*). Endogenous compounds include steroids, prostanoids, eicosanoids, fat-soluble vitamins, fatty acids, mammalian alkaloids, leukotrienes, biogenic amines and phytoalexins (Nelson, *supra*). P450 metabolism involves such reactions as epoxidation, hydroxylation, dealkylation, N-hydroxylation, sulfoxidation, desulfuration and reductive dehalogenation. These reactions generally make the compound more water soluble, which is conducive for excretion, and more electrophilic. These electrophilic products can have detrimental effects if they react with DNA or other cellular constituents. However, they can react through conjugation with low molecular weight hydrophilic substances resulting in glucoronidation, sulfation, acetylation, amino acid conjugation or glutathione conjugation typically leading to inactivation and elimination as described, e.g., in Klaassen et al., *Toxicology*, 3rd ed, Macmillan, New York, 1986, incorporated herein by reference.

P450s are heme thiolate proteins consisting of a heme moiety bound to a single polypeptide chain of 45,000 to 55,000 Da. The iron of the heme prosthetic group is located at the center of a protoporphyrin ring. Four ligands of the heme iron can be attributed to the porphyrin ring. The fifth ligand is a thiolate anion from a cysteinyl residue of the polypeptide. The sixth ligand is probably a hydroxyl group from an amino acid residue, or a moiety with a similar field strength such as a water molecule as described, e.g., in Goeptar et al., *Critical Reviews in Toxicology* 25(1):25-65 (1995), incorporated herein by reference.

Monooxygenation reactions catalyzed by cytochromes P450 in a eukaryotic membrane-bound system require the transfer of electrons from NADPH to P450 via NADPH-cytochrome P450 reductase (*CPR*) as described, e.g., in Taniguchi et al., *Arch. Biochem. Biophys.* 232:585 (1984), incorporated herein by reference. *CPR* genes are now also referred to as *NCP* genes. See, e.g., Debacker et al., *Antimicrobial Agents and Chemotherapy*, 45:1660 (2001). *CPR* is a flavoprotein of approximately 78,000 Da containing 1 mol of flavin adenine dinucleotide (FAD) and 1 mol of flavin mononucleotide (FMN) per mole of enzyme as described, e.g., in Potter et al., *J. Biol. Chem.* 258:6906 (1983), incorporated herein by reference. The FAD moiety of *CPR* is the site of electron entry into the enzyme, whereas FMN is the electron-donating site to P450 as described, e.g., in Vermilion et al., *J. Biol. Chem.* 253:8812 (1978), incorporated herein by reference. The overall reaction is as follows:



Binding of a substrate to the catalytic site of P450 apparently results in a conformational change initiating electron transfer from *CPR* to P450. Subsequent to the transfer of the first electron, O_2 binds to the Fe_2^+ -P450 substrate complex to form Fe_3^+ -P450-substrate complex. This complex is then reduced by a second electron from *CPR*, or, in some cases, NADH via cytochrome b5 and NADH-cytochrome b5 reductase as described, e.g., in Guengerich et al., *Arch. Biochem. Biophys.* 205:365 (1980), incorporated herein by reference. One atom of this reactive oxygen is introduced into the substrate, while the

other is reduced to water. The oxygenated substrate then dissociates, regenerating the oxidized form of the cytochrome P450 as described, e.g., in Klassen, Amdur and Doull, *Casarett and Doull's Toxicology*, Macmillan, New York (1986), incorporated herein by reference.

The P450 reaction cycle can be short-circuited in such a way that O₂ is reduced to O₂⁻ and/or H₂O₂ instead of being utilized for substrate oxygenation. This side reaction is often referred to as the "uncoupling" of cytochrome P450 as described, e.g., in Kuthen et al., *Eur. J. Biochem.* 126:583 (1982) and Poulos et al., *FASEBJ.* 6:674 (1992), both of which are incorporated herein by reference. The formation of these oxygen radicals may lead to oxidative cell damage as described, e.g., in Mukhopadhyay, *J. Biol. Chem.* 269(18):13390-13397 (1994) and Ross et al., *Biochem. Pharm.* 49(7):979-989 (1995), both of which are incorporated herein by reference. It has been proposed that cytochrome b5's effect on P450 binding to the *CPR* results in a more stable complex which is less likely to become "uncoupled" as described, e.g., in Yamazaki et al., *Arch. Biochem. Biophys.* 325(2):174-182 (1996), incorporated herein by reference.

P450 families are assigned based upon protein sequence comparisons. Notwithstanding a certain amount of heterogeneity, a practical classification of P450s into families can be obtained based on deduced amino acid sequence similarity. P450s with amino acid sequence similarity of between about 40 - 80% are considered to be in the same family, with sequences of about > 55% belonging to the same subfamily. Those with sequence similarity of about < 40% are generally listed as members of different P450 gene families (Nelson, *supra*). A value of about > 97% is taken to indicate allelic variants of the same gene, unless proven otherwise based on catalytic activity, sequence divergence in non-translated regions of the gene sequence, or chromosomal mapping.

The most highly conserved region is the HR2 consensus containing the invariant cysteine residue near the carboxyl terminus which is required for heme binding as described, e.g., in Gotoh et al. *J. Biochem.* 93:807-817 (1983) and Motohashi et al., *J. Biochem.* 101:879-997 (1987), both of which are incorporated herein by reference. Additional consensus regions, including the central region of helix I and the transmembrane region, have also been identified, as described, e.g., in Goeptar et al., *supra*

and Kalb et al., *PNAS*. 85:7221-7225 (1988), incorporated herein by reference, although the HR2 cysteine is the only invariant amino acid among P450s.

Short chain ($\leq C12$) aliphatic dicarboxylic acids (diacids) are important industrial intermediates in the manufacture of diesters and polymers, and find application as thermoplastics, plasticizing agents, lubricants, hydraulic fluids, agricultural chemicals, pharmaceuticals, dyes, surfactants, and adhesives. The high price and limited availability of short chain diacids are due to constraints imposed by the existing chemical synthesis.

Long-chain diacids (aliphatic α , ω -dicarboxylic acids with carbon numbers of 12 or greater, hereafter also referred to as diacids) ($HOOC-(CH_2)_n-COOH$) are a versatile family of chemicals with demonstrated and potential utility in a variety of chemical products including plastics, adhesives, and fragrances. Unfortunately, the full market potential of diacids has not been realized because chemical processes produce only a limited range of these materials at a relatively high price. In addition, chemical processes for the production of diacids have a number of limitations and disadvantages. All the chemical processes are restricted to the production of diacids of specific carbon chain lengths. For example, the dodecanedioic acid process starts with butadiene. The resulting product diacids are limited to multiples of four-carbon lengths and, in practice, only dodecanedioic acid is made. The dodecanedioic process is based on nonrenewable petrochemical feedstocks. The multireaction conversion process produces unwanted byproducts, which result in yield losses, NO_x pollution and heavy metal wastes.

Long-chain diacids offer potential advantages over shorter chain diacids, but their high selling price and limited commercial availability prevent widespread growth in many of these applications. Biocatalysis offers an innovative way to overcome these limitations with a process that produces a wide range of diacid products from renewable feedstocks. However, there is no commercially viable bioprocess to produce long chain diacids from renewable resources.

SUMMARY OF THE INVENTION

An isolated nucleic acid is provided which encodes a *CPRA* protein having the amino acid sequence set forth in SEQ ID NO: 83 or SEQ ID NO: 117. An isolated

nucleic acid is also provided which includes a coding region defined by nucleotides 1006-3042 as set forth in SEQ ID NO: 81. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 83 or SEQ ID NO: 117. A vector is provided which includes a nucleotide sequence encoding *CPRA* protein including an amino acid sequence as set forth in SEQ ID NO: 83 or SEQ ID NO: 117. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CPRA* protein having an amino acid sequence as set forth in SEQ ID NO: 83 or SEQ ID NO: 117. A method of producing a *CPRA* protein including an amino acid sequence as set forth in SEQ ID NO: 83 or SEQ ID NO: 117 is also provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 83 or SEQ ID NO: 117; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid is provided which encodes a *CPRB* protein having the amino acid sequence set forth in SEQ ID NO: 84 or SEQ ID NO: 118. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 1033-3069 as set forth in SEQ ID NO: 82. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 84 or SEQ ID NO: 118. A vector is provided which includes a nucleotide sequence encoding *CPRB* protein including an amino acid sequence as set forth in SEQ ID NO: 84 or SEQ ID NO: 118. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CPRB* protein having an amino acid sequence as set forth in SEQ ID NO: 84 or SEQ ID NO: 118. A method of producing a *CPRB* protein including an amino acid sequence as set forth in SEQ ID NO: 84 or SEQ ID NO: 118 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 84 or SEQ ID NO: 118; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid is provided which encodes a *CYP52A1A* protein having the amino acid sequence set forth in SEQ ID NO: 95 or SEQ ID NO: 110. An isolated nucleic acid is provided which includes a coding region defined by nucleotides

1177-2748 as set forth in SEQ ID NO: 85. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 95 or SEQ ID NO: 110. A vector is provided which includes a nucleotide sequence encoding *CYP52A1A* protein including an amino acid sequence as set forth in SEQ ID NO: 95 or SEQ ID NO: 110. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A1A* protein having an amino acid sequence as set forth in SEQ ID NO: 95 or SEQ ID NO: 110. A method of producing a *CYP52A1A* protein including an amino acid sequence as set forth in SEQ ID NO: 95 or SEQ ID NO: 110 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 95 or SEQ ID NO: 110; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52A2A* protein is provided which has the amino acid sequence set forth in SEQ ID NO: 96. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 1199-2767 as set forth in SEQ ID NO: 86. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 96. A vector is provided which includes a nucleotide sequence encoding *CYP52A2A* protein including an amino acid sequence as set forth in SEQ ID NO: 96. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A2A* protein having an amino acid sequence as set forth in SEQ ID NO: 96. A method of producing a *CYP52A2A* protein including an amino acid sequence as set forth in SEQ ID NO: 96 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 96; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52A2B* protein is provided which has the amino acid sequence set forth in SEQ ID NO: 97. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 1072-2640 as set forth in SEQ ID NO: 87. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 97. A vector is provided which includes a nucleotide

sequence encoding *CYP52A2B* protein including an amino acid sequence as set forth in SEQ ID NO: 97. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A2B* protein having an amino acid sequence as set forth in SEQ ID NO: 97. A method of producing a *CYP52A2B* protein including an amino acid sequence as set forth in SEQ ID NO: 97 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 97; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52A3A* protein is provided which has the amino acid sequence set forth in SEQ ID NO: 98. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 1126-2748 as set forth in SEQ ID NO: 88. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 98. A vector is provided which includes a nucleotide sequence encoding *CYP52A3A* protein including an amino acid sequence as set forth in SEQ ID NO: 98. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A3A* protein having an amino acid sequence as set forth in SEQ ID NO: 98. A method of producing a *CYP52A3A* protein including an amino acid sequence as set forth in SEQ ID NO: 98 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 98; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52A3B* protein is provided having the amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 913-2535 as set forth in SEQ ID NO: 89. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111. A vector is provided which includes a nucleotide sequence encoding *CYP52A3B* protein including an amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A3B*

protein having an amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111. A method of producing a *CYP52A3B* protein including an amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52A5A* protein is provided having the amino acid sequence set forth in SEQ ID NO: 100 or SEQ ID NO: 112. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 1103-2656 as set forth in SEQ ID NO: 90. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 100 or SEQ ID NO: 112. A vector is provided which includes a nucleotide sequence encoding *CYP52A5A* protein including an amino acid sequence as set forth in SEQ ID NO: 100 or SEQ ID NO: 112. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A5A* protein having an amino acid sequence as set forth in SEQ ID NO: 100 or SEQ ID NO: 112. A method of producing a *CYP52A5A* protein including an amino acid sequence as set forth in SEQ ID NO: 100 or SEQ ID NO: 112 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 100 or SEQ ID NO: 112; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52A5B* protein is provided having the amino acid sequence as set forth in SEQ ID NO: 101 or SEQ ID NO: 113. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 1142-2695 as set forth in SEQ ID NO: 91. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 101 or SEQ ID NO: 113. A vector is provided which includes a nucleotide sequence encoding *CYP52A5B* protein including the amino acid sequence as set forth in SEQ ID NO: 101 or SEQ ID NO: 113. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A5B* protein having the amino acid sequence as set forth in SEQ ID NO:

101 or SEQ ID NO: 113. A method of producing a *CYP52A5B* protein including an amino acid sequence as set forth in SEQ ID NO: 101 or SEQ ID NO: 113 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 101 or SEQ ID NO: 113; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52A8A* protein is provided having the amino acid sequence set forth in SEQ ID NO: 102 or SEQ ID NO: 114. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 464-2002 as set forth in SEQ ID NO: 92. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 102 or SEQ ID NO: 114. A vector is provided which includes a nucleotide sequence encoding *CYP52A8A* protein including an amino acid sequence as set forth in SEQ ID NO: 102 or SEQ ID NO: 114. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A8A* protein having an amino acid sequence as set forth in SEQ ID NO: 102 or SEQ ID NO: 114. A method of producing a *CYP52A8A* protein including an amino acid sequence as set forth in SEQ ID NO: 102 or SEQ ID NO: 114 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 102 or SEQ ID NO: 114; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52A8B* protein is provided having the amino acid sequence set forth in SEQ ID NO: 103 or SEQ ID NO: 115. An isolated nucleic acid is provided which includes a coding region defined by nucleotides 1017-2555 as set forth in SEQ ID NO: 93. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 103 or SEQ ID NO: 115. A vector is provided which includes a nucleotide sequence encoding *CYP52A8B* protein including an amino acid sequence as set forth in SEQ ID NO: 103 or SEQ ID NO: 115. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52A8B* protein having an amino acid sequence as set forth in SEQ ID NO: 103 or SEQ ID NO: 115. A method of producing a *CYP52A8B* protein including an

amino acid sequence as set forth in SEQ ID NO: 103 or SEQ ID NO: 115 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 103 or SEQ ID NO: 115; and b) culturing the cell under conditions favoring the expression of the protein.

An isolated nucleic acid encoding a *CYP52D4A* protein is provided having the amino acid sequence set forth in SEQ ID NO: 104 or SEQ ID NO: 116. An isolated nucleic acid is provided including a coding region defined by nucleotides 767-2266 as set forth in SEQ ID NO: 94. An isolated protein is provided which includes an amino acid sequence as set forth in SEQ ID NO: 104 or SEQ ID NO: 116. A vector is provided which includes a nucleotide sequence encoding *CYP52D4A* protein including an amino acid sequence as set forth in SEQ ID NO: 104 or SEQ ID NO: 116. A host cell is provided which is transfected or transformed with the nucleic acid encoding *CYP52D4A* protein having an amino acid sequence as set forth in SEQ ID NO: 104 or SEQ ID NO: 116. A method of producing a *CYP52D4A* protein including an amino acid sequence as set forth in SEQ ID NO: 104 or SEQ ID NO: 116 is provided which includes a) transforming a suitable host cell with a DNA sequence that encodes the protein having the amino acid sequence as set forth in SEQ ID NO: 104 or SEQ ID NO: 116; and b) culturing the cell under conditions favoring the expression of the protein.

A method for discriminating members of a gene family by quantifying the amount of target mRNA in a sample is provided which includes a) providing an organism containing a target gene; b) culturing the organism with an organic substrate which causes upregulation in the activity of the target gene; c) obtaining a sample of total RNA from the organism at a first point in time; d) combining at least a portion of the sample of the total RNA with a known amount of competitor RNA to form an RNA mixture, wherein the competitor RNA is substantially similar to the target mRNA but has a lesser number of nucleotides compared to the target mRNA; e) adding reverse transcriptase to the RNA mixture in a quantity sufficient to form corresponding target DNA and competitor DNA; (f) conducting a polymerase chain reaction in the presence of at least one primer specific for at least one substantially non-homologous region of the

target DNA within the gene family, the primer also specific for the competitor DNA; g) repeating steps (c-f) using increasing amounts of the competitor RNA while maintaining a substantially constant amount of target RNA; h) determining the point at which the amount of target DNA is substantially equal to the amount of competitor DNA; i) quantifying the results by comparing the ratio of the concentration of unknown target to the known concentration of competitor; and j) obtaining a sample of total RNA from the organism at another point in time and repeating steps (d-i).

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CPRA* genes; b) increasing, in the host cell, the number of *CPRA* genes which encode a *CPRA* protein having the amino acid sequence as set forth in SEQ ID NO: 83 or SEQ ID NO: 117; c) culturing the host cell in media containing an organic substrate which upregulates the *CPRA* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CPRA* protein having an amino acid sequence as set forth in SEQ ID NO: 83 or SEQ ID NO: 117 is provided which includes a) transforming a host cell having a naturally occurring amount of *CPRA* protein with an increased copy number of a *CPRA* gene that encodes the *CPRA* protein having the amino acid sequence as set forth in SEQ ID NO: 83 or SEQ ID NO: 117; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CPRA* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CPRB* genes; b) increasing, in the host cell, the number of *CPRB* genes which encode a *CPRB* protein having the amino acid sequence as set forth in SEQ ID NO: 84 or SEQ ID NO: 118; c) culturing the host cell in media containing an organic substrate which upregulates the *CPRB* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CPRB* protein having an amino acid sequence as set forth in SEQ ID NO: 84 or SEQ ID NO: 118 is provided which includes a) transforming a host cell having a naturally occurring amount of *CPRB*

protein with an increased copy number of a *CPRB* gene that encodes the *CPRB* protein having the amino acid sequence as set forth in SEQ ID NO: 84 or SEQ ID NO: 118; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CPRB* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A1A* genes; b) increasing, in the host cell, the number of *CYP52A1A* genes which encode a *CYP52A1A* protein having the amino acid sequence as set forth in SEQ ID NO: 95 or SEQ ID NO: 110; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52A1A* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A1A* protein having an amino acid sequence as set forth in SEQ ID NO: 95 or SEQ ID NO: 110 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A1A* protein with an increased copy number of a *CYP52A1A* gene that encodes the *CYP52A1A* protein having the amino acid sequence as set forth in SEQ ID NO: 95 or SEQ ID NO: 110; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A1A* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A2A* genes; b) increasing, in the host cell, the number of *CYP52A2A* genes which encode a *CYP52A2A* protein having the amino acid sequence as set forth in SEQ ID NO: 96; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52A2A* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A2A* protein having an amino acid sequence as set forth in SEQ ID NO: 96 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A2A* protein with an increased copy number of a *CYP52A2A* gene that encodes the *CYP52A2A* protein

having the amino acid sequence as set forth in SEQ ID NO: 96; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A2A* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A2B* genes; b) increasing, in the host cell, the number of *CYP52A2B* genes which encode a *CYP52A2B* protein having the amino acid sequence as set forth in SEQ ID NO: 97; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52A2B* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A2B* protein having an amino acid sequence as set forth in SEQ ID NO: 97 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A2B* protein with an increased copy number of a *CYP52A2B* gene that encodes the *CYP52A2B* protein having the amino acid sequence as set forth in SEQ ID NO: 97; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A2B* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A3A* genes; b) increasing, in the host cell, the number of *CYP52A3A* genes which encode a *CYP52A3A* protein having the amino acid sequence as set forth in SEQ ID NO: 98; c) culturing the host cell in media containing an organic substrate which upregulates *CYP52A3A* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A3A* protein having an amino acid sequence as set forth in SEQ ID NO: 98 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A3A* protein with an increased copy number of a *CYP52A3A* gene that encodes the *CYP52A3A* protein having the amino acid sequence as set forth in SEQ ID NO: 98; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A3A* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A3B* genes; b) increasing, in the host cell, the number of *CYP52A3B* genes which encode a *CYP52A3B* protein having the amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52A3B* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A3B* protein having an amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A3B* protein with an increased copy number of a *CYP52A3B* gene that encodes the *CYP52A3B* protein having the amino acid sequence as set forth in SEQ ID NO: 99 or SEQ ID NO: 111; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A3B* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A5A* genes; b) increasing, in the host cell, the number of *CYP52A5A* genes which encode a *CYP52A5A* protein having the amino acid sequence as set forth in SEQ ID NO: 100 or SEQ ID NO: 112; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52A5A* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A5A* protein having an amino acid sequence as set forth in SEQ ID NO: 100 or SEQ ID NO: 112 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A5A* protein with an increased copy number of a *CYP52A5A* gene that encodes the *CYP52A5A* protein having the amino acid sequence as set forth in SEQ ID NO: 100 or SEQ ID NO: 112; and b) culturing the cell and thereby increasing expression of the

protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A5A* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A5B* genes; b) increasing, in the host cell, the number of *CYP52A5B* genes which encode a *CYP52A5B* protein having the amino acid sequence as set forth in SEQ ID NO: 101 or SEQ ID NO: 113; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52A5B* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A5B* protein having an amino acid sequence as set forth in SEQ ID NO: 101 or SEQ ID NO: 113 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A5B* protein with an increased copy number of a *CYP52A5B* gene that encodes the *CYP52A5B* protein having the amino acid sequence as set forth in SEQ ID NO: 101 or SEQ ID NO: 113; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A5B* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A8A* genes; b) increasing, in the host cell, the number of *CYP52A8A* genes which encode a *CYP52A8A* protein having the amino acid sequence as set forth in SEQ ID NO: 102 or SEQ ID NO: 114; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52A8A* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A8A* protein having an amino acid sequence as set forth in SEQ ID NO: 102 or SEQ ID NO: 114 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A8A* protein with an increased copy number of a *CYP52A8A* gene that encodes the *CYP52A8A* protein having the amino acid sequence as set forth in SEQ ID NO: 102 or

SEQ ID NO: 114; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A8A* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52A8B* genes; b) increasing, in the host cell, the number of *CYP52A8B* genes which encode a *CYP52A8B* protein having the amino acid sequence as set forth in SEQ ID NO: 103 or SEQ ID NO: 115; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52A8B* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52A8B* protein having an amino acid sequence as set forth in SEQ ID NO: 103 or SEQ ID NO: 115 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52A8B* protein with an increased copy number of a *CYP52A8B* gene that encodes the *CYP52A8B* protein having the amino acid sequence as set forth in SEQ ID NO: 103 or SEQ ID NO: 115; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52A8B* gene.

A method for increasing production of a dicarboxylic acid is provided which includes a) providing a host cell having a naturally occurring number of *CYP52D4A* genes; b) increasing, in the host cell, the number of *CYP52D4A* genes which encode a *CYP52D4A* protein having the amino acid sequence as set forth in SEQ ID NO: 104 or SEQ ID NO: 116; c) culturing the host cell in media containing an organic substrate which upregulates the *CYP52D4A* gene, to effect increased production of dicarboxylic acid.

A method for increasing the production of a *CYP52D4A* protein having an amino acid sequence as set forth in SEQ ID NO: 104 or SEQ ID NO: 116 is provided which includes a) transforming a host cell having a naturally occurring amount of *CYP52D4A* protein with an increased copy number of a *CYP52D4A* gene that encodes the

CYP52D4A protein having the amino acid sequence as set forth in SEQ ID NO: 104 or SEQ ID NO: 116; and b) culturing the cell and thereby increasing expression of the protein compared with that of a host cell containing a naturally occurring copy number of the *CYP52D4A* gene.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic representation of cloning vector pTriplEx from Clontech™ Laboratories, Inc. Selected restriction sites within the multiple cloning site are shown.

Figure 2A is a map of the ZAP Express™ vector.

Figure 2B is a schematic representation of cloning phagemid vector pBK-CMV.

Figure 3 is a double stranded DNA sequence of a portion of the 5 prime coding region of the *CYP52A5A* gene (SEQ ID NO: 36), the non-coding or antisense sequence (SEQ ID NO: 108), primer 7581-97F (SEQ ID NO: 47) and primer 7581-97M (SEQ ID NO: 48).

Figure 4 is a diagrammatic representation of highly conserved regions of *CYP* and *CPR* gene protein sequences. Helix I represents the putative substrate binding site and HR2 represents the heme binding region. The FMN, FAD and NADPH binding regions are indicated below the *CPR* gene.

Figure 5 is a diagrammatic representation of the plasmid pHKM1 containing the truncated *CPRA* gene present in the pTriplEx vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 6 is a diagrammatic representation of the plasmid pHKM4 containing the truncated *CPRA* gene present in the pTriplEx vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 7 is a diagrammatic representation of the plasmid pHKM9 containing the *CPRB* gene (SEQ ID NO: 82) present in the pBK-CMV vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 8 is a diagrammatic representation of the plasmid pHKM11 containing the *CYP52A1A* gene (SEQ ID NO: 85) present in the pBK-CMV vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 9 is a diagrammatic representation of the plasmid pHKM12 containing the *CYP52A8A* gene (SEQ ID NO: 92) present in the pBK-CMV vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 10 is a diagrammatic representation of the plasmid pHKM13 containing the *CYP52D4A* gene (SEQ ID NO: 94) present in the pBK-CMV vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 11 is a diagrammatic representation of the plasmid pHKM14 containing the *CYP52A2B* gene (SEQ ID NO: 87) present in the pBK-CMV vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 12 is a diagrammatic representation of the plasmid pHKM15 containing the *CYP52A8B* gene (SEQ ID NO: 93) present in the pBK-CMV vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figures 13A-13D show the complete DNA sequences including regulatory and coding regions for the *CPRA* gene (SEQ ID NO: 81) and *CPRB* gene (SEQ ID NO: 82) from *C. tropicalis* ATCC 20336. Figures 13A-13D show regulatory and coding region alignment of these sequences. Asterisks indicate conserved nucleotides. The start codons are underlined and the last amino acid coding codons immediately before the stop codon are underlined.

Figure 14 shows the amino acid sequence of the *CPRA* (SEQ ID NO: 83) and *CPRB* (SEQ ID NO: 84) proteins from *C. tropicalis* ATCC 20336 and alignment of these amino acid sequences. Asterisks indicate residues which are not conserved.

Figures 15A-15M show the complete DNA sequences including regulatory and coding regions for the following genes from *C. tropicalis* ATCC 20366: *CYP52A1A* (SEQ ID NO: 85), *CYP52A2A* (SEQ ID NO: 86), *CYP52A2B* (SEQ ID NO: 87), *CYP52A3A* (SEQ ID NO: 88), *CYP52A3B* (SEQ ID NO: 89), *CYP52A5A* (SEQ ID NO: 90), *CYP52A5B* (SEQ ID NO: 91), *CYP52A8A* (SEQ ID NO: 92), *CYP52A8B* (SEQ ID NO: 93), and *CYP52D4A* (SEQ ID NO: 94). Figures 15A-15M show regulatory and coding region alignment of these sequences. Asterisks indicate conserved nucleotides. The start codons are underlined and the last amino acid coding codons immediately before the stop codon are underlined.

Figures 16A-16C show the amino acid sequences encoding the *CYP52A1A* (SEQ ID NO: 95), *CYP52A2A* (SEQ ID NO: 96), *CYP52A2B* (SEQ ID NO: 97), *CYP52A3A* (SEQ ID NO: 98), *CYP52A3B* (SEQ ID NO: 99), *CYP52A5A* (SEQ ID NO: 100), *CYP52A5B* (SEQ ID NO: 101), *CYP52A8A* (SEQ ID NO: 102), *CYP52A8B* (SEQ ID NO: 103) and *CYP52D4A* (SEQ ID NO: 104) proteins from *C. tropicalis* ATCC 20336. Asterisks indicate identical residues and dots indicate conserved residues.

Figure 17 is a diagrammatic representation of the pTAg PCR product cloning vector (commercially available from R&D Systems, Minneapolis, MN).

Figure 18 is a plot of the log ratio (U/C) of unknown target DNA product to competitor DNA product versus the concentration of competitor mRNA. The plot is used to calculate the target messenger RNA concentration in a quantitative competitive reverse transcription polymerase chain reaction (QC-RT-PCR).

Figure 19 is a graph showing the relative induction of *C. tropicalis* ATCC 20962 *CYP52A5A* (SEQ ID NO: 90) by the addition of the fatty acid substrate Emersol® 267 to the growth medium.

Figure 20 is a graph showing the induction of *C. tropicalis* ATCC 20962 *CYP52* and *CPR* genes by Emersol® 267. P450 genes *CYP52A3A* (SEQ ID NO: 88), *CYP52A3B* (SEQ ID NO: 89), and *CYP52D4A* (SEQ ID NO: 94) are expressed at levels below the detection level of the QC-RT-PCR assay.

Figure 21 is a scheme to integrate selected genes into the genome of *Candida tropicalis* strains and recovery of *URA3A* selectable marker.

Figure 22 is a schematic representation of the transformation of *C. tropicalis* H5343 *ura3* with *CYP* and/or *CPR* genes. Only one *URA3* locus needs to be functional. There are a total of 6 possible *ura3* targets (5 *ura3A* loci-2 *pox4* disruptions, 2 *pox 5* disruptions, 1 *ura3A* locus; and 1 *ura3B* locus).

Figure 23 is the complete DNA sequence (SEQ ID NO: 105) encoding *URA3A* from *C. tropicalis* ATCC 20336 and the amino acid sequence of the encoded protein (SEQ ID NO: 106).

Figure 24 is a schematic representation of the plasmid pURain, the base vector for integrating selected genes into the genome of *C. tropicalis*. The detailed construction of pURain is described in the text.

Figure 25 is a schematic representation of the plasmid pNEB193 cloning vector (commercially available from New England Biolabs, Beverly, MA).

Figure 26 is a diagrammatic representation of the plasmid pPA15 containing the truncated *CYP52A2A* gene present in the pTriplEx vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 27 is a schematic representation of pURA2in, the base vector is constructed in pNEB193 which contains the 8 bp recognition sequences for *Asc I*, *Pac I* and *Pme I*. *URA3A* (SEQ ID NO: 105) and *CYP52A2A* (SEQ ID NO: 86) do not contain these 8 bp recognition sites. *URA3A* is inverted so that the transforming fragment

will attempt to recircularize prior to integration. An *Asc I/Pme I* fragment was used to transform H5343 *ura*.

Figure 28 shows a scheme to detect integration of *CYP52A2A* gene (SEQ ID NO: 86) into the genome of H5343 *ura*. In all cases, hybridization band intensity could reflect the number of integrations.

Figure 29 is a diagrammatic representation of the plasmid pPA57 containing the truncated *CYP52A3A* gene present in the pTriplEx vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 30 is a diagrammatic representation of the plasmid pPA62 containing the truncated *CYP52A3B* gene present in the pTriplEx vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 31 is a diagrammatic representation of the plasmid pPAL3 containing the truncated *CYP52A5A* gene present in the pTriplEx vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 32 is a diagrammatic representation of the plasmid pPA5 containing the truncated *CYP52A5A* gene present in the pTriplEx vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 33 is a diagrammatic representation of the plasmid pPA18 containing the truncated *CYP52D4A* gene present in the pTriplEx vector. A detailed restriction map of only the sequenced region is shown at the top. The bar indicates the open reading frame. The direction of transcription is indicated by an arrow under the open reading frame.

Figure 34 is a graph showing the expression of *CYP52A1* (SEQ ID NO: 85), *CYP52A2* (SEQ ID NO: 86) and *CYP52A5* genes (SEQ ID NOS: 90 and 91) from *C. tropicalis* 20962 in a fermentor run upon the addition of amounts of the substrate oleic acid or tridecane in a spiking experiment.

Figure 35 depicts a scheme used for the extraction and analysis of diacids and monoacids from fermentation broths.

Figure 36 is a graph showing the induction of expression of *CYP52A1A*, *CYP52A2A* and *CYP52A5A* in a fermentor run upon addition of the substrate octadecane. No induction of *CYP52A3A* or *CYP52A3B* was observed under these conditions.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

Diacid productivity is improved according to the present invention by selectively increasing enzymes which are known to be important to the oxidation of organic substrates such as fatty acids composing the desired feed. According to the present invention, ten *CYP* genes and two *CPR* genes of *C. tropicalis* have been identified and characterized that relate to participation in the ω -hydroxylase complex catalyzing the first step in the ω -oxidation pathway. In addition, a novel quantitative competitive reverse transcription polymerase chain reaction (QC-RT-PCR) assay is used to measure gene expression in the fermentor under conditions of induction by one or more organic substrates as defined herein. Based upon QC-RT-PCR results, three *CYP* genes, *CYP52A1*, *CYP52A2* and *CYP52A5*, have been identified as being of greater importance for the ω -oxidation of long chain fatty acids. Amplification of the *CPR* gene copy number improves productivity. The QC-RT-PCR assay indicates that both *CYP* and *CPR* genes appear to be under tight regulatory control.

In accordance with the present invention, a method for discriminating members of a gene family by quantifying the amount of target mRNA in a sample is provided which includes a) providing an organism containing a target gene; b) culturing the organism with an organic substrate which causes upregulation in the activity of the target gene; c) obtaining a sample of total RNA from the organism at a first point in time;

d) combining at least a portion of the sample of the total RNA with a known amount of competitor RNA to form an RNA mixture, wherein the competitor RNA is substantially similar to the target mRNA but has a lesser number of nucleotides compared to the target mRNA; e) adding reverse transcriptase to the RNA mixture in a quantity sufficient to form corresponding target DNA and competitor DNA; (f) conducting a polymerase chain reaction in the presence of at least one primer specific for at least one substantially non-homologous region of the target DNA within the gene family, the primer also specific for the competitor DNA; g) repeating steps (c-f) using increasing amounts of the competitor RNA while maintaining a substantially constant amount of target RNA; h) determining the point at which the amount of target DNA is substantially equal to the amount of competitor DNA; i) quantifying the results by comparing the ratio of the concentration of unknown target to the known concentration of competitor; and j) obtaining a sample of total RNA from the organism at another point in time and repeating steps (d-i).

In addition, modification of existing promoters and/or the isolation of alternative promoters provides increased expression of *CYP* and *CPR* genes. Strong promoters are obtained from at least four sources: random or specific modifications of the *CYP52A2* promoter, *CYP52A5* promoter, *CYP52A1* promoter, the selection of a strong promoter from available *Candida* β -oxidation genes such as *POX4* and *POX5*, or screening to select another suitable *Candida* promoter.

Promoter strength can be directly measured using QT-RT-PCR to measure *CYP* and *CPR* gene expression in *Candida* cells isolated from fermentors. Enzymatic assays and antibodies specific for *CYP* and *CPR* proteins are used to verify that increased promoter strength is reflected by increased synthesis of the corresponding enzymes. Once a suitable promoter is identified, it is fused to the selected *CYP* and *CPR* genes and introduced into *Candida* for construction of a new improved production strain. It is contemplated that the coding region of the *CYP* and *CPR* genes can be fused to suitable promoters or other regulatory sequences which are well known to those skilled in the art.

In accordance with the present invention, studies on *C. tropicalis* ATCC 20336 have identified six unique *CYP* genes and four potential alleles. QC-RT-PCR

analyses of cells isolated during the course of the fermentation bioconversions indicate that at least three of the *CYP* genes are induced by fatty acids and at least two of the *CYP* genes are induced by alkanes. See Figure 34. Two of the *CYP* genes are highly induced indicating participation in the ω -hydroxylase complex which catalyzes the rate limiting step in the oxidation of fatty acids to the corresponding diacids.

The biochemical characterizations of each P450 enzyme herein is used to tailor the *C. tropicalis* host for optimal diacid productivity and is used to select P450 enzymes to be amplified based upon the fatty acid content of the feedstream. *CYP* gene(s) encoding P450 enzymes that have a low specific activity for the fatty acid or alkane substrate of choice are targeted for inactivation, thereby reducing the physiological load on the cell.

Since it has been demonstrated that *CPR* can be limiting in yeast systems, the removal of non-essential P450s from the system can free electrons that are being used by non-essential P450s and make them available to the P450s important for diacid productivity. Moreover, the removal of non-essential P450s can make available other necessary but potentially limiting components of the P450 system (i.e., available membrane space, heme and/or NADPH).

Diacid productivity is thus improved by selective integration, amplification, and over expression of *CYP* and *CPR* genes in the *C. tropicalis* production host.

It should be understood that host cells into which one or more copies of desired *CYP* and/or *CPR* genes have been introduced can be made to include such genes by any technique known to those skilled in the art. For example, suitable host cells include procaryotes such as *Bacillus sp.*, *Pseudomonas sp.*, *Actinomyces sp.*, *Escherichia sp.*, *Mycobacterium sp.*, and eukaryotes such as yeast, algae, insect cells, plant cells and filamentous fungi. Suitable host cells are preferably yeast cells such as *Yarrowia*, *Bebaromyces*, *Saccharomyces*, *Schizosaccharomyces*, and *Pichia* and more preferably those of the *Candida* genus. Preferred species of *Candida* are *tropicalis*, *maltosa*, *apicola*, *paratropicalis*, *albicans*, *cloacae*, *guilliermondii*, *intermedia*, *lipolytica*, *parapsilosis* and *zeylenoides*. Certain preferred strains of *Candida tropicalis* are listed in U.S. Patent No. 5,254,466, incorporated herein by reference.

Vectors such as plasmids, phagemids, phages or cosmids can be used to transform or transfect suitable host cells. Host cells may also be transformed by introducing into a cell a linear DNA vector(s) containing the desired gene sequence. Such linear DNA may be advantageous when it is desirable to avoid introduction of non-native (foreign) DNA into the cell. For example, DNA consisting of a desired target gene(s) flanked by DNA sequences which are native to the cell can be introduced into the cell by electroporation, lithium acetate transformation, spheroplasting and the like. Flanking DNA sequences can include selectable markers and/or other tools for genetic engineering.

It should be understood that, depending on whether a transformed organism utilizes the universal genetic code or the non-universal genetic code known, e.g., in connection with *C. tropicalis*, slight differences can be manifest in the amino acid sequences of protein-products. Thus, nucleotide sequences containing a CTG codon produce proteins containing a CTG encoded leucine in prokaryotes such as *E. coli* and a CTG encoded serine in non-universal coding eukaryotes such as *C. tropicalis*. For example, the *CYP52A1A* gene contains one CTG codon starting at position 1354 which is translated as a leucine in *E. coli* and a serine in *C. tropicalis*, leading to two versions of the CYP52A1A protein (SEQ. ID. NO: 95 and SEQ. ID. NO: 110); the *CYP52A3B* gene contains one CTG codon starting at position 2449 which is translated as a leucine in *E. coli* and a serine in *C. tropicalis*, leading to two versions of the CYP52A3B protein (SEQ. ID. NO: 99 and SEQ. ID. NO: 111); the *CYP52A5A* gene contains two CTG codons starting, respectively, at positions 1883 and 2570, which are translated as leucine in *E. coli* and serine in *C. tropicalis*, leading to two versions of the CYP52A5A protein (SEQ. ID. NO: 100 and SEQ. ID. NO: 112); the *CYP52A5B* gene contains two CTG codons starting, respectively, at positions 1922 and 2609, which are translated as leucine in *E. coli* and serine in *C. tropicalis*, leading to two versions of the CYP52A5B protein (SEQ. ID. NO: 101 and SEQ. ID. NO: 113); the *CYP52A8A* gene contains one CTG codon starting at position 659, which is translated as a leucine in *E. coli* and a serine in *C. tropicalis*, leading to two versions of the CYP52A8B protein (SEQ. ID. NO: 103 and SEQ. ID. NO: 115); the *CYP52D4A* gene contains three CTG codons starting, respectively, at positions 1247, 1412 and 1757, which are translated as leucine in *E. coli* and as serine in *C. tropicalis*,

leading to two versions of the CYP5234A protein (SEQ. ID. NO: 104 and SEQ. ID. NO: 116); the *CPRA* (*NCPIA*) gene contains one CTG codon starting at position 1153 which is translated as a leucine in *E. coli* and as a serine in *C. tropicalis*, leading to two versions of the CPRA (NCP1A) protein (SEQ. ID. NO: 83 and SEQ. ID. NO: 117); the *CPRG* (*NCPIB*) gene contains one CTG codon starting at position 1180 which is translated as a leucine in *E. coli* and as a serine in *C. tropicalis*, leading to two versions of the CPRB (NCP1B) protein (SEQ. ID. NO: 84 and SEQ. ID. NO: 118).

A suitable organic substrate herein can be any organic compound that is biooxidizable to a mono- or polycarboxylic acid. Such a compound can be any saturated or unsaturated aliphatic compound or any carbocyclic or heterocyclic aromatic compound having at least one terminal methyl group, a terminal carboxyl group and/or a terminal functional group which is oxidizable to a carboxyl group by biooxidation. A terminal functional group which is a derivative of a carboxyl group may be present in the substrate molecule and may be converted to a carboxyl group by a reaction other than biooxidation. For example, if the terminal group is an ester that neither the wild-type *C. tropicalis* nor the genetic modifications described herein will allow hydrolysis of the ester functionality to a carboxyl group, then a lipase can be added during the fermentation step to liberate free fatty acids. Suitable organic substrates include, but are not limited to, saturated fatty acids, unsaturated fatty acids, alkanes, alkenes, alkynes and combinations thereof.

Alkanes are a type of saturated organic substrate which are useful herein. The alkanes can be linear or cyclic, branched or straight chain, substituted or unsubstituted. Particularly preferred alkanes are those having from about 4 to about 25 carbon atoms, examples of which include but are not limited to butane, hexane, octane, nonane, dodecane, tridecane, tetradecane, octadecane and the like.

Examples of unsaturated organic substrates which can be used herein include but are not limited to internal olefins such as 2-pentene, 2-hexene, 3-hexene, 9-octadecene and the like; unsaturated carboxylic acids such as 2-hexenoic acid and esters thereof, oleic acid and esters thereof including triglyceryl esters having a relatively high oleic acid content, erucic acid and esters thereof including triglyceryl esters having a relatively high erucic acid content, ricinoleic acid and esters thereof including triglyceryl esters having

a relatively high ricinoleic acid content, linoleic acid and esters thereof including triglyceryl esters having a relatively high linoleic acid content; unsaturated alcohols such as 3-hexen-1-ol, 9-octadecen-1-ol and the like; unsaturated aldehydes such as 3-hexen-1-al, 9-octadecen-1-al and the like. In addition to the above, an organic substrate which can be used herein include alicyclic compounds having at least one internal carbon-carbon double bond and at least one terminal methyl group, a terminal carboxyl group and/or a terminal functional group which is oxidizable to a carboxyl group by biooxidation. Examples of such compounds include but are not limited to 3,6-dimethyl, 1,4-cyclohexadiene; 3-methylcyclohexene; 3-methyl-1, 4-cyclohexadiene and the like.

Examples of the aromatic compounds that can be used herein include but are not limited to arenes such as o-, m-, p-xylene; o-, m-, p-methyl benzoic acid; dimethyl pyridine, and the like. The organic substrate can also contain other functional groups that are biooxidizable to carboxyl groups such as an aldehyde or alcohol group. The organic substrate can also contain other functional groups that are not biooxidizable to carboxyl groups and do not interfere with the biooxidation such as halogens, ethers, and the like.

Examples of saturated fatty acids which may be applied to cells incorporating the present *CYP* and *CPR* genes include caproic, enanthic, caprylic, pelargonic, capric, undecylic, lauric, myristic, pentadecanoic, palmitic, margaric, stearic, arachidic, behenic acids and combinations thereof. Examples of unsaturated fatty acids which may be applied to cells incorporating the present *CYP* and *CPR* genes include palmitoleic, oleic, erucic, linoleic, linolenic acids and combinations thereof. Alkanes and fractions of alkanes may be applied which include chain links from C12 to C24 in any combination. An example of a preferred fatty acid mixtures are Emersol® 267 and Tallow, both commercially available from Henkel Chemicals Group, Cincinnati, OH. The typical fatty acid composition of Emersol® 267 and Tallow is as follows:

	<u>TALLOW</u>	<u>E267</u>
C14:0	3.5%	2.4%
C14:1	1.0%	0.7%
C15:0	0.5%	—
C16:0	25.5%	4.6%
C16:1	4.0%	5.7%
C17:0	2.5%	—
C17:1	—	5.7%
C18:0	19.5%	1.0%
C18:1	41.0%	69.9%
C18:2	2.5%	8.8%
C18:3	—	0.3%
C20:0	0.5%	—
C20:1	—	0.9%

The following examples are meant to illustrate but not to limit the invention. All relevant microbial strains and plasmids are described in Table 1 and Table 2, respectively.

Table 1. List of *Escherichia coli* and *Candida tropicalis* strains

<i>E. Coli</i> STRAIN	GENOTYPE	SOURCE
XL1Blue- MRF'	<i>endA1, gyrA96, hsdR17, lac, recA1, relA1, supE44, thi-1, [F' lacI⁺Z M15, proAB, Tn10]</i>	Stratagene, La Jolla, CA
BM25.8	<i>SupE44, thi (lac-proAB) [F' traD36, proAB, lacI⁺Z M15] λimm434 (kan^r)P1 (cam^r) hsdR (<i>t_{lir}-m_{lir}</i>)</i>	Clontech, Palo Alto, CA
XL0LR	<i>(mcrA)183 (mcrCB-hsdSMR-mrr)173 endA1 thi-1 recA1 gyrA96 relA1 lac [F' proAB lacI⁺Z M15 Tn10 (Tet) Su^r (nonsuppressing λ' (lambda resistant)</i>	Stratagene, La Jolla, CA

<i>C. tropicalis</i> STRAIN	GENOTYPE	SOURCE
ATCC20336	Wild-type	American Type Culture Collection, Rockville, MD
ATCC750	Wild-type	American Type Culture Collection, Rockville, MD
ATCC 20962	<i>ura3A/ura3B, pox4A::ura3A/pox4B::ura3A, pox5::ura3A/pox5::URA3A</i>	Henkel
H5343 ura-	<i>ura3A/ura3B, pox4A::ura3A/pox4B::ura3A, pox5::ura3A/pox5::URA3A, ura3-</i>	Henkel
HDC1	<i>ura3A/ura3B, pox4A::ura3A/pox4B::ura3A, pox5::ura3A/pox5::URA3A, ura3::URA3A-CYP52A2A</i>	Henkel
HDC5	<i>ura3A/ura3B, pox4A::ura3A/pox4B::ura3A, pox5::ura3A/pox5::URA3A, ura3::URA3A-CYP52A3A</i>	Henkel
HDC10	<i>ura3A/ura3B, pox4A::ura3A/pox4B::ura3A, pox5::ura3A/pox5::URA3A, ura3::URA3A-CPRB</i>	Henkel
HDC15	<i>ura3A/ura3B, pox4A::ura3A/pox4B::ura3A, pox5::ura3A/pox5::URA3A, ura3::URA3A-CYP52A5A</i>	Henkel
HDC20	<i>ura3A/ura3B, pox4A::ura3A/pox4B::ura3A, pox5::ura3A/pox5::URA3A, ura3::URA3A-CYP52A2A + CPR B (CYP and CPR have opposite 5' to 3' orientation with respect to each other)</i>	Henkel

HDC23	<i>ura3A/ura3B</i> , <i>pox4A::ura3A/pox4B::ura3A</i> , <i>pox5::ura3A/pox5::URA3A</i> , <i>ura3::URA3A-CYP52A2A + CPR B</i> (<i>CYP</i> and <i>CPR</i> have same 5' to 3' orientation with respect to each other)	Henkel
-------	--	--------

Table 2. List of plasmids isolated from genomic libraries and constructed for use in gene integrations.

Plasmid	Base vector	Insert	Insert Size	Plasmid size	Description
pURain	pNEB193	<i>URA3A</i>	1706 bp	4399 bp	pNEB193 with the <i>URA3A</i> gene inserted in the <i>AscI</i> - <i>PmeI</i> site, generating a <i>PacI</i> site
pURA 2in	pURain	<i>CYP52A2 A</i>	2230 bp	6629 bp	pURain containing a PCR <i>CYP52A2A</i> allele containing <i>PacI</i> restriction sites
pURA REDB in	pURain	<i>CPRB</i>	3266 bp	7665 bp	pURain containing a PCR <i>CPRB</i> allele containing <i>PacI</i> restriction sites
pHKM1	pTriplEx	Truncated <i>CPRA</i> gene	Approx. 3.8 kb	Approx. 7.4 kb	A truncated <i>CPRA</i> gene obtained by first screening library containing the 5' untranslated region and 1.2 kb open reading frame
pHKM4	PTriplEx	Truncated <i>CPRA</i> gene	Approx. 5 kb	Approx. 8.6 kb	A truncated <i>CPRA</i> gene obtained by screening second library containing the 3' untranslated region end sequence
pHKM9	pBC-CMV	<i>CPRB</i> gene	Approx. 5.3 kb	Approx. 9.8 kb	<i>CPRB</i> allele isolated from the third library
pHKM11	pBC-CMV	<i>CYP52A1 A</i>	Approx. 5 kb	Approx. 9.5 kb	<i>CYP52A1A</i> isolated from the third library
pHKM12	pBC-CMV	<i>CYP52A8 A</i>	Approx. 7.5 kb	Approx. 12 kb	<i>CYP52A8A</i> isolated from the third library
pHKM13	pBC-CMV	<i>CYP52D4 A</i>	Approx. 7.3 kb	Approx. 11.8 kb	<i>CYP52D4A</i> isolated from the third library
pHKM14	pBC-CMV	<i>CYP52A2 B</i>	Approx. 6 kb	Approx. 10.5 kb	<i>CYP52A2B</i> isolated from the third library
pHKM15	pBC-CMV	<i>CYP52A8 B</i>	Approx. 6.6 kb	Approx. 11.1 kb	<i>CYP52A8B</i> isolated from the third library

pPAL3	pTriplEx	<i>CYP52A5</i> <i>A</i>	4.4 kb	Approx. 8.1 kb	<i>CYP52A5A</i> isolated from the 1st library
pPA5	pTriplEx	<i>CYP52A5</i> <i>B</i>	4.1 kb	Approx. 7.8 kb	<i>CYP52A5B</i> isolated from the 2nd library
pPA15	pTriplEx	<i>CYP52A2</i> <i>A</i>	6.0 kb	Approx. 9.7 kb	<i>CYP52A2A</i> isolated from the 2nd library
pPA57	pTriplEx	<i>CYP52A3</i> <i>A</i>	5.5 kb	Approx. 9.2 kb	<i>CYP52A3A</i> isolated from the 2nd library
pPA62	pTriplEx	<i>CYP52A3</i> <i>B</i>	6.0 kb	Approx. 9.7 kb	<i>CYP52A3B</i> isolated from the 2nd library

EXAMPLE 1

Purification of Genomic DNA from *Candida tropicalis* ATCC 20336

A. Construction of Genomic Libraries

50 ml of YEPD broth (see Table 9) was inoculated with a single colony of *C. tropicalis* 20336 from YEPD agar plate and grown overnight at 30°C. 5 ml of the overnight culture was inoculated into 100 ml of fresh YEPD broth and incubated at 30°C for 4 to 5 hr with shaking. Cells were harvested by centrifugation, washed twice with sterile distilled water and resuspended in 4 ml of spheroplasting buffer (1 M Sorbitol, 50 mM EDTA, 14 mM mercaptoethanol) and incubated for 30 min at 37°C with gentle shaking. 0.5 ml of 2 mg/ml zymolyase (ICN Pharmaceuticals, Inc., Irvine, CA) was added and incubated at 37°C with gentle shaking for 30 to 60 min. Spheroplast formation was monitored by SDS lysis. Spheroplasts were harvested by brief centrifugation (4,000 rpm, 3 min) and were washed once with the spheroplast buffer without mercaptoethanol. Harvested spheroplasts were then suspended in 4 ml of lysis buffer (0.2 M Tris/pH 8.0, 50 mM EDTA, 1% SDS) containing 100 µg/ml RNase (Qiagen Inc., Chatsworth, CA) and incubated at 37°C for 30 to 60 min.

Proteins were denatured and extracted twice with an equal volume of chloroform/isoamyl alcohol (24:1) by gently mixing the two phases by hand inversions. The two phases were separated by centrifugation at 10,000 rpm for 10 min and the aqueous phase containing the high-molecular weight DNA was recovered. To the aqueous layer NaCl was added to a final concentration of 0.2 M and the DNA was

precipitated by adding 2 vol of ethanol. Precipitated DNA was spooled with a clean glass rod and resuspended in TE buffer (10 mM Tris/pH 8.0, 1 mM EDTA) and allowed to dissolve overnight at 4°C. To the dissolved DNA, RNase free of any DNase activity (Qiagen Inc., Chatsworth, CA) was added to a final concentration of 50 µg/ml and incubated at 37°C for 30 min. Then protease (Qiagen Inc., Chatsworth, CA) was added to a final concentration of 100 µg/ml and incubated at 55 to 60°C for 30 min. The solution was extracted once with an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1) and once with equal volume of chloroform/isoamyl alcohol (24:1). To the aqueous phase 0.1 vol of 3 M sodium acetate and 2 volumes of ice cold ethanol (200 proof) were added and the high molecular weight DNA was spooled with a glass rod and dissolved in 1 to 2 ml of TE buffer.

B. Genomic DNA Preparation for PCR Amplification of *CYP* and *CPR* Genes

Five 5 ml of YPD medium was inoculated with a single colony and grown at 30°C overnight. The culture was centrifuged for 5 min at 1200 x g. The supernatant was removed by aspiration and 0.5 ml of a sorbitol solution (0.9 M sorbitol, 0.1 M Tris-Cl pH 8.0, 0.1 M EDTA) was added to the pellet. The pellet was resuspended by vortexing and 1 µl of 2-mercaptoethanol and 50 µl of a 10 µg/ml zymolyase solution were added to the mixture. The tube was incubated at 37°C for 1 hr on a rotary shaker (200 rpm). The tube was then centrifuged for 5 min at 1200 x g and the supernatant was removed by aspiration. The protoplast pellet was resuspended in 0.5 ml 1x TE (10 mM Tris-Cl pH 8.0, 1 mM EDTA) and transferred to a 1.5 ml microcentrifuge tube. The protoplasts were lysed by the addition of 50 µl 10% SDS followed by incubation at 65°C for 20 min. Next, 200 µl of 5M potassium acetate was added and after mixing, the tube was incubated on ice for at least 30 min. Cellular debris was removed by centrifugation at 13,000 x g for 5 min. The supernatant was carefully removed and transferred to a new microfuge tube. The DNA was precipitated by the addition of 1 ml 100% (200 proof) ethanol followed by centrifugation for 5 min at 13,000 x g. The DNA pellet was washed with 1 ml 70 %

ethanol followed by centrifugation for 5 min at 13,000 x g. After partially drying the DNA under a vacuum, it was resuspended in 200 µl of 1x TE. The DNA concentration was determined by ratio of the absorbance at 260 nm / 280 nm ($A_{260/280}$).

EXAMPLE 2

Construction of *Candida tropicalis* 20336 Genomic Libraries

Three genomic libraries of *C. tropicalis* were constructed, two at Clontech Laboratories, Inc., (Palo Alto, CA) and one at Henkel Corporation (Cincinnati, OH).

A. Clontech Libraries

The first Clontech library was made as follows: Genomic DNA was prepared from *C. tropicalis* 20336 as described above, partially digested with *EcoRI* and size fractionated by gel electrophoresis to eliminate fragments smaller than 0.6 kb. Following size fractionation, several ligations of the *EcoRI* genomic DNA fragments and lambda (λ) TriplEx™ vector (Figure 1) arms with *EcoRI* sticky ends were packaged into λ phage heads under conditions designed to obtain one million independent clones. The second genomic library was constructed as follows: Genomic DNA was digested partially with *Sau3A1* and size fractionated by gel electrophoresis. The DNA fragments were blunt ended using standard protocols as described, e.g., in Sambrook et al, *Molecular Cloning: A Laboratory Manual*, 2ed. Cold Spring Harbor Press, USA (1989), incorporated herein by reference. The strategy was to fill in the *Sau3A1* overhangs with Klenow polymerase (Life Technologies, Grand Island, NY) followed by digestion with S1 nuclease (Life Technologies, Grand Island, NY). After S1 nuclease digestion the fragments were end filled one more time with Klenow polymerase to obtain the final blunt-ended DNA fragments. *EcoRI* linkers were ligated to these blunt-ended DNA fragments followed by ligation into the λ TriplEx vector. The resultant library contained approximately 2×10^6 independent clones with an average insert size of 4.5 kb.

B. Henkel Library

The third genomic library was constructed at Henkel Corporation using λ ZAP ExpressTM vector (Stratagene, La Jolla, CA) (Figure 2). Genomic DNA was partially digested with *Sau3A1* and fragments in the range of 6 to 12 kb were purified from an agarose gel after electrophoresis of the digested DNA. These DNA fragments were then ligated to *Bam*HI digested λ ZAP ExpressTM vector arms according to manufacturers protocols. Three ligations were set up to obtain approximately 9.8×10^5 independent clones. All three libraries were pooled and amplified according to manufacturer instructions to obtain high-titre ($>10^9$ plaque forming units/ml) stock for long-term storage. The titre of packaged phage library was ascertained after infection of *E. coli* XL1Blue-MRF' cells. *E. coli* XL1Blue-MRF' were grown overnight in either in LB medium or NZCYM (Table 9) containing 10 mM MgSO₄ and 0.2% maltose at 37°C or 30°C, respectively with shaking. Cells were then centrifuged and resuspended in 0.5 to 1 volume of 10 mM MgSO₄. 200 μ l of this *E. coli* culture was mixed with several dilutions of packaged phage library and incubated at 37°C for 15 min. To this mixture 2.5 ml of LB top agarose or NZCYM top agarose (maintained at 60°C) (see Table 9) was added and plated on LB agar or NCZYM agar (see Table 9) present in 82 mm petri dishes. Phage were allowed to propagate overnight at 37°C to obtain discrete plaques and the phage titre was determined.

EXAMPLE 3

Screening of Genomic Libraries

Both λ TriplExTM and λ ZAP ExpressTM vectors are phagemid vectors that can be propagated either as phage or plasmid DNA (after conversion of phage to plasmid). Therefore, the genomic libraries constructed in these vectors can be screened either by plaque hybridization (screening of lambda form of library) or by colony hybridization (screening plasmid form of library after phage to plasmid conversion). Both vectors are capable of expressing the cloned genes and the main difference is the

mechanism of excision of plasmid from the phage DNA. The cloning site in λ TriplExTM is located within a plasmid which is present in the phage and is flanked by *loxP* site (Figure 1). When λ TriplExTM is introduced into *E. coli* strain BM25.8 (supplied by Clontech), the *Cre* recombinase present in BM25.8 promotes the excision and circularization of plasmid pTriplEx from the phage λ TriplExTM at the *loxP* sites. The mechanism of excision of plasmid pBK-CMV from phage λ ZAP ExpressTM is different. It requires the assistance of a helper phage such as ExAssistTM (Stratagene) and an *E. coli* strain such as XLOR (Stratagene). Both pTriplEx and pBK-CMV can replicate autonomously in *E. coli*.

A. Screening Genomic Libraries (Plasmid Form)

1) Colony Lifts

A single colony of *E. coli* BM25.8 was inoculated into 5 ml of LB containing 50 μ g/ml kanamycin, 10 mM MgSO₄ and 0.1% maltose and grown overnight at 31°C, 250 rpm. To 200 μ l of this overnight culture ($\sim 4 \times 10^8$ cells) 1 μ l of phage library ($2 - 5 \times 10^6$ plaque forming units) and 150 μ l LB broth were added and incubated at 31°C for 30 min after which 400 μ l of LB broth was added and incubated at 31°C, 225 rpm for 1 h. This bacterial culture was diluted and plated on LB agar containing 50 μ g/ml ampicillin (Sigma Chemical Company, St. Louis, MO) and kanamycin (Sigma Chemical Company) to obtain 500 to 600 colonies/plate. The plates were incubated at 37°C for 6 to 7 hrs until the colonies became visible. The plates were then stored at 4°C for 1.5 h before placing a Colony/Plaque ScreenTM Hybridization Transfer Membrane disc (DuPont NEN Research Products, Boston, MA) on the plate in contact with bacterial colonies. The transfer of colonies to the membrane was allowed to proceed for 3 to 5 min. The membrane was then lifted and placed on a fresh LB agar (see Table 9) plate containing 200 μ g/ml of chloramphenicol with the side exposed to the bacterial colonies facing up. The plates containing the membranes were then incubated at 37°C overnight in order to allow full development of the bacterial colonies. The LB agar plates from

which colonies were initially lifted were incubated at 37°C overnight and stored at 4°C for future use. The following morning the membranes containing bacterial colonies were lifted and placed on two sheets of Whatman 3M (Whatman, Hillsboro, OR) paper saturated with 0.5 N NaOH and left at room temperature (RT) for 3 to 6 min to lyse the cells. Additional treatment of membranes was as described in the protocol provided by NEN Research Products.

2) DNA Hybridizations

Membranes were dried overnight before hybridizing to oligonucleotide probes prepared using a non-radioactive ECL™ 3'-oligolabelling and detection system from Amersham Life Sciences (Arlington Heights, IL). DNA labeling, prehybridization and hybridizations were performed according to manufacturer's protocols. After hybridization, membranes were washed twice at room temperature in 5 X SSC, 0.1% SDS (in a volume equivalent to 2 ml/cm² of membrane) for 5 min each followed by two washes at 50°C in 1X SSC, 0.1% SDS (in a volume equivalent to 2 ml/cm² of membrane) for 15 min each. The hybridization signal was then generated and detected with Hyperfilm ECL™ (Amersham) according to manufacturer's protocols. Membranes were aligned to plates containing bacterial colonies from which colony lifts were performed and colonies corresponding to positive signals on X-ray were then isolated and propagated in LB broth. Plasmid DNA's were isolated from these cultures and analyzed by restriction enzyme digestions and by DNA sequencing.

B. Screening Genomic Libraries (Plaque Form)

1) λ Library Plating

E. coli XL1Blue-MRF' cells were grown overnight in LB medium (25 ml) containing 10 mM MgSO₄ and 0.2% maltose at 37°C, 250 rpm. Cells were then centrifuged (2,200 x g for 10 min) and resuspended in 0.5 volumes of 10 mM MgSO₄. 500 µl of this *E. coli* culture was mixed with a phage suspension containing 25,000

amplified lambda phage particles and incubated at 37°C for 15 min. To this mixture 6.5 ml of NZCYM top agarose (maintained at 60°C) (see Chart) was added and plated on 80 - 100 ml NCZYM agar (see Chart) present in a 150 mm petridish. Phage were allowed to propagate overnight at 37°C to obtain discrete plaques. After overnight growth plates were stored in a refrigerator for 1-2 hr before plaque lifts were performed.

2) Plaque Lift and DNA Hybridizations

Magna Lift™ nylon membranes (Micron Separations, Inc., Westborough, MA) were placed on the agar surface in complete contact with λ plaques and transfer of plaques to nylon membranes was allowed to proceed for 5 min at RT. After plaque transfer the membrane was placed on 2 sheets of Whatman 3M™ (Whatman, Hillsboro, OR) filter paper saturated with a 0.5 N NaOH, 1.0 M NaCl solution and left for 10 min at RT to denature DNA. Excess denaturing solution was removed by blotting briefly on dry Whatman 3M paper. Membranes were then transferred to 2 sheets of Whatman 3M™ paper saturated with 0.5 M Tris-HCl (pH 8.0), 1.5 M NaCl and left for 5 min to neutralize. Membranes were then briefly washed in 200 - 500 ml of 2 X SSC, dried by air and baked for 30 - 40 min at 80°C. The membranes were then probed with labelled DNA.

Membranes were prewashed with a 200 - 500 ml solution of 5 X SSC, 0.5% SDS, 1 mM EDTA (pH 8.0) for 1 - 2 hr at 42°C with shaking (60 rpm) to get rid of bacterial debris from the membranes. The membranes were prehybridized for 1 - 2 hr at 42°C with (in a volume equivalent to 0.125 - 0.25 ml/cm² of membrane) ECL Gold™ buffer (Amersham) containing 0.5 M NaCl and 5% blocking reagent. DNA fragments that were used as probes were purified from agarose gel using a QIAEX II™ gel extraction kit (Qiagen Inc., Chatsworth, CA) according to manufacturers protocol and labeled using an Amersham ECL™ direct nucleic acid labeling kit (Amersham). Labeled DNA (5 - 10 ng/ml hybridization solution) was added to the prehybridized membranes and the hybridization was allowed to proceed overnight. The following day membranes

were washed with shaking (60 rpm) twice at 42°C for 20 min each time in (in a volume equivalent to 2 ml/cm² of membrane) a buffer containing either 0.1 (high stringency) or 0.5 (low stringency) X SSC, 0.4% SDS and 360 g/l urea. This was followed by two 5 min washes at room temperature in (in a volume equivalent to 2 ml/cm² of membrane) 2 X SSC. Hybridization signals were generated using the ECLTM nucleic acid detection reagent and detected using Hyperfilm ECLTM (Amersham).

Agar plugs which contained plaques corresponding to positive signals on the X-ray film were taken from the master plates using the broad-end of Pasteur pipet. Plaques were selected by aligning the plates with the x-ray film. At this stage, multiple plaques were generally taken. Phage particles were eluted from the agar plugs by soaking in 1 ml SM buffer (Sambrook et al., *supra*) overnight. The phage eluate was then diluted and plated with freshly grown *E. coli* XL1Blue-MRF' cells to obtain 100 - 500 plaques per 85 mm NCZYM agar plate. Plaques were transferred to Magna Lift nylon membranes as before and probed again using the same probe. Single well-isolated plaques corresponding to signals on X - ray film were picked by removing agar plugs and eluting the phage by soaking overnight in 0.5 ml SM buffer.

C. Conversion of λ Clones to Plasmid Form

The lambda clones isolated were converted to plasmid form for further analysis. Conversion from the plaque to the plasmid form was accomplished by infecting the plaques into *E. coli* strain BM25.8. The *E. coli* strain was grown overnight at 31°C, 250 rpm in LB broth containing 10 mM MgSO₄ and 0.2% maltose until the OD₆₀₀ reached 1.1 - 1.4. Ten milliliters of the overnight culture was removed and mixed with 100 μ l of 1 M MgCl₂. A 200 μ l volume of cells was removed, mixed with 150 μ l of eluted phage suspension and incubated at 31°C for 30 min. LB broth (400 μ l) was added to the tube and incubation was continued at 31°C for 1 hr with shaking, 250 rpm. 1 - 10 μ l of the infected cell suspension was plated on LB agar containing 100 μ g/ml ampicillin (Sigma, St. Louis, MO). Well-isolated colonies were picked and grown overnight in 5 ml LB

broth containing 100 µg/ml ampicillin at 37°C, 250 rpm. Plasmid DNA was isolated from these cultures and analyzed. To convert the λZAP Express™ vector to plasmid form *E. coli* strains XL1Blue-MRF' and XLOR were used. The conversion was performed according to the manufacturer's (Stratagene) protocols for single-plaque excision.

EXAMPLE 4

Transformation of *C. tropicalis* H5343 *ura*

A. Transformation of *C. tropicalis* H5343 by Electroporation

5 ml of YEPD was inoculated with *C. tropicalis* H5343 *ura* from a frozen stock and incubated overnight on a New Brunswick shaker at 30°C and 170 rpm. The next day, 10 µl of the overnight culture was inoculated into 100 ml YEPD and growth was continued at 30°C, 170 rpm. The following day the cells were harvested at an OD₆₀₀ of 1.0 and the cell pellet was washed one time with sterile ice-cold water. The cells were resuspended in ice-cold sterile 35 % Polyethylene glycol (4,000 MW) to a density of 5x10⁸ cells/ml. A 0.1 ml volume of cells were utilized for each electroporation. The following electroporation protocol was followed: 1.0 µg of transforming DNA was added to 0.1 ml cells, along with 5 µg denatured, sheared calf thymus DNA and the mixture was allowed to incubate on ice for 15 min. The cell solution was then transferred to an ice-cold 0.2 cm electroporation cuvette, tapped to make sure the solution was on the bottom of the cuvette and electroporated. The cells were electroporated using an Invitrogen electroporator (Carlsbad, CA) at 450 Volts, 200 Ohms and 250 µF. Following electroporation, 0.9 ml SOS media (1M Sorbitol, 30% YEPD, 10 mM CaCl₂) was added to the suspension. The resulting culture was grown for 1 hr at 30°C, 170 rpm. Following the incubation, the cells were pelleted by centrifugation at 1500 x g for 5 min. The electroporated cells were resuspended in 0.2 ml of 1M sorbitol and plated on synthetic complete media minus uracil (SC - uracil) (Nelson, *supra*). In some cases the electroporated cells were plated directly onto SC - uracil. Growth of transformants was monitored for 5 days. After three days,

several transformants were picked and transferred to SC-uracil plates for genomic DNA preparation and screening.

B. Transformation of *C. tropicalis* Using Lithium Acetate

The following protocol was used to transform *C. tropicalis* in accordance with the procedures described in *Current Protocols in Molecular Biology*, Supplement 5, 13.7.1 (1989), incorporated herein by reference.

5 ml of YEPD was inoculated with *C. tropicalis* H5343 *ura⁻* from a frozen stock and incubated overnight on a New Brunswick shaker at 30°C and 170 rpm. The next day, 10 µl of the overnight culture was inoculated into 50 ml YEPD and growth was continued at 30°C, 170 rpm. The following day the cells were harvested at an OD₆₀₀ of 1.0. The culture was transferred to a 50 ml polypropylene tube and centrifuged at 1000 X g for 10 min. The cell pellet was resuspended in 10 ml sterile TE (10mM Tris-Cl and 1mM EDTA, pH 8.0). The cells were again centrifuged at 1000 X g for 10 min and the cell pellet was resuspended in 10 ml of a sterile lithium acetate solution [LiAc (0.1 M lithium acetate, 10 mM Tris-Cl, pH 8.0, 1 mM EDTA)]. Following centrifugation at 1000 X g for 10 min., the pellet was resuspended in 0.5 ml LiAc. This solution was incubated for one hour at 30°C while shaking gently at 50 rpm. A 0.1 ml aliquot of this suspension was incubated with 5 µg of transforming DNA at 30°C with no shaking for 30 min. A 0.7 ml PEG solution (40 % wt/vol polyethylene glycol 3340, 0.1 M lithium acetate, 10 mM Tris-Cl, pH 8.0, 1 mM EDTA) was added and incubated at 30°C for 45 min. The tubes were then placed at 42°C for 5 min. A 0.2 ml aliquot was plated on synthetic complete media minus uracil (SC - uracil) (Kaiser et al. *Methods in Yeast Genetics*, Cold Spring Harbor Laboratory Press, USA, 1994, incorporated herein by reference). Growth of transformants was monitored for 5 days. After three days, several transformants were picked and transferred to SC-uracil plates for genomic DNA preparation and screening.

EXAMPLE 5

Plasmid DNA Isolation

Plasmid DNA were isolated from *E. coli* cultures using Qiagen plasmid isolation kit (Qiagen Inc., Chatsworth, CA) according to manufacturer's instructions.

EXAMPLE 6

DNA Sequencing and Analysis

DNA sequencing was performed at Sequetech Corporation (Mountain View, CA) using Applied Biosystems automated sequencer (Perkin Elmer, Foster City, CA). DNA sequences were analyzed with MacVector and GeneWorks software packages (Oxford Molecular Group, Campbell, CA).

EXAMPLE 7

PCR Protocols

PCR amplification was carried out in a Perkin Elmer Thermocycler using the AmpliTaqGold enzyme (Perkin Elmer Cetus, Foster City, CA) kit according to manufacturer's specifications. Following successful amplification, in some cases, the products were digested with the appropriate enzymes and gel purified using QiaexII (Qiagen, Chatsworth, CA) as per manufacturer instructions. In specific cases the Ultima Taq polymerase (Perkin Elmer Cetus, Foster City, CA) or the Expand Hi-Fi Taq

polymerase (Boehringer Mannheim, Indianapolis, IN) were used per manufacturer's recommendations or as defined in Table 3.

Table 3. PCR amplification conditions used with different primer combinations.

PRIMER COMBINATION	<i>Taq</i>	TEMPLATE DENATURING CONDITION	ANNEALING TEMP/TIME	EXTENSION TEMP/TIME	CYCLE Number
3674-41-1/ 41-2/ 41-4 + 3674-41-4	Ampli- <i>Taq</i> Gold	94 C/30 sec	55 C/30 sec	72 C/1 min	30
URA Primer 1a URA Primer 1b	Ampli- <i>Taq</i> Gold	95 C/1 min	70 C/1 min	72 C/2 min	35
URA Primer 2a URA Primer 2b	Ampli- <i>Taq</i> Gold	95 C/1 min	70 C/1 min	72 C/2 min	35
<i>CYP2A</i> #1 <i>CYP2A</i> #2	Ampli- <i>Taq</i> Gold	95 C/1 min	70 C/1 min	72 C/2 min	35
<i>CYP3A</i> #1 <i>CYP3A</i> #2	Ultma <i>Taq</i>	95 C/1 min	70 C/1 min	72 C/1 min	30
<i>CPR B</i> #1 <i>CPR B</i> #2	Expand Hi-Fi <i>Taq</i>	94 C/15 sec 94 C/15 sec	50 C/30 sec 50 C/30 sec	68 C/3 min 68 C/3 min +20 sec/cycle	10 15
<i>CYP5A</i> #1 <i>CYP5A</i> #2	Expand Hi-Fi <i>Taq</i>	94 C/15 sec 94 C/15 sec	50 C/30 sec 50 C/30 sec	68 C/3 min 68 C/3 min +20 sec/cycle	10 15

Table 4 below contains a list of primers (SEQ ID NOS: 1-35) used for PCR amplification to construct gene integration vectors or to generate probes for gene detection and isolation.

Table 4. Primer table for PCR amplification to construct gene integration vectors, to generate probes for gene isolation and detection and to obtain DNA sequence of constructs. (A- deoxyadenosine triphosphate [dATP], G- deoxyguanosine triphosphate [dGTP], C- deoxycytosine triphosphate [dCTP], T- deoxythymidine triphosphate [dTTP], Y- dCTP or dTTP, R- dATP or dGTP, W- dATP or dTTP, M- dATP or dCTP, N- dATP or dCTP or dGTP or dTTP).

Target gene(s)	Patent Primer Name	Lab Primer Name	Sequence (5' to 3')	PCR Product Size
<i>CYP52A2</i> A	CYP2A#1	3659-72M	<i>CCTTAATTAAATGCACGAAGCGGAGATAAAAG</i> (SEQ ID NO: 1)	2230 bp
	CYP2A#2	3659-72N	<i>CCTTAATTAAAGCATAAGCTTGCTCGAGTCT</i> (SEQ ID NO: 2)	
<i>CYP52A3</i> A	CYP3A#1	3659-72O	<i>CCTTAATTAAACGCAATGGGAACATGGAGTG</i> (SEQ ID NO: 3)	2154 bp
	CYP3A#2	3659-72P	<i>CCTTAATTAAATCGCACTACGGTTATTGGTATCAG</i> (SEQ ID NO: 4)	
<i>CYP52A5</i> A	CYP5A#1	3659-72K	<i>CCTTAATTAAATCAAAGTACGTTTCAGGCGG</i> (SEQ ID NO: 5)	3298 bp
	CYP5A#2	3659-72L	<i>CCTTAATTAAAGGCAGACAACAACCTGGCAAAGTC</i> (SEQ ID NO: 6)	
<i>CPRB</i>	CPRB#1	3698-20A	<i>CCTTAATTAAAGAGGTCGTTGGTTGAGTTTTTC</i> (SEQ ID NO: 7)	3266 bp
	CPRB#2	3698-20B	<i>CCTTAATTAAATTGATAATGACGTTGCGGG</i> (SEQ ID NO: 8)	
<i>URA3A</i>	URA Primer 1a	3698-7C	<i>AGGCGCGCGGAGTCCAAAAAGACCAACCTCTG</i> (SEQ ID NO: 9)	956 bp
	URA Primer 1b	3698-7D	<i>CCTTAATTAAATACGTGGATACCTTCAAGCAAGTG</i> (SEQ ID NO: 10)	
<i>URA3A</i>	URA Primer 2a	3698-7A	<i>CCTTAATTAAAGCTCACGAGTTTTGGGATTTTCGA</i> G (SEQ ID NO: 11)	750 bp
	URA Primer 2b	3698-7B	<i>GGGTTTAAACCGCAGAGGTGGTCTTTTTGGAC</i> TC (SEQ ID NO: 12)	
			<i>GGGTTTAAAC</i> - <i>Pme</i> I restriction site (SEQ ID NO: 13)	

			AGGCGCGCC - <i>AscI</i> restriction site (SEQ ID NO: 14)	
			CCTTAATTAA - <i>PacI</i> restriction site (SEQ ID NO: 15)	
<i>CPR</i>	FMN1	3674-41-1	TCYCAAACWGGTACWGCWGAA (SEQ ID NO: 16)	
<i>CPR</i>	FMN2	3674-41-2	GGTTTGGGTAAATCWACTTAT (SEQ ID NO: 17)	
<i>CPR</i>	FAD	3674-41-3	CGTTATTAYTCYATTTCTTC (SEQ ID NO: 18)	
<i>CPR</i>	NADPH	3674-41-4	GCMACACCRGTACCTGGACC (SEQ ID NO: 19)	
<i>CPR</i>	PRK1.F3	PRK1.F3	ATCCCAATCGTAATCAGC (SEQ ID NO: 20)	
<i>CPR</i>	PRK1.F5	PRK1.F5	ACTTGTCTTCGTTTAGCA (SEQ ID NO: 21)	
<i>CPR</i>	PRK4.R20	PRK4.R20	CTACGTCTGTGGTGATGC (SEQ ID NO: 22)	
<i>CYP</i>	UCup1	UCup1	CGNGAYACNACNGCNGG (SEQ ID NO: 23)	
<i>CYP</i>	UCup2	UCup2	AGRGAYACNACNGCNGG (SEQ ID NO: 24)	
<i>CYP</i>	UCdown1	UCdown1	AGNGCRAAYTGYTGNCC (SEQ ID NO: 25)	
<i>CYP</i>	UCdown2	UCdown2	YAANGCRAAYTGYTGNCC (SEQ ID NO: 26)	
<i>CYP</i>	HemeB1	HemeB1	ATTCAACGGTGGTCCAAGAATCTGTTTGG (SEQ ID NO: 27)	
<i>CYP</i>	2,3,5P	2,3,5P	GAGCTATGTTGAGACCACAGTTTGC (SEQ ID NO: 28)	
<i>CYP</i>	2,3,5M	2,3,5M	CTTCAGTTAAAGCAAATTGTTTGGCC (SEQ ID NO: 29)	
pTriplEx vector	Triplex5'	Triplex5'	CTCGGGAAGCGCGCCATTGTGTGG (SEQ ID NO: 30)	
pTriplEx vector	Triplex3'	Triplex3'	TAATACGACTCACTATAGGGCGAATTGGC (SEQ ID NO: 31)	
<i>CYP</i>	Cyp52a	Cyp52a	TGRYTCAAACCATCTYTCTGG (SEQ ID NO: 32)	
<i>CYP</i>	Cyp52b	Cyp52b	GGACCGGCGTTAAAGGG (SEQ ID NO: 33)	
<i>CYP</i>	Cyp52c	Cyp52c	CATAGTCGWATYATGCTTAGACC (SEQ ID NO: 34)	
<i>CYP</i>	Cyp52d	Cyp52d	GGACCACCATTGAATGG (SEQ ID NO: 35)	

EXAMPLE 8

Yeast Colony PCR Procedure for Confirmation of Gene Integration into the Genome of *C. tropicalis*

Single yeast colonies were removed from the surface of transformation plates, suspended in 50 μ l of spheroplasting buffer (50mM KCl, 10mM Tris-HCl, pH 8.3, 1.0 mg/ml Zymolyase, 5% glycerol) and incubated at 37°C for 30 min. Following incubation, the solution was heated for 10 min at 95°C to lyse the cells. Five μ l of this solution was used as a template in PCR. Expand Hi-Fi *Taq* polymerase (Boehringer Mannheim, Indianapolis, IN) was used in PCR coupled with a gene-specific primer (gene to be integrated) and a *URA3* primer. If integration did occur, amplification would yield a PCR product of predicted size confirming the presence of an integrated gene.

EXAMPLE 9

Fermentation Method for Gene Induction Studies

A fermentor was charged with a semi-synthetic growth medium having the composition 75 g/l glucose (anhydrous), 6.7 g/l Yeast Nitrogen Base (Difco Laboratories), 3 g/l yeast extract, 3 g/l ammonium sulfate, 2 g/l monopotassium phosphate, 0.5 g/l sodium chloride. Components were made as concentrated solutions for autoclaving then added to the fermentor upon cooling: final pH approximately 5.2. This charge was inoculated with 5-10% of an overnight culture of *C. tropicalis* ATCC 20962 prepared in YM medium (Difco Laboratories) as described in the methods of Examples 17 and 20 of US Patent 5,254,466, which is incorporated herein by reference. *C. tropicalis* ATCC 20962 is a POX 4 and POX 5 disrupted *C. tropicalis* ATCC 20336. Air and agitation were supplied to maintain the dissolved oxygen at greater than about 40% of saturation versus air. The pH was maintained at about 5.0 to 8.5 by the addition of 5N caustic soda on pH control. Both a fatty acid feedstream (commercial oleic acid in this example) having a typical composition: 2.4% C₁₄; 0.7% C_{14:1}; 4.6% C₁₆; 5.7% C_{16:1}; 5.7% C_{17:1}; 1.0% C₁₈; 69.9% C_{18:1}; 8.8% C_{18:2}; 0.30% C_{18:3}; 0.90% C_{20:1} and a glucose co-substrate feed were added in a feedbatch mode beginning near the end of exponential growth. Caustic was added on pH control during

hundred microliters of each ethanol treated sample were transferred to a Qiagen RNeasy spin column, followed by centrifugation at 8,000 x g for 15 sec. The flow through was discarded and the column reloaded with the remaining sample (700 µl) and re-centrifuged at 8,000 x g for 15 sec. The column was washed once with 700 µl of buffer RW1*, and centrifuged at 8,000 x g for 15 sec and the flow through discarded. The column was placed in a new 2 ml collection tube and washed with 500 µl of RPE* buffer and the flow through discarded. The RPE* wash was repeated with centrifugation at 8,000 x g for 2 min and the flow through discarded. The spin column was transferred to a new 1.5 ml collection tube and 100 µl of RNase free water added to the column followed by centrifugation at 8,000 x g for 15 seconds. An additional 75 µl of RNase free water was added to the column followed by centrifugation at 8,000 x g for 2 min. RNA eluted in the water flow through was collected for further purification.

The RNA eluate was then treated to remove contaminating DNA. Twenty microliters of 10X DNase I buffer (0.5 M tris (pH 7.5), 50 mM CaCl₂, 100 mM MgCl₂), 10 µl of RNase-free DNase I (2 Units/µl, Ambion Inc., Austin, Texas) and 40 units Rnasin (Promega Corporation, Madison, Wisconsin) were added to the RNA sample. The mixture was then incubated at 37°C for 15 to 30 min. Samples were placed on ice and 250 µl Lysis buffer RLT* and 250 µl ethanol (200 proof) added. The samples were then mixed by inversion. The samples were transferred to Qiagen RNeasy spin columns and centrifuged at 8,000 x g for 15 sec and the flow through discarded. Columns were placed in new 2 ml collection tubes and washed twice with 500 µl of RPE* wash buffer and the flow through discarded. Columns were transferred to new 1.5 ml eppendorf tubes and RNA was eluted by the addition of 100 µl of DEPC treated water followed by centrifugation at 8,000 x g for 15 sec. Residual RNA was collected by adding an additional 50 µl of RNase free water to the spin column followed by centrifugation at full speed for 2 min. 10 µl of the RNA preparation was removed and quantified by the (A₂₆₀ 280) method. RNA was stored at -70°C. Yields were found to be 30-100 µg total RNA per 2.0 ml of fermentation broth.

the bioconversion of fatty acids to diacids to maintain the pH in the desired range. Typically, samples for gene induction studies were collected just prior to starting the fatty acid feed and over the first 10 hours of bioconversion. Determination of fatty acid and diacid content was determined by a standard methyl ester protocol using gas liquid chromatography (GLC). Gene induction was measured using the QC-RT-PCR protocol described in this application.

EXAMPLE 10

RNA Preparation

The first step of this protocol involves the isolation of total cellular RNA from cultures of *C. tropicalis*. The cellular RNA was isolated using the Qiagen RNeasy Mini Kit (Qiagen Inc., Chatsworth, CA) as follows: 2 ml samples of *C. tropicalis* cultures were collected from the fermentor in a standard 2 ml screw capped Eppendorf style tubes at various times before and after the addition of the fatty acid or alkane substrate. Cell samples were immediately frozen in liquid nitrogen or a dry-ice/alcohol bath after their harvesting from the fermentor. To isolate total RNA from the samples, the tubes were allowed to thaw on ice and the cells pelleted by centrifugation in a microfuge for 5 minutes (min) at 4°C and the supernatant was discarded while keeping the pellet ice-cold.

The microfuge tubes were filled 2/3 full with ice-cold Zirconia/Silica beads (0.5 mm diameter, Biospec Products, Bartlesville, OK) and the tube filled to the top with ice-cold RLT* lysis buffer (* buffer included with the Qiagen RNeasy Mini Kit). Cell rupture was achieved by placing the samples in a mini bead beater (Biospec Products, Bartlesville, OK) and immediately homogenized at full speed for 2.5 min. The samples were allowed to cool in a ice water bath for 1 minute and the homogenization/cool process repeated two more times for a total of 7.5 min homogenization time in the beadbeater. The homogenized cells samples were microfuged at full speed for 10 min and 700 µl of the RNA containing supernatant removed and transferred to a new eppendorf tube. 700 µl of 70% ethanol was added to each sample followed by mixing by inversion. This and all subsequent steps were performed at room temperature. Seven

EXAMPLE 11

Quantitative Competitive Reverse Transcription Polymerase Chain Reaction (QC-RT-PCR) Protocol

QC-RT-PCR is a technique used to quantitate the amount of a specific RNA in a RNA sample. This technique employs the synthesis of a specific DNA molecule that is complementary to an RNA molecule in the original sample by reverse transcription and its subsequent amplification by polymerase chain reaction. By the addition of various amounts of a competitor RNA molecule to the sample one can determine the concentration of the RNA molecule of interest (in this case the mRNA transcripts of the *CYP* and *CPR* genes). The levels of specific mRNA transcripts were assayed over time in response to the addition of fatty acid and/or alkane substrates to the growth medium of fermentation grown *C. tropicalis* cultures for the identification and characterization of the genes involved in the oxidation of these substrates. This approach can be used to identify the *CYP* and *CPR* genes involved in the oxidation of any given substrate based upon their transcriptional regulation.

A. Primer Design

The first requirement for QC-RT-PCR is the design of the primer pairs to be used in the reverse transcription and subsequent PCR reactions. These primers need to be unique and specific to the gene of interest. As there is a family of genetically similar *CYP* genes present in *C. tropicalis* 20336, care had to be taken to design primer pairs that would be discriminating and only amplify the gene of interest, in this example the *CYP52A5* gene. In this manner, unique primers directed to substantially non-homologous (aka variable) regions within target members of a gene family are constructed. What constitutes substantially non-homologous regions is determined on a case by case basis. Such unique primers should be specific enough to anneal the non-homologous region of the target gene without annealing to other non-target members of the gene family. By

comparing the known sequences of the members of a gene family, non-homologous regions are identified and unique primers are constructed which will anneal to those regions. It is contemplated that non-homologous regions herein would typically exhibit less than about 85% homology but can be more homologous depending on the positions which are conserved and stringency of the reaction. After conducting PCR, it may be helpful to check the reaction product to assure it represents the unique target gene product. If not, the reaction conditions can be altered in terms of stringency to focus the reaction to the desired target. Alternatively a new primer or new non-homologous region can be chosen. Due to the high level of homology between the genes of the *CYP52A* family, the most variable 5 prime region of the *CYP52A5* coding sequence was targeted for the design of the primer pairs. In Figure 3, a portion of the 5 prime coding region for the *CYP52A5A* (SEQ ID NO: 36) allele of *C. tropicalis* 20336 is shown. The boxed sequences in Figure 3 are the sequences of the forward and backwards primers (SEQ ID NOS: 47 and 48) used to quantitate expression of both alleles of this gene. The actual reverse primer (SEQ ID NO: 48) contains one less adenine than that shown in Figure 3. Primers used to measure the expression of specific *C. tropicalis* 20336 genes using the QC-RT-PCR protocol are listed in Table 5 (SEQ ID NOS: 37-58).

Table 5. Primer used to measure *C. tropicalis* gene expression in the QC-RT-PCR reactions.

Primer Name	Direction	Target	Sequence
3737-89F	F	<i>CYP52A1A</i>	CCGATGAAGTTTTTCGACGAGTACCC (SEQ ID NO: 37)
3737-89B	B	<i>CYP52A1A</i>	AAGGCTTTAACGTGTCCAATCTGGTC (SEQ ID NO: 38)
alk2aF1	F	<i>CYP52A2A</i>	ATTATCGCCACATACTTCACCAAATGG (SEQ ID NO: 39)
alk2aB5	B	<i>CYP52A2A</i>	CGAGATCGTGGATACGCTGGAGTG (SEQ ID NO: 40)
7581-178-3	F	<i>CYP52A3A</i>	GCCACTCGGTAACCTTTGTCAGGGAC (SEQ ID NO: 41)
7581-178-4	B	<i>CYP52A3A</i>	CATTGAACTGAGTAGCCAAAACAGCC (SEQ ID NO: 42)
3737-50F	F	<i>CYP52A3A</i> & <i>CYP52A3B</i>	CCTACGTTTGGTATCGCTACTCCGTTG (SEQ ID NO: 43)
3737-50B	B	<i>CYP52A3A</i> & <i>CYP52A3B</i>	TTTCCAGCCAGCACCGTCCAAG (SEQ ID NO: 44)
3737-175F	F	<i>CYP52D4A</i>	GCAGAGCCGATCTATGTTGCGTCC (SEQ ID NO: 45)
3737-175B	B	<i>CYP52D4A</i>	TCATTGAATGCTTCCAGGAACCTCG (SEQ ID NO: 46)
7581-97-F	F	<i>CYP52A5A</i> & <i>CYP52A5B</i>	AAGAGGGCAGGGCTCAAGAG (SEQ ID NO: 47)
7581-97-M	B	<i>CYP52A5A</i> & <i>CYP52A5B</i>	TCCATGTGAAGATCCCATCAC (SEQ ID NO: 48)
4P-2	F	<i>CYP52A8A</i>	CTTGAAGGCCGTGTTGAACG (SEQ ID NO: 49)
4M-1	B	<i>CYP52A8A</i>	CAGGATTTGTCTGAGTTGCCG (SEQ ID NO: 50)
3737-52F	F	<i>POX4A</i> & <i>POX4B</i>	CCATTGCCTTGAGATACGCCATTGGTAG (SEQ ID NO: 51)
3737-52B	B	<i>POX4A</i> & <i>POX4B</i>	AGCCTTGGTGTCGTTCTTTTCAACGG (SEQ ID NO: 52)
3737-53F	F	<i>POX5A</i>	TTGGGTTTGTGTTTCTCCTGTGTCCG (SEQ ID NO: 53)
3737-53B	B	<i>POX5A</i>	CCTTTGACCTTCAATCTGGCGTAGACG (SEQ ID NO: 54)
F33	F	<i>CPRA</i>	GGTTTGCTGAATACGCTGAAGGTGATG (SEQ ID NO: 55)
B63	B	<i>CPRA</i>	TGGAGCTGAACAACTCTCTCGTCTCGG (SEQ ID NO: 56)
3737-133F	F	<i>CPRA</i> & <i>CPRB</i>	TTCCTCAACACGGACAGCGG (SEQ ID NO: 57)

3737-133B	B	CPRA & CPRB	AGTCAACCAGGTGTGGAACTCGTC (SEQ ID NO: 58)
-----------	---	----------------	---

F-Forward B-Backward

B. Design and Synthesis of the Competitor DNA Template

The competitor RNA is synthesized *in vitro* from a competitor DNA template that has the T7 polymerase promoter and preferably carries a small deletion of e.g., about 10 to 25 nucleotides relative to the native target RNA sequence. The DNA template for the *in-vitro* synthesis of the competitor RNA is synthesized using PCR primers that are between 46 and 60 nucleotides in length. In this example, the primer pairs for the synthesis of the *CYP52A5* competitor DNA are shown in Tables 6 and 7 (SEQ ID NOS: 59 AND 60).

Table 6. Forward and Reverse primers used to synthesize the competitor RNA template for the QC-RT-PCR measurement of *CYP52A5A* gene expression.

Forward Primer	<i>CYP52A5A</i>	GGATCCTAATACGACTCACTATAGGGAGG AAGAGGGCAGGGCTCAAGAG (SEQ ID NO: 59)
Reverse Primer	<i>CYP52A5A</i>	TCCATGTGAAGATCCCATCACGAGTGTGC CTCTTGCCCAAAG (SEQ ID NO: 60)

Table 7. Primers for the synthesis of the QC-RT-PCR competitor RNA templates

Primer Name	Direction	Target	Sequence 5'-3'
3737-89C	F	<i>CYP52A1A</i>	GGATCCTAATACGACTCACTATAGGGAGGCCGAT GAAGTTTTCGACGAGTACCC (SEQ ID NO: 61)
3737-89D	B	<i>CYP52A1A</i>	AAGGCTTTAACGTGTCCAATCTGGTC AACATAGCTCTGGAGTGCTTCCAACC (SEQ ID NO: 62)
7581-137-A	F	<i>CYP52A2A</i>	GGATCCTAATACGACTCACTATAGGGAGGATTAT CGCCACATACTCACCAAATGG (SEQ ID NO: 63)

7581-137-B	B	<i>CYP52A2A</i>	CGAGATCGTGGATACGCTGGAGTGCCTCGCTCTT CTTCTTCAACAATTCAAG (SEQ ID NO: 64)
7581-137-D	B	<i>CYP52A3A</i>	CATTGAACTGAGTAGCCAAAACAGCCCATGGTTT CAATCAATGGGAGGC (SEQ ID NO: 65)
7581-137-C	F	<i>CYP52A3A</i>	GGATCCTAATACGACTCACTATAGGGAGGGCCAC TCGGTAACTTTGTCAGGGAC (SEQ ID NO: 66)
3737-50-D	F	<i>CYP52A3A</i> & <i>CYP52A3B</i>	GGATCCTAATACGACTCACTATAGGGAGGCCTAC GTTTGGTATCGCTACTCCGTTG (SEQ ID NO: 67)
3737-50-C	B	<i>CYP52A3A</i> & <i>CYP52A3B</i>	TTCCAGCCAGCACCGTCCAAGCAACAAGGAGTA CAAGAAATCGTGTC (SEQ ID NO: 68)
3737-175C	F	<i>CYP52D4A</i>	GGATCCTAATACGACTCACTATAGGGAGGGCAGA GCCGATCTATGTTGCGTCC (SEQ ID NO: 69)
3737-175D	B	<i>CYP52D4A</i>	TCATTGAATGCTTCCAGGAACCTCGCCACATCCAT CGAGAACCCG (SEQ ID NO: 70)
7581-97-A	F	<i>CYP52A5A</i> & <i>CYP52A5B</i>	GGATCCTAATACGACTCACTATAGGGAGGAAGAG GGCAGGGCTCAAGAG (SEQ ID NO: 59)
7581-97-B	B	<i>CYP52A5A</i> & <i>CYP52A5B</i>	TCCATGTGAAGATCCCATCACGAGTGTGCCTCTT GCCCAAAG (SEQ ID NO: 60)
4P-2/T7	F	<i>CYP52A8A</i>	GGATCCTAATACGACTCACTATAGGGAGGCTTGA AGGCCGTGTTGAACG (SEQ ID NO: 71)
4M-3/4M-1	B	<i>CYP52A8A</i>	CAGGATTTGTCTGAGTTGCCGCTGATCAAGATA GGATCCTTGCCG (SEQ ID NO: 72)
3737-26-D	F	<i>CPRA</i>	GGATCCTAATACGACTCACTATAGGGAGGGGTTT GCTGAATACGCTGAAGGTGATG (SEQ ID NO: 73)
3737-26-C	B	<i>CPRA</i>	TGGAGCTGAACAACTCTCTCGTCTCGGTGGTCG AATGGACCTTGGTCAAG (SEQ ID NO: 74)
3737-133C	F	<i>CPRA</i> & <i>CPRB</i>	GGATCCTAATACGACTCACTATAGGGAGGTTCTT CAACACGGACAGCGG (SEQ ID NO: 75)
3737-133D	B	<i>CPRA</i> & <i>CPRB</i>	AGTCAACCAGGTGTGGAACCTCGTCGGTGGCAACA ATGAAAAACACCAAG (SEQ ID NO: 76)
3737-52-C	F	<i>POX4A</i> & <i>POX4B</i>	GGATCCTAATACGACTCACTATAGGGAGGCCATT GCCTTGAGATACGCCATTGGTAG (SEQ ID NO: 77)
3737-52-D	B	<i>POX4A</i> & <i>POX4B</i>	AGCCTTGGTGTCTGTTCTTTTCAACGGAAGGTGGT CTCGATGCTGTGTTCAACC (SEQ ID NO: 78)

3737-53-C	F	<i>POX5A</i>	GGATCCTAATACGACTCACTATAGGGAGGTTGGG TTTGTTTGTTCCTGTGTCCG (SEQ ID NO: 79)
3737-53-D	B	<i>POX5A</i>	CCTTTGACCTTCAATCTGGCGTAGACGCAGCACC ACCGATCCACCACTTG (SEQ ID NO: 80)

F=Forward B=Backward

The forward primer (SEQ ID NO: 59) contains the T7 promoter consensus sequence "GGATCCTAATACGA CTCACTATAGGG AGG" (SEQ ID NO: 109) fused to the primer 7581-97-F sequence (SEQ ID NO: 47). The Reverse Primer (SEQ ID NO: 60) contains the sequence of primer 7581-97M (SEQ ID NO: 48) followed by the 20 bases of upstream sequence with a 18 base pair deletion between the two blocks of the *CYP52A5* sequence. The forward primer was used with the corresponding reverse primer to synthesize the competitor DNA template. The primer pairs were combined in a standard *Taq* Gold polymerase PCR reaction according to the manufacturer's recommended conditions (Perkin-Elmer/Applied Biosystems, Foster City, CA). The PCR reaction mix contained a final concentration of 250 nM each primer and 10 ng *C. tropicalis* chromosomal DNA for template. The reaction mixture was placed in a thermocycler for 25 to 35 cycles using the highest annealing temperature possible during the PCR reactions to assure a homogeneous PCR product (in this case 62°C). The PCR products were either gel purified or filtered purified to remove un-incorporated nucleotides and primers. The competitor template DNA was then quantified using the ($A_{260/280}$) method. Primers used in QC-RT-PCR experiments for the synthesis of various competitive DNA templates are listed in Table 7 (SEQ ID NOS: 61-80).

C. Synthesis of the Competitor RNA

Competitor template DNA was transcribed *In-Vitro* to make the competitor RNA using the Megascript T7 kit from Ambion Biosciences (Ambion Inc., Austin, Texas). 250 nanograms (ng) of competitor DNA template and the *in-vitro* transcription reagents are mixed according to the directions provided by the manufacturer. The reaction mixture was incubated for 4 hours at 37°C. The resulting RNA preparations were then checked by gel electrophoresis for the conditions giving the highest yields and quality of competitor RNA. This often required optimization according to the manufacturer's specifications. The DNA template was then removed using DNase I as described in the Ambion kit. The RNA competitor was then quantified by the ($A_{260/280}$) method. Serial dilution's of the RNA (1 ng/ μ l to 1 femtogram (fg)/ μ l) were made for use in the QC-RT-PCR reactions and the original stocks stored at -70°C.

D. QC-RT-PCR Reactions

QC-RT-PCR reactions were performed using rTth polymerase from Perkin-Elmer(Perkin-Elmer/Applied Biosystems, Foster City, CA) according to the manufacturer's recommended conditions. The reverse transcription reaction was performed in a 10 μ l volume with a final concentrations of 200 μ M for each dNTP, 1.25 units rTth polymerase, 1.0 mM MnCl₂, 1X of the 10X buffer supplied with the Enzyme from the manufacturer, 100 ng of total RNA isolated from a fermentor grown culture of *C. tropicalis* and 1.25 μ M of the appropriate reverse primer. To quantitate *CYP52A5* expression in *C. tropicalis* an appropriate reverse primer was 7581-97M (SEQ ID NO: 48). Several reaction mixes were prepared for each RNA sample characterized. To quantitate *CYP52A5* expression a series of 8 to 12 of the previously described QC-RT-PCR reaction mixes were aliquoted to different reaction tubes. To each tube 1 μ l of a serial dilution containing from 100 pg to 100 fg *CYP52A5* competitor RNA per μ l was added bringing the final reaction mixtures up to the final volume of 10 μ l. The QC-RT-PCR reaction mixtures were mixed and incubated at 70°C for 15 min according to the manufacturer's recommended times for reverse transcription to occur. At the completion of the 15 minute incubation, the sample temperature was reduced to 4°C to stop the reaction and 40 μ l of the PCR reaction mix added to the reaction to bring the total volume up to

50 µl. The PCR reaction mix consists of an aqueous solution containing 0.3125 µM of the forward primer 7581-97F (SEQ ID NO: 47), 3.125 mM MgCl and 1X chelating buffer supplied with the enzyme from Perkin-Elmer. The reaction mixtures were placed in a thermocycler (Perkin-Elmer GeneAmp PCR System 2400, Perkin-Elmer/Applied Biosystems, Foster City, CA) and the following PCR cycle performed: 94°C for 1 min. followed by 94°C for 10 seconds followed by 58°C for 40 seconds for 17 to 22 cycles. The PCR reaction was completed with a final incubation at 58°C for 2 min followed by 4°C. In some reactions where no detectable PCR products were produced the samples were returned the thermocycler for additional cycles, this process was repeated until enough PCR products were produced to quantify using HPLC. The number of cycles necessary to produce enough PCR product is a function of the amount of the target mRNA in the 100 ng of total cellular RNA. In cultures where the *CYP52A5* gene is highly expressed there is sufficient *CYP52A5* mRNA message present and less PCR cycles (≤ 17) are required to produce quantifiable amount of PCR product. The lower the concentrations of the target mRNA present the more PCR cycles are required to produce a detectable amount of product. These QC-RT-PCR procedures were applied to all the target genes listed in Table 5 using the respective primers indicated therein.

E. HPLC Quantification

Upon completion of the QC-RT-PCR reactions the samples were analyzed and quantitated by HPLC. Five to fifteen microliters of the QC-RT-PCR reaction mix was injected into a Waters Bio-Compatible 625 HPLC with an attached Waters 484 tunable detector. The detector was set to measure a wave length of 254 nm. The HPLC contained a Sarasep brand DNASep™ column (Sarasep, Inc., San Jose, CA) which was placed within the oven and the temperature set for 52°C. The column was installed according to the manufacturer's recommendation of having 30 cm. of heated PEEK tubing installed between the injector and the column. The system was configured with a Sarasep brand Guard column positioned before the injector. In addition, there was a 0.22 µm filter disk just before the column, within the oven. Two Buffers were used to create an elution gradient to resolve and quantitate the PCR products from the QC-RT-PCR reactions. Buffer-A consists of 0.1 M tri-ethyl ammonium acetate (TEAA) and 5% acetonitrile (volume to volume). Buffer-B consists of 0.1 M TEAA and 25% acetonitrile

(volume to volume). The QC-RT-PCR samples were injected into the HPLC and the linear gradient of 75% buffer-A/ 25% buffer-B to 45% buffer-A/ 55% B was run over 6 min at a flow rate of 0.85 ml per minute. The QC-RT-PCR product of the competitor RNA being 18 base pairs smaller is eluted from the HPLC column before the QC-RT-PCR product from the *CYP52A5* mRNA(U). The amount of the QC-RT-PCR products are plotted and quantitated with an attached Waters Corporation 745 data module. The log ratios of the amount of *CYP52A5* mRNA QC-RT-PCR product (U) to competitor QC-RT-PCR product (C), as measured by peak areas, was plotted and the amount of competitor RNA required to equal the amount of *CYP52A5* mRNA product determined. In the case of each of the target genes listed in Table 5, the competitor RNA contained fewer base pairs as compared to the native target mRNA and eluted before the native mRNA in a manner similar to that demonstrated by *CYP52A5*. HPLC quantification of the genes was conducted as above.

EXAMPLE 12

Evaluation of New Strains in Shake Flasks

The *CYP* and *CPR* amplified strains such as strains HDC10, HDC15, HDC20 and HDC23 (Table 1) and H5343 were evaluated for diacid production in shake flasks. A single colony for each strain was transferred from a YPD agar plate into 5 ml of YPD broth and grown overnight at 30°C, 250 rpm. An inoculum was then transferred into 50 ml of DCA2 medium (Table 9) and grown for 24 h at 30°C, 300 rpm. The cells were centrifuged at 5000 rpm for 5 min and resuspended in 50 ml of DCA3 medium (Table 9) and grown for 24 h at 30°C, 300 rpm. 3% oleic acid w/v was added after 24 h growth in DCA3 medium and the cultures were allowed to bioconvert oleic acid for 48 h. Samples were harvested and the diacid and monoacid concentrations were analyzed as per the scheme given in Figure 35. Each strain was tested in duplicate and the results shown in Table 8 represent the average value from two flasks.

Table 8. Bioconversion of oleic acid by different recombinant strains of *Candida tropicalis*

Strain	Conversion to Oleic diacid (%)	Specific Conversion (g diacid/g biomass
H5343	41.9	0.53
HDC 10-2	50.5	0.85
HDC 15	54.4	0.85
HDC 20-1	45.1	0.72
HDC 20-2	45.3	0.58
HDC 23-2	55.2	0.84
HDC 23-3	58.8	0.89

EXAMPLE 13

Cloning and Characterization of *C. tropicalis* 20336^c Cytochrome P450 Monooxygenase (*CYP*) and Cytochrome P450 NADPH Oxidoreductase (*CPR*) Genes

To clone *CYP* and *CPR* genes several different strategies were employed. Available *CYP* amino acid sequences were aligned and regions of similarity were observed (Figure 4). These regions corresponded to described conserved regions seen in other cytochrome P450 families (Goeptar et al., *supra* and Kalb et al. *supra*). Proteins from eight eukaryotic cytochrome P450 families share a segmented region of sequence similarity. One region corresponded to the HR2 domain containing the invariant cysteine residue near the carboxyl terminus which is required for heme binding while the other region corresponded to the central region of the I helix thought to be involved in substrate recognition (Figure 4). Degenerate oligonucleotide primers corresponding to these highly conserved regions of the *CYP52* gene family present in *Candida maltosa* and *Candida tropicalis* ATCC 750 were designed and used to amplify DNA fragments of *CYP* genes from *C. tropicalis* 20336 genomic

DNA. These discrete PCR fragments were then used as probes to isolate full-length *CYP* genes from the *C. tropicalis* 20336 genomic libraries. In a few instances oligonucleotide primers corresponding to highly conserved regions were directly used as probes to isolate full-length *CYP* genes from genomic libraries. In the case of *CPR* a heterologous probe based upon the known DNA sequence for the *CPR* gene from *C. tropicalis* 750 was used to isolate the *C. tropicalis* 20336 *CPR* gene.

A. Cloning of the *CPR* Gene from *C. tropicalis* 20336

1) Cloning of the *CPRA* Allele

Approximately 25,000 phage particles from the first genomic library of *C. tropicalis* 20336 were screened with a 1.9 kb *Bam*HI-*Nde*I fragment from plasmid pCU3RED (See Picattagio et al., Bio/Technology 10:894-898 (1992), incorporated herein by reference) containing most of the *C. tropicalis* 750 *CPR* gene. Five clones that hybridized to the probe were isolated and the plasmid DNA from these lambda clones was rescued and characterized by restriction enzyme analysis. The restriction enzyme analysis suggested that all five clones were identical but it was not clear that a complete *CPR* gene was present.

PCR analysis was used to determine if a complete *CPR* gene was present in any of the five clones. Degenerate primers were prepared for highly conserved regions of known *CPR* genes (See Sutter et al., *J. Biol. Chem.* 265:16428-16436 (1990), incorporated herein by reference) (Figure 4). Two Primers were synthesized for the FMN binding region (FMN1, SEQ ID NO: 16 and FMN2, SEQ ID NO: 17). One primer was synthesized for the FAD binding region (FAD, SEQ ID NO: 18), and one primer for the NADPH binding region (NADPH, SEQ ID NO: 19) (Table 4). These four primers were used in PCR amplification experiments using as a template plasmid DNA isolated from four of the five clones described above. The FMN (SEQ ID NOS: 16 and 17) and FAD (SEQ ID NO: 18) primers served as forward primers and the NADPH primer (SEQ ID NO: 19) as the reverse primer in the PCR reactions. When different combinations of forward and reverse primers were used, no PCR products were obtained from any of the plasmids. However, all primer combinations amplified expected size products with a plasmid containing the *C. tropicalis* 750 *CPR* gene (positive

control). The most likely reason for the failure of the primer pairs to amplify a product, was that all four of clones contained a truncated *CPR* gene. One of the four clones (pHKM1) was sequenced using the Triplex 5' (SEQ ID NO: 30) and the Triplex 3' (SEQ ID NO: 31) primers (Table 4) which flank the insert and the multiple cloning site on the cloning vector, and with the degenerate primer based upon the NADPH binding site described above. The NADPH primer (SEQ ID NO: 19) failed to yield any sequence data and this is consistent with the PCR analysis. Sequences obtained with Triplex primers were compared with *C. tropicalis* 750 *CPR* sequence using the MacVector™ program (Oxford Molecular Group, Campbell, CA). Sequence obtained with the Triplex 3' primer (SEQ ID NO: 31) showed similarity to an internal sequence of the *C. tropicalis* 750 *CPR* gene confirming that pHKM1 contained a truncated version of a 20336 *CPR* gene. pHKM1 had a 3.8 kb insert which included a 1.2 kb coding region of the *CPR* gene accompanied by 2.5 kb of upstream DNA (Figure 5). Approximately 0.85 kb of the 20336 *CPR* gene encoding the C-terminal portion of the *CPR* protein is missing from this clone.

Since the first Clontech library yielded only a truncated *CPR* gene, the second library prepared by Clontech was screened to isolate a full-length *CPR* gene. Three putative *CPR* clones were obtained. The three clones, having inserts in the range of 5-7 kb, were designated pHKM2, pHKM3 and pHKM4. All three were characterized by PCR using the degenerate primers described above. Both pHKM2 and pHKM4 gave PCR products with two sets of internal primers. pHKM3 gave a PCR product only with the FAD (SEQ ID NO: 18) and NADPH (SEQ ID NO: 19) primers suggesting that this clone likely contained a truncated *CPR* gene. All three plasmids were partially sequenced using the two Triplex primers and a third primer whose sequence was selected from the DNA sequence near the truncated end of the *CPR* gene present in pHKM1. This analysis confirmed that both pHKM2 & 4 have sequences that overlap pHKM1 and that both contained the 3' region of *CPR* gene that is missing from pHKM1. Portions of inserts from pHKM1 and pHKM4 were sequenced and a full-length *CPR* gene was identified. Based on the DNA sequence and PCR analysis, it was concluded that pHKM1 contained the putative promoter region and 1.2 kb of sequence encoding a portion (5' end) of a *CPR* gene. pHKM4 had 1.1 kb of DNA that overlapped pHKM1 and contained the remainder (3' end) of a *CPR* gene along with a downstream untranslated region (Figure 6).

Together these two plasmids contained a complete *CPRA* gene with an upstream promoter region. *CPRA* is 4206 nucleotides in length (SEQ ID NO: 81) and includes a regulatory region and a protein coding region (defined by nucleotides 1006-3042) which is 2037 base pairs in length and codes for a putative protein of 679 amino acids (SEQ ID NO: 83) (Figures 13 and 14). In Figure 13, the asterisks denote conserved nucleotides between *CPRA* and *CPRB*, bold denotes protein coding nucleotides, and the start and stop codons are underlined. The *CPRA* protein, when analyzed by the protein alignment program of the GeneWorks™ software package (Oxford Molecular Group, Campbell, CA), showed extensive homology to *CPR* proteins from *C. tropicalis* 750 and *C. maltosa*.

2) Cloning of the *CPRB* Allele

To clone the second *CPRB* allele, the third genomic library, prepared by Henkel, was screened using DNA fragments from pHKM1 and pHKM4 as probes. Five clones were obtained and these were sequenced with the three internal primers used to sequence *CPRA*. These primers were designated PRK1.F3 (SEQ ID NO: 20), PRK1.F5 (SEQ ID NO: 21) and PRK4.R20 (SEQ ID NO: 22) (Table 4). and the two outside primers (M13 -20 and T3 [Stratagene]) for the polylinker region present in the pBK-CMV cloning vector. Sequence analysis suggested that four of these clones, designated pHKM5 to 8, contained inserts which were identical to the *CPRA* allele isolated earlier. All four seemed to contain a full length *CPR* gene. The fifth clone was very similar to the *CPRA* allele, especially in the open reading frame region where the identity was very high. However, there were significant differences in the 5' and 3' untranslated regions. This suggested that the fifth clone was the allele to *CPRA*. The plasmid was designated pHKM9 (Figure 7) and a 4.14 kb region of this plasmid was sequenced and the analysis of this sequence confirmed the presence of the *CPRB* allele (SEQ ID NO: 82), which includes a regulatory region and a protein coding region (defined by nucleotides 1033-3069) (Figure 13). The amino acid sequence of the *CPRB* protein is set forth in SEQ ID NO: 84 (Figure 14).

B. Cloning of *C. tropicalis* 20336 (*CYP*) Genes

1) Cloning of *CYP52A2A*, *CYP52A3A* & *3B* and *CYP52A5A* & *5B*

Clones carrying *CYP52A2A*, *A3A*, *A3B*, *A5A* and *A5B* genes were isolated from the first and second Clontech genomic libraries using an oligonucleotide probe (HemeB1, SEQ ID NO: 27) whose sequence was based upon the amino acid sequence for the highly conserved heme binding region present throughout the *CYP52* family. The first and second libraries were converted to the plasmid form and screened by colony hybridizations using the HemeB1 probe (SEQ ID NO: 27) (Table 4). Several potential clones were isolated and the plasmid DNA was isolated from these clones and sequenced using the HemeB1 oligonucleotide (SEQ ID NO: 27) as a primer. This approach succeeded in identifying five *CYP52* genes. Three of the *CYP* genes appeared unique, while the remaining two were classified as alleles. Based upon an arbitrary choice of homology to *CYP52* genes from *Candida maltosa*, these five genes and corresponding plasmids were designated *CYP52A2A* (pPA15 [Figure 26]), *CYP52A3A* (pPA57 [Figure 29]), *CYP52A3B* (pPA62 [Figure 30]), *CYP52A5A* (pPAL3 [Figure 31]) and *CYP52A5B* (pPA5 [Figure 32]). The complete DNA sequence including regulatory and protein coding regions of these five genes was obtained and confirmed that all five were *CYP52* genes (Figure 15). In Figure 15, the asterisks denote conserved nucleotides among the *CYP* genes. Bold indicates the protein coding nucleotides of the *CYP* genes, and the start and stop codons are underlined. The *CYP52A2A* gene as represented by SEQ ID NO: 86 has a protein coding region defined by nucleotides 1199-2767 and the encoded protein has an amino acid sequence as set forth in SEQ ID NO: 96. The *CYP52A3A* gene as represented by SEQ ID NO: 88 has a protein encoding region defined by nucleotides 1126-2748 and the encoded protein has an amino acid sequence as set forth in SEQ ID NO: 98. The *CYP52A3B* gene as represented by SEQ ID NO: 89 has a protein coding defined by nucleotides 913-2535 and the encoded protein has an amino acid sequence as set forth in SEQ ID NO: 99. The *CYP52A5A* gene as represented by SEQ ID NO: 90 has a protein coding region defined by nucleotides 1103-2656 and the encoded protein has an amino acid sequence as set forth in SEQ ID NO: 100. The *CYP52A5B* gene as represented by SEQ ID NO: 91 has a protein coding

region defined by nucleotides 1142-2695 and the encoded protein has an amino acid sequence as set forth in SEQ ID NO: 101.

2) Cloning of *CYP52A1A* and *CYP52A8A*

CYP52A1A and *CYP52A8A* genes were isolated from the third genomic library using PCR fragments as probes. The PCR fragment probe for *CYP52A1* was generated after PCR amplification of 20336 genomic DNA with oligonucleotide primers that were designed to amplify a region from the Helix I region to the HR2 region using all available *CYP52* genes from National Center for Biotechnology Information. Degenerate forward primers UCup1 (SEQ ID NO: 23) and UCup2 (SEQ ID NO: 24) were designed based upon an amino acid sequence (-RDTTAG-) from the Helix I region (Table 4). Degenerate primers UCdown1 (SEQ ID NO: 25) and UCdown2 (SEQ ID NO: 26) were designed based upon an amino acid sequence (-GQQFAL-) from the HR2 region (Table 4). For the reverse primers, the DNA sequence represents the reverse complement of the corresponding amino acid sequence. These primers were used in pairwise combinations in a PCR reaction with Stoffel *Taq* DNA polymerase (Perkin-Elmer Cetus, Foster City, CA) according to the manufacturer's recommended procedure. A PCR product of approximately 450 bp was obtained. This product was purified from agarose gel using Gene-clean™ (Bio 101, LaJolla, CA) and ligated to the pTAG™ vector (Figure 17) (R&D systems, Minneapolis, MN) according to the recommendations of the manufacturer. No treatment was necessary to clone into pTAG because it employs the use of the TA cloning technique. Plasmids from several transformants were isolated and their inserts were characterized. One plasmid contained the PCR clone intact. The DNA sequence of the PCR fragment (designated 44*CYP3*, SEQ ID NO: 107) shared homology with the DNA sequences for the *CYP52A1* gene of *C. maltosa* and the *CYP52A3* gene of *C. tropicalis* 750. This fragment was used as a probe in isolating the *C. tropicalis* 20336 *CYP52A1* homolog. The third genomic library was screened using the 44*CYP3* PCR probe (SEQ ID NO: 107) and a clone (pHKM11) that contained a full-length *CYP52* gene was obtained (Figure 8). The clone contained a gene having regulatory and protein coding regions. An open reading frame of 1572 nucleotides encoded a *CYP52* protein of 523 amino acids (Figures 15 and 16). This *CYP52*

gene was designated *CYP52A1A* (SEQ ID NO: 85) since its putative amino acid sequence (SEQ ID NO: 95) was most similar to the *CYP52A1* protein of *C. maltosa*. The protein coding region of the *CYP52A1A* gene is defined by nucleotides 1177-2748 of SEQ ID NO: 85.

A similar approach was taken to clone *CYP52A8A*. A PCR fragment probe for *CYP52A8* was generated using primers for highly conserved sequences of *CYP52A3*, *CYP52A2* and *CYP52A5* genes of *C. tropicalis* 750. The reverse primer (primer 2,3,5,M) (SEQ ID NO: 29) was designed based on the highly conserved heme binding region (Table 4). The design of the forward primer (primer 2,3,5,P) (SEQ ID NO: 28) was based upon a sequence conserved near the N-terminus of the *CYP52A3*, *CYP52A2* and *CYP52A5* genes from *C. tropicalis* 750 (Table 4). Amplification of 20336 genomic DNA with these two primers gave a mixed PCR product. One amplified PCR fragment was 1006 bp long (designated DCA1002). The DNA sequence for this fragment was determined and was found to have 85% identity to the DNA sequence for the *CYP52D4* gene of *C. tropicalis* 750. When this PCR product was used to screen the third genomic library one clone (pHKM12) was identified that contained a full-length *CYP52* gene along with 5' and 3' flanking sequences (Figure 9). The *CYP52* gene included regulatory and protein coding regions with an open reading frame of 1539 nucleotides long which encoded a putative *CYP52* protein of 512 amino acids (Figures 15 and 16). This gene was designated as *CYP52A8A* (SEQ ID NO: 92) since its amino acid sequence (SEQ ID NO: 102) was most similar to the *CYP52A8* protein of *C. maltosa*. The protein coding region of the *CYP52A8A* gene is defined by nucleotides 464-2002 of SEQ ID NO: 92. The amino acid sequence of the *CYP52A8A* protein is set forth in SEQ ID NO: 102.

3) Cloning of *CYP52D4A*

The screening of the second genomic library with the HemeB1 (SEQ ID NO: 27) primer (Table 4) yielded a clone carrying a plasmid (pPA18) that contained a truncated gene having homology with the *CYP52D4* gene of *C. maltosa* (Figure 33). A 1.3 to 1.5-kb *EcoRI*-*SstI* fragment from pPA18 containing part of the truncated *CYP* gene was isolated and used as a probe to screen the third genomic library for a full length *CYP52* gene. One clone (pHKM13) was isolated and found to contain a full-length *CYP* gene with extensive 5' and 3' flanking

sequences (Figure 10). This gene has been designated as *CYP52D4A* (SEQ ID NO: 94) and the complete DNA including regulatory and protein coding regions (coding region defined by nucleotides 767-2266) and putative amino acid sequence (SEQ ID NO: 104) of this gene is shown in Figures 15 and 16. *CYP52D4A* (SEQ ID NO: 94) shares the greatest homology with the *CYP52D4* gene of *C. maltosa*.

4) Cloning of *CYP52A2B* and *CYP52A8B*

A mixed probe containing *CYP52A1A*, *A2A*, *A3A*, *D4A*, *A5A* and *A8A* genes was used to screen the third genomic library and several putative positive clones were identified. Seven of these were sequenced with the degenerate primers Cyp52a (SEQ ID NO: 32), Cyp52b (SEQ ID NO: 33), Cyp52c (SEQ ID NO: 34) and Cyp52d (SEQ ID NO: 35) shown in Table 4. These primers were designed from highly conserved regions of the four *CYP52* subfamilies, namely *CYP52A*, *B*, *C* & *D*. Sequences from two clones, pHKM14 and pHKM15 (Figures 11 and 12), shared considerable homology with DNA sequence of the *C. tropicalis* 20336 *CYP52A2* and *CYP52A8* genes, respectively. The complete DNA (SEQ ID NO: 87) including regulatory and protein coding regions (coding region defined by nucleotides 1072-2640) and putative amino acid sequence (SEQ ID NO: 97) of the *CYP52* gene present in pHKM14 suggested that it is *CYP52A2B* (Figures 15 and 16). The complete DNA (SEQ ID NO: 93) including regulatory and protein coding regions (coding region defined by nucleotides 1017-2555) and putative amino acid sequence (SEQ ID NO: 103) of the *CYP52* gene present in pHKM15 suggested that it is *CYP52A8B* (Figures 15 and 16).

EXAMPLE 14

Identification of *CYP* and *CPR* Genes Induced by Selected Fatty Acid and Alkane Substrates

Genes whose transcription is turned on by the presence of selected fatty acid or alkane substrates have been identified using the QC-RT-PCR assay. This assay was used to measure (*CYP*) and (*CPR*) gene expression in fermentor grown cultures *C. tropicalis* ATCC 20962. This method involves the isolation of total cellular RNA from cultures of *C. tropicalis* and

the quantification of a specific mRNA within that sample through the design and use of sequence specific QC-RT-PCR primers and an RNA competitor. Quantification is achieved through the use of known concentrations of highly homologous competitor RNA in the QC-RT-PCR reactions. The resulting QC-RT-PCR amplified cDNA's are separated and quantitated through the use of ion pairing reverse phase HPLC. This assay was used to characterize the expression of *CYP52* genes of *C. tropicalis* ATCC 20962 in response to various fatty acid and alkane substrates. Genes which were induced were identified by the calculation of their mRNA concentration at various times before and after induction. Figure 18 provides an example of how the concentration of mRNA for *CYP52A5* can be calculated using the QC-RT-PCR assay. The log ratio of unknown (U) to competitor product (C) is plotted versus the concentration of competitor RNA present in the QC-RT-PCR reactions. The concentration of competitor which results in a log ratio of U/C of zero, represents the point where the unknown messenger RNA concentration is equal to the concentration of the competitor. Figure 18 allows for the calculation of the amount of *CYP52A5* message present in 100 ng of total RNA isolated from cell samples taken at 0, 1, and 2 hours after the addition of Emersol® 267 in a fermentor run. From this analysis, it is possible to determine the concentration of the *CYP52A5* mRNA present in 100 ng of total cellular RNA. In the plot contained in Figure 18 it takes 0.46 pg of competitor to equal the number of mRNA's of *CYP52A5* in 100 ng of RNA isolated from cells just prior (time 0) to the addition of the substrate, Emersol® 267. In cell samples taken at one and two hours after the addition of Emersol® 267 it takes 5.5 and 8.5 pg of competitor RNA, respectively. This result demonstrates that *CYP52A5* (SEQ ID NOS: 90 and 91) is induced more than 18 fold within two hours after the addition of Emersol® 267. This type of analysis was used to demonstrate that *CYP52A5* (SEQ ID NO: 90 and 91) is induced by Emersol® 267. Figure 19 shows the relative amounts of *CYP52A5* (SEQ ID NOS: 90 and 91) expression in fermentor runs with and without Emersol® 267 as a substrate. The differences in the *CYP52A5* (SEQ. ID NOS: 90 and 91) expression patterns are due to the addition of Emersol® 267 to the fermentation medium.

This analysis clearly demonstrates that expression of *CYP52A5* (SEQ ID NOS: 90 and 91) in *C. tropicalis* 20962 is inducible by the addition of Emersol® 267 to the growth medium. This analysis was performed to characterize the expression of *CYP52A2A* (SEQ ID

NO: 86) , *CYP52A3AB* (SEQ ID NOS: 88 and 89) , *CYP52A8A* (SEQ ID NO: 92) , *CYP52A1A* (SEQ ID NO: 85), *CYP52D4A* (SEQ ID NO: 94) and *CPRB* (SEQ ID NO: 82) in response to the presence of Emersol® 267 in the fermentation medium (Figure 20). The results of these analysis' indicate, that like the *CYP52A5* gene (SEQ ID NOS: 90 and 91) of *C. tropicalis* 20962, the *CYP52A2A* gene (SEQ ID NO: 86) is inducible by Emersol® 267. A small induction is observed for *CYP52A1A* (SEQ ID NO: 85) and *CYP52A8A* (SEQ ID NO: 92). In contrast, any induction for *CYP52D4A* (SEQ ID NO: 94), *CYP52A3A* (SEQ ID NO: 88), *CYP52A3B* (SEQ ID NO: 89) is below the level of detection of the assay. *CPRB* (SEQ ID NO: 82) is moderately induced by Emersol® 267, four to five fold. The results of these analysis are summarized in Figure 20. Figure 34 provides an example of selective induction of *CYP52A* genes. When pure fatty acid or alkanes are spiked into a fermentor containing *C. tropicalis* 20962 or a derivative thereof, the transcriptional activation of *CYP52A* genes was detected using the QC-RT-PCR assay. Figure 34 shows that pure oleic acid (C18:1) strongly induces *CYP52A2A* (SEQ ID NO: 86) while inducing *CYP52A5* (SEQ ID NOS: 90 and 91). In the same fermentor addition of pure alkane (tridecane) shows strong induction of both *CYP52A2A* (SEQ ID NO: 86) and *CYP52A1A* (SEQ ID NO: 85). However, tridecane did not induce *CYP52A5* (SEQ ID NOS: 90 and 91) . In a separate fermentation using ATCC 20962, containing pure octadecane as the substrate, induction of *CYP52A2A*, *CYP52A5A* and *CYP52A1A* is detected (see Figure 36). The foregoing demonstrates selective induction of particular *CYP* genes by specific substrates, thus providing techniques for selective metabolic engineering of cell strains. For example, if tridecane modification is desired, organisms engineered for high levels of *CYP52A2A* (SEQ ID NO: 86) and *CYP52A1A* (SEQ ID NO: 85) activity are indicated. If oleic acid modification is desired, organisms engineered for high levels of *CYP52A2A* (SEQ ID NO: 86) activity are indicated.

EXAMPLE 15

Integration of Selected *CYP* and *CPR* Genes into the Genome of *Candida tropicalis*

In order to integrate selected genes into the chromosome of *C. tropicalis* 20336 or its descendants, there has to be a target DNA sequence, which may or may not be an intact gene, into which the genes can be inserted. There must also be a method to select for the integration event. In some cases the target DNA sequence and the selectable marker are the same and, if so, then there must also be a method to regain use of the target gene as a selectable marker following the integration event. In *C. tropicalis* and its descendants, one gene which fits these criteria is *URA3A*, encoding orotidine-5'-phosphate decarboxylase. Using it as a target for integration, *ura⁻* variants of *C. tropicalis* can be transformed in such a way as to regenerate a *URA⁺* genotype via homologous recombination (Figure 21). Depending upon the design of the integration vector, one or more genes can be integrated into the genome at the same time. Using a split *URA3A* gene oriented as shown in Figure 22, homologous integration would yield at least one copy of the gene(s) of interest which are inserted between the split portions of the *URA3A* gene. Moreover, because of the high sequence similarity between *URA3A* and *URA3B* genes, integration of the construct can occur at both the *URA3A* and *URA3B* loci. Subsequently, an oligonucleotide designed with a deletion in a portion of the *URA* gene based on the identical sequence across both the *URA3A* and *URA3B* genes, can be utilized to yield *C. tropicalis* transformants which are once again *ura⁻* but which still carry one or more newly integrated genes of choice (Figure 21). *ura⁻* variants of *C. tropicalis* can also be isolated via other methods such as classical mutagenesis or by spontaneous mutation. Using well established protocols, selection of *ura⁻* strains can be facilitated by the use of 5-fluoroorotic acid (5-FOA) as described, e.g., in Boeke et al., *Mol. Gen. Genet.* 197:345-346, (1984), incorporated herein by reference. The utility of this approach for the manipulation of *C. tropicalis* has been well documented as described, e.g., in Picataggio et al., *Mol. and Cell. Biol.* 11:4333-4339 (1991); Rohrer et al., *Appl. Microbiol. Biotechnol.* 36:650-654 (1992); Picataggio et al., *Bio/Technology* 10:894-898 (1992); U.S. Patent No. 5,648,247; U.S. Patent No. 5,620,878; U.S. Patent No. 5,204,252; U.S. Patent No. 5,254,466, all of which are incorporated herein by reference.

A. Construction of a URA Integration Vector, pURain.

Primers were designed and synthesized based on the 1712 bp sequence of the *URA3A* gene of *C. tropicalis* 20336 (see Figure 23). The nucleotide sequence of the *URA3A* gene of *C. tropicalis* 20336 is set forth in SEQ ID NO: 105 and the amino acid sequence of the encoded protein is set forth in SEQ ID NO: 106. *URA3A* Primer Set #1a (SEQ ID NO: 9) and #1b (SEQ ID NO: 10) (Table 4) was used in PCR with *C. tropicalis* 20336 genomic DNA to amplify *URA3A* sequences between nucleotide 733 and 1688 as shown in Figure 23. The primers are designed to introduce unique 5' *Asc*I and 3' *Pac*I restriction sites into the resulting amplified *URA3A* fragment. *Asc*I and *Pac*I sites were chosen because these sites are not present within *CYP* or *CPR* genes identified to date. *URA3A* Primer Set #2 was used in PCR with *C. tropicalis* 20336 genomic DNA as a template, to amplify *URA3A* sequences between nucleotide 9 and 758 as shown in Figure 23. *URA3A* Primer set #2a (SEQ ID NO: 11) and #2b (SEQ ID NO: 12) (Table 4) was designed to introduce unique 5' *Pac*I and 3' *Pme*I restriction sites into the resulting amplified *URA3A* fragment. The *Pme*I site is also not present within *CYP* and *CPR* genes identified to date. PCR fragments of the *URA3A* gene were purified, restricted with *Asc*I, *Pac*I and *Pme*I restriction enzymes and ligated to a gel purified, QiaexII cleaned *Asc*I-*Pme*I digest of plasmid pNEB193 (Figure 25) purchased from New England Biolabs (Beverly, MA). The ligation was performed with an equimolar number of DNA termini at 16 °C for 16 hr using T4 DNA ligase (New England Biolabs). Ligations were transformed into *E. coli* XL1-Blue cells (Stratagene, LaJolla, CA) according to manufacturers recommendations. White colonies were isolated, grown, plasmid DNA isolated and digested with *Asc*I-*Pme*I to confirm insertion of the modified *URA3A* into pNEB193. The resulting base integration vector was named pURain (Figure 24).

B. Amplification of *CYP52A2A*, *CYP52A3A*, *CYP52A5A* and *CPRB* from *C. tropicalis* 20336 Genomic DNA

The genes encoding *CYP52A2A*, (SEQ ID NO: 86) and *CYP52A3A* (SEQ ID NO: 88) from *C. tropicalis* 20336 were amplified from genomic clones (pPA15 and pPA57, respectively) (Figures 26 and 29) via PCR using primers (Primer *CYP2A*#1, SEQ ID NO: 1 and Primer *CYP2A*#2, SEQ ID NO: 2 for *CYP52A2A*) (Primer *CYP3A*#1, SEQ ID NO: 3 and Primer *CYP3A*#2, SEQ ID NO: 4 for *CYP52A3A*) to introduce *PacI* cloning sites. These PCR primers were designed based upon the DNA sequence determined for *CYP52A2A* (SEQ ID NO: 86) (Figure 15). The *AmpliTaq* Gold PCR kit (Perkin Elmer Cetus, Foster City, CA) was used according to manufacturers specifications. The *CYP52A2A* PCR amplification product was 2,230 base pairs in length, yielding 496 bp of DNA upstream of the *CYP52A2A* start codon and 168 bp downstream of the stop codon for the *CYP52A2A* ORF. The *CYP52A3A* PCR amplification product was 2154 base pairs in length, yielding 437bp of DNA upstream of the *CYP52A3A* start codon and 97bp downstream of the stop codon for the *CYP52A3A* ORF. The *CYP52A3A* PCR amplification product was 2154 base pairs in length, yielding 437bp of DNA upstream of the *CYP52A3A* start codon and 97bp downstream of the stop codon for the *CYP52A3A* ORF.

The gene encoding *CYP52A5A* (SEQ ID NO: 90) from *C. tropicalis* 20336 was amplified from genomic DNA via PCR using primers (Primer *CYP 5A*#1, SEQ ID NO: 5 and Primer *CYP 5A*#2, SEQ ID NO: 6) to introduce *PacI* cloning sites. These PCR primers were designed based upon the DNA sequence determined for *CYP52A5A* (SEQ ID NO: 90). The *Expand Hi-Fi Taq* PCR kit (Boehringer Mannheim, Indianapolis, IN) was used according to manufacturers specifications. The *CYP52A5A* PCR amplification product was 3,298 base pairs in length.

The gene encoding *CPRB* (SEQ ID NO: 82) from *C. tropicalis* 20336 was amplified from genomic DNA via PCR using primers (*CPR B*#1, SEQ ID NO: 7 and *CPR B*#2, SEQ ID NO: 8) based upon the DNA sequence determined for *CPRB* (SEQ ID NO: 82) (Figure 13). These primers were designed to introduce unique *PacI* cloning sites. The *Expand Hi-Fi Taq* PCR kit (Boehringer Mannheim, Indianapolis, IN) was used according to manufacturers specifications. The *CPRB* PCR product was 3266 bp in length, yielding 747 bp of DNA upstream of the *CPRB* start codon and 493 bp downstream of the stop codon for the *CPRB*

ORF. The resulting PCR products were isolated via agarose gel electrophoresis, purified using QiaexII and digested with *PacI*. The PCR fragments were purified, desalted and concentrated using a Microcon 100 (Amicon, Beverly, MA).

The above described amplification procedures are applicable to the other genes listed in Table 5 using the respectively indicated primers.

C. Cloning of *CYP* and *CPR* Genes into pURAIin.

The next step was to clone the selected *CYP* and *CPR* genes into the pURAIin integration vector. In a preferred aspect of the present invention, no foreign DNA other than that specifically provided by synthetic restriction site sequences are incorporated into the DNA which was cloned into the genome of *C. tropicalis*, i.e., with the exception of restriction site DNA only native *C. tropicalis* DNA sequences are incorporated into the genome. pURAIin was digested with *PacI*, Qiaex II cleaned, and dephosphorylated with Shrimp Alkaline Phosphatase (SAP) (United States Biochemical, Cleveland, OH) according the manufacturer's recommendations. Approximately 500 ng of *PacI* linearized pURAIin was dephosphorylated for 1 hr at 37°C using SAP at a concentration of 0.2 Units of enzyme per 1 pmol of DNA termini. The reaction was stopped by heat inactivation at 65°C for 20 min.

The *CYP52A2A* *PacI* fragment derived using the primer shown in Table 4 was ligated to plasmid pURAIin which had also been digested with *PacI*. *PacI* digested pURAIin was dephosphorylated, and ligated to the *CYP52A2A* ULTMA PCR product as described previously. The ligation mixture was transformed into *E. coli* XL1 Blue MRF' (Stratagene) and 2 resistant colonies were selected and screened for correct constructs which should contain vector sequence, the inverted *URA3A* gene, and the amplified *CYP52A2A* gene (SEQ ID NO: 86) of 20336. *AscI-PmeI* digestion identified one of the two constructs, plasmid pURA2in, as being correct (Figure 27). This plasmid was sequenced and compared to *CYP52A2A* (SEQ ID NO: 86) to confirm that PCR did not introduce DNA base changes that would result in an amino acid change.

Prior to its use, the *CPRB* *PacI* fragment derived using the primers shown in Table 4 was sequenced and compared to *CPRB* (SEQ ID NO: 82) to confirm that PCR did not introduce DNA base pair changes that would result in an amino acid change. Following

confirmation, *CPRB* (SEQ ID NO: 82) was ligated to plasmid pURain which had also been digested with *PacI*. *PacI* digested pURain was dephosphorylated, and ligated to the *CPR* Expand Hi-Fi PCR product as described previously. The ligation mixture was transformed into *E. coli* XL1 Blue MRF' (Stratagene) and several resistant colonies were selected and screened for correct constructs which should contain vector sequence, the inverted *URA3A* gene, and the amplified *CPRB* gene (SEQ ID NO: 82) of 20336. *AscI-PmeI* digestion confirmed a successful construct, pURAREDBin.

In a manner similar to the above, each of the other *CYP* and *CPR* genes disclosed herein are cloned into pURain. *PacI* fragments of these genes, whose sequences are given in Figures 13 and 15, are derivable by methods known to those skilled in the art.

1) Construction of Vectors Used to Generate HDC 20 and HDC 23

A previously constructed integration vector containing *CPRB* (SEQ ID NO: 82), pURAREDBin, was chosen as the starting vector. This vector was partially digested with *PacI* and the linearized fragment was gel-isolated. The active *PacI* was destroyed by treatment with T4 DNA polymerase and the vector was re-ligated. Subsequent isolation and complete digestion of this new plasmid yielded a vector now containing only one active *PacI* site. This fragment was gel-isolated, dephosphorylated and ligated to the *CYP52A2A* *PacI* fragment. Vectors that contain the *CYP52A2A* (SEQ ID NO: 86) and *CPRB* (SEQ ID NO: 82) genes oriented in the same direction, pURain *CPR* 2A S, as well as opposite directions (5' ends connected), pURain *CPR* 2A O, were generated.

D. Confirmation of *CYP* Integration (Figure 21 for Integration Scheme) into the Genome of *C. tropicalis*

Based on the construct, pURA2in, used to transform H5343 *ura⁻*, a scheme to detect integration was devised. Genomic DNA from transformants was digested with *Dra* III and *Spe* I which are enzymes that cut within the *URA3A*, and *URA3B* genes but not within the integrated *CYP52A2A* gene. Digestion of genomic DNA where an integration had occurred at the *URA3A* or *URA3B* loci would be expected to result in a 3.5 kb or a 3.3 kb fragment, respectively (Figure 28). Moreover, digestion of the same genomic DNA with *PacI* would yield a

2.2 kb fragment characteristic for the integrated *CYP52A2A* gene (Figure 28). Southern hybridizations of these digests with fragments of the *CYP52A2A* gene were used to screen for these integration events. Intensity of the band signal from the Southern using *PacI* digestion was used as a measure of the number of integration events, ((i.e. the more copies of the *CYP52A2A* gene (SEQ ID NO: 86) which are present, the stronger the hybridization signal)).

C. tropicalis H5343 transformed *URA* prototrophs were grown at 30°C, 170 rpm, in 10 ml SC-uracil media for preparation of genomic DNA. Genomic DNA was isolated by the method described previously. Genomic DNA was digested with *SpeI* and *DraIII*. A 0.95% agarose gel was used to prepare a Southern hybridization blot. The DNA from the gel was transferred to a MagnaCharge nylon filter membrane (MSI Technologies, Westboro, MA) according to the alkaline transfer method of Sambrook et al., *supra*. For the Southern hybridization, a 2.2 kb *CYP52A2A* DNA fragment was used as a hybridization probe. 300 ng of *CYP52A2A* DNA was labeled using a ECL Direct labeling and detection system (Amersham) and the Southern was processed according to the ECL kit specifications. The blot was processed in a volume of 30 ml of hybridization fluid corresponding to 0.125 ml/cm². Following a prehybridization at 42°C for 1 hr, 300 ng of *CYP52A2A* probe was added and the hybridization continued for 16 hr at 42°C. Following hybridization, the blots were washed two times for 20 min each at 42 °C in primary wash containing urea. Two 5 min secondary washes at RT were conducted, followed by detection according to directions. The blots were exposed for 16 hours (hr) as recommended.

Integration was confirmed by the detection of a *SpeI*-*DraIII* 3.5 kb fragment from the genomic DNA of the transformants but not with the *C. tropicalis* 20336 control. Subsequently, a *PacI* digestion of the genomic DNA of the positive transformants, followed by a Southern hybridization using an *CYP52A2A* gene probe, confirmed integration by the detection of a 2.2 kb fragment. The resulting *CYP52A2A* integrated strain was named HDC1 (see Table 1).

In a manner similar to the above, each of the genes contained in the *PacI* fragments which are described in Section 3c above were confirmed for integration into the genome of *C. tropicalis*.

Transformants generated by transformation with the vectors, pURAI in *CPR* 2A S or pURAI in *CPR* 2A O, were analyzed by Southern hybridization for integration of both the *CYP52A2A* (SEQ ID NO: 86) and *CPRB* (SEQ ID NO: 82) genes tandemly. Three strains were generated in which the *CYP52A2A* (SEQ ID NO: 86) and *CPRB* (SEQ ID NO: 82) genes integrated are in the opposite orientation (HDC 20-1, HDC 20-2 and HDC 20-3) and three were generated with the *CYP52A2A* (SEQ ID NO: 86) and *CPRB* (SEQ ID NO: 82) genes integrated in the same orientation (HDC 23-1, HDC 23-2 and HDC 23-3), Table 1.

E. Confirmation of *CPRB* Integration into H5343 *ura*

Seven transformants were screened by colony PCR using *CPRB* primer #2 (SEQ ID NO: 8) and a *URA3A*-specific primer. In five of the transformants, successful integration was detected by the presence of a 3899 bp PCR product. This 3899 bp PCR product represents the *CPRB* gene adjacent to the *URA3A* gene in the genome of H5343 thereby confirming integration. The resulting *CPRB* integrated strains were named HDC10-1 and HDC10-2 (see Table 1).

F. Strain Evaluation.

As determined by quantitative PCR, when compared to parent H5343, HDC10-1 contained three additional copies of the reductase gene and HDC10-2 contained four additional copies of the reductase gene. Evaluations of HDC20-1, HDC20-2 and HDC20-3 based on Southern hybridization data indicates that HDC20-1 contained multiple integrations, i.e., 2 to 3 times that of HDC20-2 or HDC20-3. Evaluations of HDC23-1, HDC23-2, and HDC23-3 based on Southern hybridization data indicates that HDC23-3 contained multiple integrations, i.e., 2 to 3 times that of HDC23-1 or HDC23-2. The data in Table 8 indicates that the integration of components of the ω -hydroxylase complex have a positive effect on the improvement of *Candida tropicalis* ATCC 20962 as a biocatalyst. The results indicate that *CYP52A5A* (SEQ ID NO: 90) is an important gene for the conversion of oleic acid to diacid. Surprisingly, tandem integrations of *CYP* and *CPR* genes oriented in the opposite direction (HDC 20 strains) seem to

Table 9

Media Composition

LB Broth

Bacto Tryptone	10 g
Bacto Yeast Extract	5 g
Sodium Chloride	10 g
Distilled Water	1,000 ml

LB Agar

Bacto Tryptone	10 g
Bacto Yeast Extract	5 g
Sodium Chloride	10 g
Agar	15 g
Distilled Water	1,000 ml

LB Top Agarose

Bacto Tryptone	10 g
Bacto Yeast Extract	5 g
Sodium Chloride	10 g
Agarose	7 g
Distilled Water	1,000 ml

NZCYM Broth

Bacto Casein Digest	10 g
Bacto Casamino Acids	1 g
Bacto Yeast Extract	5 g
Sodium Chloride	5 g
Magnesium Sulfate 0.98 g (anhydrous)	
Distilled Water	1,000 ml

NZCYM Agar

Bacto Casein Digest	10 g
Bacto Casamino Acids	1 g
Bacto Yeast Extract	5 g
Sodium Chloride	5 g

Magnesium Sulfate 0.98 g

(anhydrous)

Agar	15 g
Distilled Water	1,000 ml

NZCYM Top Agarose

Bacto Casein Digest	10 g
Bacto Casamino Acids	1 g
Bacto Yeast Extract	5 g
Sodium Chloride	5 g

Magnesium Sulfate 0.98 g

(anhydrous)

Agarose	7 g
Distilled Water	1,000 ml

YEPD Broth

Bacto Yeast Extract	10 g
Bacto Peptone	20 g
Glucose	20 g
Distilled Water	1,000 ml

YEPD Agar*

Bacto Yeast Extract	10 g
Bacto Peptone	20 g
Glucose	20 g
Agar	20 g
Distilled Water	1,000 ml

SC - uracil*

Bacto-yeast nitrogen base without amino acids	6.7g
Glucose	20g
Bacto-agar	20g
Drop-out mix	2g
Distilled water	1,000ml

<u>DCA2 medium</u>	g/l
Peptone	3.0
Yeast Extract	6.0
Sodium Acetate	3.0
Yeast Nitrogen Base (Difco)	6.7
Glucose (anhydrous)	50.0
Potassium Phosphate (dibasic, trihydrate)	7.2
Potassium Phosphate (monobasic, anhydrous)	9.3

<u>DCA3 medium</u>	g/l
0.3 M Phosphate buffer containing, pH 7.5	
Glycerol	50
Yeast Nitrogen base (Difco)	6.7

Drop-out mix

Adenine	0.5g	Alanine	2g
Arginine	2g	Asparagine	2g
Aspartic acid	2g	Cysteine	2g
Glutamine	2g	Glutamic acid	2g
Glycine	2g	Histidine	2g
Inositol	2g	Isoleucine	2g
Leucine	10g	Lysine	2g
Methionine	2g	para-Aminobenzoic acid	0.2g
Phenylalanine	2g	Proline	2g
Serine	2g	Threonine	2g
Tryptophan	2g	Tyrosine	2g
Valine	2g		

*See Kaiser et al., Methods in Yeast Genetics, Cold Spring Harbor Laboratory Press, USA (1994), incorporated herein by reference.

It will be understood that various modifications may be made to the embodiments and/or examples disclosed herein. Thus, the above description should not be construed as limiting, but merely as exemplifications of preferred embodiments. Those skilled in the art will envision other modifications within the scope and spirit of the claims appended hereto.

100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000

SEQUENCE LISTING

<110> Wilson, Ron C.
 Craft, David L.
 Eirich, Dudley
 Eshoo, Mark
 Madduri, Krishna M.
 Cornett, Cathy A.
 Brenner, Alfred A.
 Tang, Maria
 Loper, John C.
 Gleeson, Martin

<120> CYTOCHROME P450 MONOOXYGENASE AND NADPH CYTOCHROME P450 OXIDOREDUCTASE
 GENES AND PROTEINS RELATED TO THE OMEGA HYDROXYLASE COMPLEX OF CANDIDA
 TROPICALIS AND METHODS RELATING THERETO

<130> 1010-16

<140> US 09/302,602

<141> 1999-04-30

<160> 118

<170> PatentIn version 3.1

<210> 1

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 1

ccttaattaa atgcacgaag cggagataaa ag 32

<210> 2

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 2

ccttaattaa gcataagctt gctcgagtct 30

<210> 3

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 3

ccttaattaa acgcaatggg aacatggagt g 31

<210> 4
 <211> 34
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 4
 ccttaattaa tcgcactacg gttattggta tcag 34

 <210> 5
 <211> 29
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 5
 ccttaattaa tcaaagtcg ttcaggcgg 29

 <210> 6
 <211> 34
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 6
 ccttaattaa ggcagacaac aacttgcaa agtc 34

 <210> 7
 <211> 31
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 7
 ccttaattaa gaggtcgttg gttgagtttt c 31

 <210> 8
 <211> 29
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 8
 ccttaattaa ttgataatga cgttgcggg 29

<210> 9
 <211> 33
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 9
 aggcgcgcgcg gagtccaaaa agaccaacct ctg 33

<210> 10
 <211> 34
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 10
 ccttaattaa tacgtggata ccttcaagca agtg 34

<210> 11
 <211> 35
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 11
 ccttaattaa gctcacgagt tttgggattt tcgag 35

<210> 12
 <211> 35
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 12
 gggtttaaac cgcagagggtt ggtctttttg gactc 35

<210> 13
 <211> 10
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 13
 gggtttaaac 10

<210> 14
<211> 9
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 14
aggcgcgcc

9

<210> 15
<211> 10
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 15
ccttaattaa

10

<210> 16
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<220>
<221> misc_feature
<222> (3)..(4)
<223> y=dCTP or dTTP

<220>
<221> misc_feature
<222> (9)..(10)
<223> w=dATP or dTTP

<220>
<221> misc_feature
<222> (15)..(16)
<223> w=dATP or dTTP

<220>
<221> misc_feature
<222> (18)..(19)
<223> w=dATP or dTTP

<400> 16
tcycaaacwg gtacgwcwga a

21

<210> 17
<211> 21
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<220>

<221> misc_feature

<222> (12)..(13)

<223> y=dCTP or dTTP

<220>

<221> misc_feature

<222> (15)..(16)

<223> w=dATP or dTTP

<400> 17

ggtttgggta aytcwactta t

21

<210> 18

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 18

cgttattatc atttcttc

18

<210> 19

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<220>

<221> misc_feature

<222> (3)..(4)

<223> m=dATP or dCTP

<220>

<221> misc_feature

<222> (9)..(10)

<223> r=dATP or dGTP

<400> 19

gcmacaccrg tacctggacc

20

<210> 20
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 20
atcccaatcg taatcagc

18

<210> 21
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 21
acttgctcttc gtttagca

18

<210> 22
<211> 18
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 22
ctacgtctgt ggtgatgc

18

<210> 23
<211> 17
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<220>

<221> misc_feature

<222> (3)..(4)

<223> n=dATP or dCTP or dGTP or dTTP

<220>

<221> misc_feature

<222> (6)..(7)

<223> Y=dCTP or dTTP

<220>

<221> misc_feature

<222> (9)..(10)
<223> n=dATP or dCTP or dGTP or dTTP

<220>
<221> misc_feature
<222> (12)..(13)
<223> n=dATP or dCTP or dGTP or dTTP

<220>
<221> misc_feature
<222> (15)..(16)
<223> n=dATP or dCTP or dGTP or dTTP

<400> 23
cnggayacna cngcngg

17

<210> 24
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<220>
<221> misc_feature
<222> (3)..(4)
<223> r=dATP or dTTP

<220>
<221> misc_feature
<222> (6)..(7)
<223> y=dCTP or dTTP

<220>
<221> misc_feature
<222> (9)..(10)
<223> n=dATP or dCTP or dGTP or dTTP

<220>
<221> misc_feature
<222> (12)..(13)
<223> n=dATP or dCTP or dGTP or dTTP

<220>
<221> misc_feature
<222> (15)..(16)
<223> n=dATP or dCTP or dGTP or dTTP

<400> 24
agrgayacna cngcngg

17

<210> 25
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<220>
<221> misc_feature
<222> (3)..(4)
<223> n=dATP or dCTP or dGTP or dTTP

<220>
<221> misc_feature
<222> (6)..(7)
<223> r=dATP or dGTP

<220>
<221> misc_feature
<222> (9)..(10)
<223> y=dCTP or dTTP

<220>
<221> misc_feature
<222> (12)..(13)
<223> y=dCTP or dTTP

<220>
<221> misc_feature
<222> (15)..(16)
<223> n=dATP or dCTP or dGTP or dTTP

<400> 25
agngcraayt gytgncc

17

<210> 26
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<220>
<221> misc_feature
<222> (1)..(2)
<223> y=dCTP or dTTP

<220>
<221> misc_feature

<222> (4)..(5)
<223> n=dATP or dCTP or dGTP or dTTP

<220>
<221> misc_feature
<222> (7)..(8)
<223> r=dATP or dGTP

<220>
<221> misc_feature
<222> (10)..(11)
<223> y=dCTP or dTTP

<220>
<221> misc_feature
<222> (13)..(14)
<223> y=dCTP or dTTP

<220>
<221> misc_feature
<222> (16)..(17)
<223> n=dATP or dCTP or dGTP or dTTP

<400> 26
yaangcraay tgytgncc

18

<210> 27
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 27
attcaacggt ggtccaagaa tctgtttgg

29

<210> 28
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 28
gagctatggt gagaccacag ttgac

25

<210> 29
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

 <400> 29
 cttcagttaa agcaaattgt ttggcc 26

 <210> 30
 <211> 25
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 30
 ctcggaagc gcgccattgt gttgg 25

 <210> 31
 <211> 29
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <400> 31
 taatacgact cactataggg cgaattggc 29

 <210> 32
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Primer

 <220>
 <221> misc_feature
 <222> (3)..(4)
 <223> r=dATP,or dGTP

 <220>
 <221> misc_feature
 <222> (4)..(5)
 <223> y=dCTP or dTTP

 <220>
 <221> misc_feature
 <222> (16)..(17)
 <223> y=dCTP or dTTP

 <400> 32
 tgrytcaaac catctytctg g 21

<210> 33
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 33
 ggaccggcgt taaagg 17

<210> 34
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 34
 catagtcgwa tyatgcttag acc 23

<210> 35
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Primer

<400> 35
 ggaccacat tgaatgg 17

<210> 36
 <211> 540
 <212> DNA
 <213> Candida tropicalis

<400> 36
 atgattgaac aactcctaga atattggtat gtcgttgtgc cagtgttgta catcatcaaa 60
 caactccttg catacacaaa gactcgcgtc ttgatgaaaa agttgggtgc tgctccagtc 120
 acaaacaagt tgtacgacaa cgctttcggg atcgtcaatg gatggaaggc tctccagttc 180
 aagaaagagg gcagggctca agagtacaac gattacaagt ttgaccactc caagaaccca 240
 agcgtgggca cctacgtcag tattcttttc ggcaccagga tcgtcgtgac caaagatcca 300
 gagaatatca aagctatattt ggcaaccag tttggtgatt tttctttggg caagaggcac 360
 actcttttta agcctttggt aggtgatggg atcttcacat tggacggcga aggttggaag 420
 cacagcagag ccatgttgag accacagttt gccagagaac aagttgctca tgtgacgtcg 480

ttggaaccac acttccagtt gttgaagaag catattctta agcacaaggg tgaatacttt 540

<210> 37
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 37
ccgatgaagt tttcgacgag taccc 25

<210> 38
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 38
aaggctttaa cgtgtccaat ctggtc 26

<210> 39
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 39
attatcgcca catacttcac caaatgg 27

<210> 40
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 40
cgagatcgtg gatacgctgg agtg 24

<210> 41
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 41

gccactcggg aactttgtca gggac

25

<210> 42

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 42

cattgaactg agtagccaaa acagcc

26

<210> 43

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 43

cctacgtttg gtatcgctac tccgttg

27

<210> 44

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 44

tttccagcca gcaccgtcca ag

22

<210> 45

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 45

gcagagccga tctatgttgc gtcc

24

<210> 46

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 46

tcattgaatg cttccaggaa cctcg 25

<210> 47
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 47
aagagggcag ggctcaagag 20

<210> 48
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 48
tccatgtgaa gatcccatca c 21

<210> 49
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 49
cttgaaggcc gtggttgaacg 20

<210> 50
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 50
caggatttgt ctgagttgcc g 21

<210> 51
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 51

ccattgcctt gagatacgcc attggtag

28

<210> 52

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 52

agccttggtg tcgttctttt caacgg

26

<210> 53

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 53

ttgggtttgt ttgtttcctg tgtccg

26

<210> 54

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 54

cctttgacct tcaatctggc gtagacg

27

<210> 55

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 55

gtttgctgaa tacgctgaag gtgatg

26

<210> 56

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 56

tgagctgaa caactctctc gtctcgg

27

<210> 57

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 57

ttcctcaaca cggacagcgg

20

<210> 58

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 58

agtcaaccag gtgtggaact cgtc

24

<210> 59

<211> 49

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 59

ggatcctaatac gactcact atagggagga agagggcagg gctcaagag

49

<210> 60

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 60

tccatgtgaa gatcccatca cgagtgtgcc tcttgcccaa ag

42

<210> 61

<211> 54

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 61

ggatcctaatac gactcact ataggaggc cgatgaagtt ttcgacgagt accc 54

<210> 62
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 62
aaggctttaa cgtgtccaat ctggtcaaca tagctctgga gtgcttccaa cc 52

<210> 63
<211> 56
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 63
ggatcctaatac gactcact ataggaggga ttatgccac atacttcacc aaatgg 56

<210> 64
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 64
cgagatcgtg gatacgctgg agtgcgtcgc tcttcttctt caacaattca ag 52

<210> 65
<211> 49
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 65
cattgaactg agtagccaaa acagcccatg gtttcaatca atgggaggc 49

<210> 66
<211> 54
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 66

ggatccta at acgactcact atagggaggg ccactcggtta actttgtcag ggac 54

<210> 67
<211> 56
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 67
ggatccta at acgactcact atagggaggg ctacgtttgg tatcgctact ccgttg 56

<210> 68
<211> 48
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 68
tttcagcca gcaccgtcca agcaacaagg agtacaagaa atcgtgtc 48

<210> 69
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 69
ggatccta at acgactcact atagggaggg cagagccgat ctatgttgcg tcc 53

<210> 70
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 70
tcattgaatg cttccaggaa cctcgccaca tccatcgaga accgg 45

<210> 71
<211> 49
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 71

ggatcctaatac gactcact ataggaggc ttgaaggccg tgttgaacg 49

<210> 72
<211> 46
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 72
caggatttgt ctgagttgcc gcctgatcaa gataggatcc ttgccg 46

<210> 73
<211> 56
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 73
ggatcctaatac gactcact ataggagggt gtttgctgaa tacgctgaag gtgatg 56

<210> 74
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 74
tggagctgaa caactctctc gtctcgggtg gtcgaatgga cccttggtca ag 52

<210> 75
<211> 49
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 75
ggatcctaatac gactcact ataggagggt tcctcaacac ggacagcgg 49

<210> 76
<211> 49
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 76

agtcaaccag gtgtggaact cgtcgggtggc aacaatgaaa aacaccaag 49

<210> 77
<211> 57
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 77
ggatcctaatac gactcact ataggagggc cattgccttg agatacgcca ttggtag 57

<210> 78
<211> 53
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 78
agccttggtg tcgttctttt caacggaagg tggctcgtat ggtgtgttca acc 53

<210> 79
<211> 55
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 79
ggatcctaatac gactcact ataggagggt tgggtttggt tgtttcctgt gtccg 55

<210> 80
<211> 50
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 80
cctttgacct tcaatctggc gtagacgcag caccaccgat ccaccacttg 50

<210> 81
<211> 4206
<212> DNA
<213> Candida tropicalis

<400> 81
catcaagatc atctatgggg ataattacga cagcaacatt gcagaaagag cgttgggtcac 60

aatcgaaaga gcctatggcg ttgccgtcgt tgaggcaaata gacagcacca acaataacga 120

tggtcccagt	gaagagcctt	cagaacagtc	cattgttgac	gcttaaggca	cggataatta	180
cgtggggcaa	aggaacgcgg	aattagttat	ggggggatca	aaagcggaag	atttgtgttg	240
cttgtgggtt	ttttccttta	tttttcatat	gatttccttg	cgcaagtaac	atgtgccaat	300
ttagtttgtg	attagcgtgc	cccacaattg	gcatcgtgga	cgggcgtgtt	ttgtcatacc	360
ccaagtctta	actagctcca	cagtctcgac	ggtgtctcga	cgatgtcttc	ttccaccctt	420
cccatgaatc	attcaaagtt	gttgggggat	ctccaccaag	ggcaccggag	ttaatgctta	480
tgtttctccc	actttgggtg	tgattggggg	agtctagtga	gttgagatt	ttcttttttt	540
cgcaggtgtc	tccgatatcg	aaatttgatg	aatatagaga	gaagccagat	cagcacagta	600
gattgccttt	gtagttagag	atgttgaaca	gcaactagtt	gaattacacg	ccaccacttg	660
acagcaagtg	cagtgaagctg	taaacgatgc	agccagagtg	tcaccaccaa	ctgacgttgg	720
gtggagtgtg	tggtgttggt	gttggcaggg	ccatattgct	aaacgaagac	aagtagcaca	780
aaacccaagc	ttaagaacaa	aaataaaaaa	aattcatacg	acaattccaa	agccattgat	840
ttacataatc	aacagtaaga	cagaaaaaac	tttcaacatt	tcaaagttcc	ctttttccta	900
ttacttcttt	tttttcttct	ttccttcttt	ccttctgttt	ttcttacttt	atcagtcttt	960
tacttgtttt	tgcaattcct	catcctcttc	ctactcctcc	tcaccatggc	tttagacaag	1020
ttagatttgt	atgtcatcat	aacattgggtg	gtcgctgtag	ccgcctattt	tgctaagaac	1080
cagttccttg	atcagcccca	ggacaccggg	ttcctcaaca	cggacagcgg	aagcaactcc	1140
agagacgtct	tgctgacatt	gaagaagaat	aataaaaaca	cggtgttggt	gtttgggtcc	1200
cagacgggta	cggcagaaga	ttacgccaac	aaattgtcca	gagaattgca	ctccagattt	1260
ggcttgaaaa	cgatgggtgc	agatttcgct	gattacgatt	gggataactt	cggagatatc	1320
accgaagaca	tcttggtgtt	tttcattgtt	gccacctatg	gtgagggtga	acctaccgat	1380
aatgccgacg	agttccacac	ctggttgact	gaagaagctg	acactttgag	taccttgaaa	1440
tacaccgtgt	tcgggttggg	taactccacg	tacgagttct	tcaatgccat	tggtagaaag	1500
tttgacagat	tggtgagcga	gaaagggtgt	gacaggtttg	ctgaatacgc	tgaagggtgat	1560
gacgggtactg	gcaccttgga	cgaagatttc	atggcctgga	aggacaatgt	ctttgacgcc	1620
ttgaagaatg	atttgaactt	tgaagaaaag	gaattgaagt	acgaaccaa	cgtgaaattg	1680
actgagagag	acgacttgct	tgctgctgac	tcccaagttt	ccttggtgga	gccaaacaag	1740
aagtacatca	actccgaggg	catcgacttg	accaagggtc	cattcgacca	caccoaccca	1800
tacttgcca	gaatcaccca	gacgagagag	ttgttcagct	ccaaggacag	acactgtatc	1860
cacgttgaat	ttgacatttc	tgaatcgaac	ttgaaataca	ccaccggtga	ccatctagct	1920

atctggccat ccaactccga cgaaaacatt aagcaatttg ccaagtgttt cggattggaa 1980
 gataaactcg aactgttat tgaattgaag gcgttggact ccacttacac catcccattc 2040
 ccaaccccaa ttacctacgg tgctgtcatt agacaccatt tagaaatctc cggtcacgtc 2100
 tcgagacaat tctttttgtc aattgctggg tttgctcctg atgaagaaac aaagaaggct 2160
 tttaccagac ttggtggtga caagcaagaa ttgcgcgcca aggtcaccgc cagaaagttc 2220
 aacattgccg atgccttggt atattcctcc aacaacgctc catggtccga tgttcctttt 2280
 gaattcctta ttgaaaacgt tccacacttg actccacgtt actactccat ttcgtcttcg 2340
 tcattgagtg aaaagcaact catcaacgtt actgcagttg ttgaagccga agaagaagct 2400
 gatggcagac cagtcactgg tgtgtcacc aacttgttga agaacgttga aattgtgcaa 2460
 aacaagactg gcgaaaagcc acttgtccac tacgatttga ggggcccaag aggcaagttc 2520
 aacaagttca agttgccagt gcatgtgaga agatccaact ttaagttgcc aaagaactcc 2580
 accaccccag ttatcttgat tggtcacagt actggtgttg cccattgag aggttttgtc 2640
 agagaaagag ttcaacaagt caagaatggt gtcaatgttg gcaagacttt gttgttttat 2700
 gggtgcagaa actccaacga ggactttttg tacaagcaag aatgggccga gtacgcttct 2760
 gttttgggtg aaaactttga gatgttcaat gccttctcca gacaagacc atccaagaag 2820
 gtttacgtcc aggataagat tttagaaaac agccaacttg tgcacgagtt gttgactgaa 2880
 ggtgccatta tctacgtctg tggtagtgcc agtagaatgg ctagagacgt gcagaccaca 2940
 atttccaaga ttgttgctaa aagcagagaa attagtgaag acaaggctgc tgaattggtc 3000
 aagtcctgga aggtccaaaa tagataccaa gaagatgttt ggtagactca aacgaatctc 3060
 tctttctccc aacgcattta tgaatcttta ttctcattga agctttacat atgttctaca 3120
 ctttattttt tttttttttt ttattattat attacgaaac ataggtcaac tatatatact 3180
 tgattaaatg ttatagaaac aataactatt atctactcgt ctacttcttt ggcattgaca 3240
 tcaacattac cgttccatt accgttgccg ttggcaatgc cgggatattt agtacagtat 3300
 ctccaatccg gatttgagct attgtagatc agctgcaagt cattctccac cttcaaccag 3360
 tacttatact tcatctttga cttcaagtcc aagtcataaa tattacaagt tagcaagaac 3420
 ttctggccat ccacgatata gacgttattc acgttattat gcgacgtatg gatgtggtta 3480
 tccttattga acttctcaaa cttcaaaaac aacccacgt cccgcaacgt cattatcaac 3540
 gacaagttct ggctcacgtc gtcggagctc gtcaagttct caattagatc gttcttgtaa 3600
 ttgatcttct ggtactttct caattgctgg aacacattgt cctcgttggt caaatagatc 3660
 ttgaacaact ttttcaacgg gatcaacttc tcaatctggg ccaagatctc cgccgggatac 3720

ttcagaaaca agtcctgcaa cccctggtcg atggtctccg ggtacaacaa gtccaagggg	3780
cagaagtgtc taggcacgtg tttcaactgg ttcaacgaac atgttcgaca gtagttcgag	3840
ttatagttat cgtacaacca ttttggtttg atttcgaaaa tgacggagct gatgccatca	3900
ttctcctggt tcctctcata gtacaactgg cacttcttcg agaggctcaa ttctcgtag	3960
ttcccgcca agatattcgg caacaagagc ccgtaccgct cacggagcat caagtctggtg	4020
ccctggttgt tcaacttggt gatgaagtc gaggtcaaga caatcaactg gatgtcgatg	4080
atctggtgcg ggaacaagtt cttgcatttt agctcgatga agtcgtacaa ctcacacgtc	4140
gagatatact cctgttcttc cttcaagagc cggatccgca agagcttgtg cttcaagtag	4200
tcgttg	4206

<210> 82
 <211> 4145
 <212> DNA
 <213> *Candida tropicalis*

<400> 82	
tatatgat atgatatac ttctgtgta attattatc gtattcgta atacttacta	60
catttttttt tctttattta tgaagaaaag gagagtctgt aagttgagtt gagtagaata	120
ggctgttgtg catacgggga gcagaggaga gtatccgacg aggaggaaact ggggtgaaatt	180
tcctctatgc tgttgcgctc tgtactgtac tgtaaattct agatttccta gaggttggtc	240
tagcaaataa agtgtttcaa gatacaattt tacaggcaag ggtaaaggat caactgatta	300
gcggaagatt ggtgttgctt gtgggggtct tttatttttc atatgatttc tttgcgcgag	360
taacatgtgc caatctagtt tatgattagc gtacctccac aattggcatc ttggacgggc	420
gtgttttgtc ttacccaag cttatttag ttccacagtc tcgacggtgt ctgcgcgatg	480
tcttctccca cccctcgcag gaatcattcg aagttgttg gggatctcct ccgcagtta	540
tgttcatgtc tttccactt tgggtgtgat tggggtagcg tagtgagttg gtgattttct	600
tttttcgcag gtgtctccga tatcgaagtt tgatgaatat aggagccaga tcagcatggt	660
atattgcctt tgtagataga gatgttgaac aacaactagc tgaattacac accaccgcta	720
aacgatgctc acaggggtgc accgccaact gacgttgggt ggagttgttg ttggcagggc	780
catattgcta aacgaagaga agtagacaa aaccaagggt taagaacaat taaaaaaatt	840
catacgacaa ttccacagcc atttacataa tcaacagcga caaatgagac agaaaaaact	900
ttcaacattt caaggttccc tttttctat tacttctttt tttctttcct tcctttcatt	960
tcctttcctt ctgcttttat tactttacca gtcttttgct tgtttttgca attcctcatc	1020

ctccctcctca ccatggcctt agacaagtta gatttgtatg tcatcataac attggtggtc 1080
 gctgtggccg cctatcttgc taagaaccag ttccttgatc agccccagga caccgggttc 1140
 ctcaacacgg acagcggaag caactccaga gacgtcttgc tgacattgaa gaagaataat 1200
 aaaaacacgt tgttgttgtt tgggtcccag accggtacgg cagaagatta cgccaacaaa 1260
 ttgtcaagag aattgcactc cagatttggc ttgaaaacca tggttgcaga tttcgtgat 1320
 tacgattggg ataacttcgg agatatcacc gaagatatct tgggtgtttt catcgttgcc 1380
 acctacggtg aggggtgaacc taccgacaat gccgacgagt tccacacctg gttgactgaa 1440
 gaagctgaca ctttgagtac tttgagatat accgtgttcg ggttgggtaa ctccacctac 1500
 gagttcttca atgctattgg tagaaaagtt gacagattgt tgagtgagaa aggtggtgac 1560
 agatttgctg aatatgctga aggtgacgac ggcactggca ccttgagcga agatttcatg 1620
 gcctggaagg ataattgtct tgacgccttg aagaatgact tgaactttga agaaaaggaa 1680
 ttgaagtacg aaccaaactg gaaattgact gagagagatg acttgtctgc tgccgactcc 1740
 caagtttctt tgggtgagcc aaacaagaag tacatcaact ccgagggcat cgacttgacc 1800
 aaggttccat tcgaccacac ccaccatac ttggccagga tcacogagac cagagagttg 1860
 ttcagctcca aggaaagaca ctgtattcac gttgaatttg acatttctga atcgaacttg 1920
 aaatacacca ccggtgacca tctagccatc tggccatcca actccgacga aaacatcaag 1980
 caatttgcca agtgtttcgg attggaagat aaactcgaca ctgttattga attgaaggca 2040
 ttggactcca cttacaccat tccattccca actccaatta cttacgggtg tgctattaga 2100
 caccatttag aaatctccgg tccagtctcg agacaattct ttttgtcgat tgctgggttt 2160
 gtcctgatg aagaaacaaa gaagacttcc accagacttg gtggtgacaa acaagaattc 2220
 gccaccaagg ttaccgcgag aaagttcaac attgccgatg ccttggtata ttcctccaac 2280
 aacactccat ggtccgatgt tccttttgag ttccttattg aaaacatcca aacttgact 2340
 ccacgttact actccatttc tcttcgtcg ttgagtgaac aacaactcat caatgttact 2400
 gcagtcgttg aggccgaaga agaagccgat ggcagaccag tcaactggtg tgttaccaac 2460
 ttgttgaaga acattgaaat tgcgcaaac aagactggcg aaaagccact tgttactac 2520
 gatttgagcg gcccaagagg caagttcaac aagttcaagt tgccagtga cgtgagaaga 2580
 tccaacttta agttgccaaa gaactccacc accccagtta tcttgattgg tccaggtact 2640
 ggtgttgccc cattgagagg tttcgttaga gaaagagttc aacaagtcaa gaatggtgtc 2700
 aatgttgcca agactttgtt gttttatggt tgcagaaact ccaacgagga cttttgtac 2760
 aagcaagaat gggccgagta cgcttctgtt ttgggtgaaa actttgagat gttcaatgcc 2820

ttctctagac aagacccatc caagaaggtt tacgtccagg ataagatttt agaaaacagc 2880
 caacttgtgc acgaattggt gaccgaaggt gccattatct acgtctgtgg tgacgccagt 2940
 agaatggcca gagacgtcca gaccacgata tccaagattg ttgccaaaag cagagaaatc 3000
 agtgaagaca aggccgctga attggtcaag tcttggaag tccaaaatag ataccaagaa 3060
 gatgtttggt agactcaaac gaatctctct ttctcccaac gcatttatga atattctcat 3120
 tgaagtttta catatgttct atatttcatt tttttttat tatattacga aacataggtc 3180
 aactatatat acttgattaa atgttataga aacaataatt attatctact cgtctacttc 3240
 tttggcattg gcattggcat tggcattggc attgccgttg ccgttggtaa tgccgggata 3300
 tttagtacag tatctccaat ccggatttga gctattgtaa atcagctgca agtcattctc 3360
 caccttcaac cagtacttat acttcatctt tgacttcaag tccaagtcac aatatattaca 3420
 agttagcaag aacttctggc catccacaat atagacgtta ttcacgttat tatgcgacgt 3480
 atggatatgg ttatccttat tgaacttctc aaacttcaa aacaaccca cgtcccga 3540
 cgtcattatc aacgacaagt tctgactcac gtctcgagg ctcgtcaagt tctcaattag 3600
 atcgttcttg ttattgatct tctggtactt tctcaactgc tggaacacat tgtcctcgtt 3660
 gttcaaatag atcttgaaca acttcttcaa gggaatcaac ttttcgatct gggccaagat 3720
 ttccgcgggg atcttcagaa acaagtctg caaccctgg tcgatggtct cggggtagaa 3780
 caagtctaag gggcagaagt gtctaggcac gtgtttcaac tgggtcaagg aacatgttcg 3840
 acagtagttc gagttatagt tatcgtacaa ccactttggc ttgatttcga aatgacgga 3900
 gctgatccca tcattctcct ggttcctttc atagtacaac tggcatttct tcgagagact 3960
 caactcctcg tagttcccg ccaagatatt cggcaacaag agcccgtagc gtcacggag 4020
 catcaagtgc tggccctggt tgttcaactt gttgatgaag tccgatgtca agacaatcaa 4080
 ctggatgtcg atgatctggt gcggaacaa gttcttgca tttagctcga tgaagtcgta 4140
 caact 4145

<210> 83
 <211> 679
 <212> PRT
 <213> CANDIDATROPICALIS

<400> 83

Met Ala Leu Asp Lys Leu Asp Leu Tyr Val Ile Ile Thr Leu Val Val
 1 5 10 15

Ala Val Ala Ala Tyr Phe Ala Lys Asn Gln Phe Leu Asp Gln Pro Gln
 20 25 30

Asp Thr Gly Phe Leu Asn Thr Asp Ser Gly Ser Asn Ser Arg Asp Val
 35 40 45

Leu Leu Thr Leu Lys Lys Asn Asn Lys Asn Thr Leu Leu Leu Phe Gly
 50 55 60

Ser Gln Thr Gly Thr Ala Glu Asp Tyr Ala Asn Lys Leu Ser Arg Glu
 65 70 75 80

Leu His Ser Arg Phe Gly Leu Lys Thr Met Val Ala Asp Phe Ala Asp
 85 90 95

Tyr Asp Trp Asp Asn Phe Gly Asp Ile Thr Glu Asp Ile Leu Val Phe
 100 105 110

Phe Ile Val Ala Thr Tyr Gly Glu Gly Glu Pro Thr Asp Asn Ala Asp
 115 120 125

Glu Phe His Thr Trp Leu Thr Glu Glu Ala Asp Thr Leu Ser Thr Leu
 130 135 140

Lys Tyr Thr Val Phe Gly Leu Gly Asn Ser Thr Tyr Glu Phe Phe Asn
 145 150 155 160

Ala Ile Gly Arg Lys Phe Asp Arg Leu Leu Ser Glu Lys Gly Gly Asp
 165 170 175

Arg Phe Ala Glu Tyr Ala Glu Gly Asp Asp Gly Thr Gly Thr Leu Asp
 180 185 190

Glu Asp Phe Met Ala Trp Lys Asp Asn Val Phe Asp Ala Leu Lys Asn
 195 200 205

Asp Leu Asn Phe Glu Glu Lys Glu Leu Lys Tyr Glu Pro Asn Val Lys
 210 215 220

Leu Thr Glu Arg Asp Asp Leu Ser Ala Ala Asp Ser Gln Val Ser Leu
 225 230 235 240

Gly Glu Pro Asn Lys Lys Tyr Ile Asn Ser Glu Gly Ile Asp Leu Thr
 245 250 255

Lys Gly Pro Phe Asp His Thr His Pro Tyr Leu Ala Arg Ile Thr Glu
 260 265 270

Thr Arg Glu Leu Phe Ser Ser Lys Asp Arg His Cys Ile His Val Glu
275 280 285

Phe Asp Ile Ser Glu Ser Asn Leu Lys Tyr Thr Thr Gly Asp His Leu
290 295 300

Ala Ile Trp Pro Ser Asn Ser Asp Glu Asn Ile Lys Gln Phe Ala Lys
305 310 315 320

Cys Phe Gly Leu Glu Asp Lys Leu Asp Thr Val Ile Glu Leu Lys Ala
325 330 335

Leu Asp Ser Thr Tyr Thr Ile Pro Phe Pro Thr Pro Ile Thr Tyr Gly
340 345 350

Ala Val Ile Arg His His Leu Glu Ile Ser Gly Pro Val Ser Arg Gln
355 360 365

Phe Phe Leu Ser Ile Ala Gly Phe Ala Pro Asp Glu Glu Thr Lys Lys
370 375 380

Ala Phe Thr Arg Leu Gly Gly Asp Lys Gln Glu Phe Ala Ala Lys Val
385 390 395 400

Thr Arg Arg Lys Phe Asn Ile Ala Asp Ala Leu Leu Tyr Ser Ser Asn
405 410 415

Asn Ala Pro Trp Ser Asp Val Pro Phe Glu Phe Leu Ile Glu Asn Val
420 425 430

Pro His Leu Thr Pro Arg Tyr Tyr Ser Ile Ser Ser Ser Ser Leu Ser
435 440 445

Glu Lys Gln Leu Ile Asn Val Thr Ala Val Val Glu Ala Glu Glu Glu
450 455 460

Ala Asp Gly Arg Pro Val Thr Gly Val Val Thr Asn Leu Leu Lys Asn
465 470 475 480

Val Glu Ile Val Gln Asn Lys Thr Gly Glu Lys Pro Leu Val His Tyr
485 490 495

Asp Leu Ser Gly Pro Arg Gly Lys Phe Asn Lys Phe Lys Leu Pro Val
500 505 510

His Val Arg Arg Ser Asn Phe Lys Leu Pro Lys Asn Ser Thr Thr Pro
515 520 525

Val Ile Leu Ile Gly Pro Gly Thr Gly Val Ala Pro Leu Arg Gly Phe
530 535 540

Val Arg Glu Arg Val Gln Gln Val Lys Asn Gly Val Asn Val Gly Lys
545 550 555 560

Thr Leu Leu Phe Tyr Gly Cys Arg Asn Ser Asn Glu Asp Phe Leu Tyr
565 570 575

Lys Gln Glu Trp Ala Glu Tyr Ala Ser Val Leu Gly Glu Asn Phe Glu
580 585 590

Met Phe Asn Ala Phe Ser Arg Gln Asp Pro Ser Lys Lys Val Tyr Val
595 600 605

Gln Asp Lys Ile Leu Glu Asn Ser Gln Leu Val His Glu Leu Leu Thr
610 615 620

Glu Gly Ala Ile Ile Tyr Val Cys Gly Asp Ala Ser Arg Met Ala Arg
625 630 635 640

Asp Val Gln Thr Thr Ile Ser Lys Ile Val Ala Lys Ser Arg Glu Ile
645 650 655

Ser Glu Asp Lys Ala Ala Glu Leu Val Lys Ser Trp Lys Val Gln Asn
660 665 670

Arg Tyr Gln Glu Asp Val Trp
675

<210> 84
<211> 679
<212> PRT
<213> CANDIDATROPICALIS

<400> 84

Met Ala Leu Asp Lys Leu Asp Leu Tyr Val Ile Ile Thr Leu Val Val
1 5 10 15

Ala Val Ala Ala Tyr Phe Ala Lys Asn Gln Phe Leu Asp Gln Pro Gln
20 25 30

Asp Thr Gly Phe Leu Asn Thr Asp Ser Gly Ser Asn Ser Arg Asp Val
 35 40 45
 Leu Leu Thr Leu Lys Lys Asn Asn Lys Asn Thr Leu Leu Leu Phe Gly
 50 55 60
 Ser Gln Thr Gly Thr Ala Glu Asp Tyr Ala Asn Lys Leu Ser Arg Glu
 65 70 75 80
 Leu His Ser Arg Phe Gly Leu Lys Thr Met Val Ala Asp Phe Ala Asp
 85 90 95
 Tyr Asp Trp Asp Asn Phe Gly Asp Ile Thr Glu Asp Ile Leu Val Phe
 100 105 110
 Phe Ile Val Ala Thr Tyr Gly Glu Gly Glu Pro Thr Asp Asn Ala Asp
 115 120 125
 Glu Phe His Thr Trp Leu Thr Glu Glu Ala Asp Thr Leu Ser Thr Leu
 130 135 140
 Arg Tyr Thr Val Phe Gly Leu Gly Asn Ser Thr Tyr Glu Phe Phe Asn
 145 150 155 160
 Ala Ile Gly Arg Lys Phe Asp Arg Leu Leu Ser Glu Lys Gly Gly Asp
 165 170 175
 Arg Phe Ala Glu Tyr Ala Glu Gly Asp Asp Gly Thr Gly Thr Leu Asp
 180 185 190
 Glu Asp Phe Met Ala Trp Lys Asp Asn Val Phe Asp Ala Leu Lys Asn
 195 200 205
 Asp Leu Asn Phe Glu Glu Lys Glu Leu Lys Tyr Glu Pro Asn Val Lys
 210 215 220
 Leu Thr Glu Arg Asp Asp Leu Ser Ala Ala Asp Ser Gln Val Ser Leu
 225 230 235 240
 Gly Glu Pro Asn Lys Lys Tyr Ile Asn Ser Glu Gly Ile Asp Leu Thr
 245 250 255
 Lys Gly Pro Phe Asp His Thr His Pro Tyr Leu Ala Arg Ile Thr Glu
 260 265 270

Thr Arg Glu Leu Phe Ser Ser Lys Glu Arg His Cys Ile His Val Glu
 275 280 285

Phe Asp Ile Ser Glu Ser Asn Leu Lys Tyr Thr Thr Gly Asp His Leu
 290 295 300

Ala Ile Trp Pro Ser Asn Ser Asp Glu Asn Ile Lys Gln Phe Ala Lys
 305 310 315 320

Cys Phe Gly Leu Glu Asp Lys Leu Asp Thr Val Ile Glu Leu Lys Ala
 325 330 335

Leu Asp Ser Thr Tyr Thr Ile Pro Phe Pro Thr Pro Ile Thr Tyr Gly
 340 345 350

Ala Val Ile Arg His His Leu Glu Ile Ser Gly Pro Val Ser Arg Gln
 355 360 365

Phe Phe Leu Ser Ile Ala Gly Phe Ala Pro Asp Glu Glu Thr Lys Lys
 370 375 380

Thr Phe Thr Arg Leu Gly Gly Asp Lys Gln Glu Phe Ala Thr Lys Val
 385 390 395 400

Thr Arg Arg Lys Phe Asn Ile Ala Asp Ala Leu Leu Tyr Ser Ser Asn
 405 410 415

Asn Thr Pro Trp Ser Asp Val Pro Phe Glu Phe Leu Ile Glu Asn Ile
 420 425 430

Gln His Leu Thr Pro Arg Tyr Tyr Ser Ile Ser Ser Ser Ser Leu Ser
 435 440 445

Glu Lys Gln Leu Ile Asn Val Thr Ala Val Val Glu Ala Glu Glu Glu
 450 455 460

Ala Asp Gly Arg Pro Val Thr Gly Val Val Thr Asn Leu Leu Lys Asn
 465 470 475 480

Ile Glu Ile Ala Gln Asn Lys Thr Gly Glu Lys Pro Leu Val His Tyr
 485 490 495

Asp Leu Ser Gly Pro Arg Gly Lys Phe Asn Lys Phe Lys Leu Pro Val
 500 505 510

His Val Arg Arg Ser Asn Phe Lys Leu Pro Lys Asn Ser Thr Thr Pro
515 520 525

Val Ile Leu Ile Gly Pro Gly Thr Gly Val Ala Pro Leu Arg Gly Phe
530 535 540

Val Arg Glu Arg Val Gln Gln Val Lys Asn Gly Val Asn Val Gly Lys
545 550 555 560

Thr Leu Leu Phe Tyr Gly Cys Arg Asn Ser Asn Glu Asp Phe Leu Tyr
565 570 575

Lys Gln Glu Trp Ala Glu Tyr Ala Ser Val Leu Gly Glu Asn Phe Glu
580 585 590

Met Phe Asn Ala Phe Ser Arg Gln Asp Pro Ser Lys Lys Val Tyr Val
595 600 605

Gln Asp Lys Ile Leu Glu Asn Ser Gln Leu Val His Glu Leu Leu Thr
610 615 620

Glu Gly Ala Ile Ile Tyr Val Cys Gly Asp Ala Ser Arg Met Ala Arg
625 630 635 640

Asp Val Gln Thr Thr Ile Ser Lys Ile Val Ala Lys Ser Arg Glu Ile
645 650 655

Ser Glu Asp Lys Ala Ala Glu Leu Val Lys Ser Trp Lys Val Gln Asn
660 665 670

Arg Tyr Gln Glu Asp Val Trp
675

<210> 85

<211> 4115

<212> DNA

<213> Candida tropicalis

<400> 85

catatgcgct aatcttcttt ttctttttat cacaggagaa actatccac cccacttcg 60

aaacacaatg acaactcctg cgtaacttgc aaattcttgt ctgactaatt gaaaactccg 120

gacgagtcag acctccagtc aaacggacag acagacaaac acttggtgcg atgttcatac 180

ctacagacat gtcaacgggt gttagacgac ggtttcttgc aaagacaggt gttggcatct 240

cgtacgatgg caactgcagg aggtgtcgac ttctccttta ggcaatagaa aaagactaag 300

agaacagcgt	ttttacaggt	tgcatgggt	aatgtagtat	tttttagtc	ccagcattct	360
gtgggttgct	ctgggtttct	agaataggaa	atcacaggag	aatgcaaatt	cagatggaag	420
aacaaagaga	taaaaaaca	aaaaaaactg	agttttgcac	caatagaatg	tttgatgata	480
tcatccactc	gctaaacgaa	tcatgtgggt	gatcttctct	ttagttttgg	tctatcataa	540
aacacatgaa	agtgaatcc	aaatacacta	cactccgggt	attgtccttc	gttttacaga	600
tgtctcattg	tcttactttt	gaggtcatag	gagttgcctg	tgagagatca	cagagattat	660
cacactcaca	tttatcgtag	tttctatct	catgctgtgt	gtctctgggt	ggttcatgag	720
tttgattgt	tgtacattaa	aggaatcgct	ggaaagcaaa	gctaactaaa	ttttctttgt	780
cacaggtaca	ctaacctgta	aaacttca	gccacgccag	tcttctctga	ttgggcaagt	840
gcacaaacta	caacctgcaa	aacagcactc	cgcttgtcac	aggttgtctc	ctctcaacca	900
acaaaaaat	aagattaaac	tttctttgct	catgcatcaa	tcggagttat	ctctgaaaga	960
gttgcccttg	tgtaatgtgt	gccaaactca	aactgcaaaa	ctaaccacag	aatgatttcc	1020
ctcacaatta	tataaactca	cccacatttc	cacagaccgt	aatttcatgt	ctcactttct	1080
cttttgctct	tcttttactt	agtcaggttt	gataacttcc	ttttttatta	ccctatctta	1140
tttatttatt	tattcattta	taccaaccaa	ccaaccatgg	ccacacaaga	aatcatcgat	1200
tctgtacttc	cgtacttgac	caaattgtac	actgtgatta	ctgcagcagt	attagtcttc	1260
cttatctcca	caaacatcaa	gaactacgtc	aaggcaaaga	aattgaaatg	tgtcgatcca	1320
ccatacttga	aggatgccgg	tctcactggg	attctgtctt	tgatcgccgc	catcaaggcc	1380
aagaacgacg	gtagattggc	taactttgcc	gatgaagttt	tcgacgagta	cccaaaccac	1440
accttctact	tgtctgttgc	cggtgctttg	aagattgtca	tgactgttga	cccagaaaac	1500
atcaaggctg	tcttggccac	ccaattca	gacttctcct	tgggtaccag	acacgcccac	1560
tttgctcctt	tgttggttga	cggtatcttc	accttgacg	gagaagggtg	gaagcactcc	1620
agagctatgt	tgagaccaca	gtttgctaga	gaccagattg	gacacgttaa	agccttgga	1680
ccacacatcc	aatcatggc	taagcagatc	aagttgaacc	agggaaagac	tttcgatatc	1740
caagaattgt	tcttttagatt	taccgtcgac	accgtactg	agttcttggt	tggatgaatcc	1800
gttcaactct	tgtacgatga	aaaattgggc	atcccaactc	caaacgaaat	cccaggaaga	1860
gaaaactttg	cgctgcttt	caacgtttcc	caacactact	tggccaccag	aagttactcc	1920
cagacttttt	actttttgac	caaccctaag	gaattcagag	actgtaacgc	caagggtccac	1980
cacttgcca	agtactttgt	caacaaggcc	ttgaacttta	ctcctgaaga	actcgaagag	2040
aatccaagt	ccggttacgt	tttctgttac	gaattgggtta	agcaaaccag	agatccaaag	2100

gtcttgcaag atcaattggt gaacattatg gttgccggaa gagacaccac tgccggtttg 2160
 ttgtcccttg ctttgtttga attggctaga caccagaga tgtgggtcaa gttgagagaa 2220
 gaaatogaag ttaactttgg tgttggtgaa gactcccgcg ttgaagaaat taccttcgaa 2280
 gccttgaaga gatgtgaata cttgaaggct atccttaacg aaaccttgcg tatgtacca 2340
 tctgttctcg tcaactttag aaccgccacc agagacacca ctttgccaag aggtgggtgg 2400
 gctaacggta ccgaccaat ctacattcct aaaggctcca ctgttgctta cgttgtctac 2460
 aagaccacc gtttgaaga atactacggt aaggacgcta acgacttcag accagaaaga 2520
 tggtttgaac catctactaa gaagttgggc tgggcttatg ttccattcaa cgggtgtcca 2580
 agagtctgct tgggtcaaca attgccttg actgaagctt cttatgtgat cactagattg 2640
 gccagatgt ttgaaactgt ctcatctgat ccaggctctg aataccctcc accaaagtgt 2700
 attcacttga ccatgagtca caacgatggt gtctttgtca agatgtaaag tagtcgatgc 2760
 tgggtattcg attacatgtg tataggaaga ttttggtttt ttattogttc ttttttttaa 2820
 tttttgttaa attagtttag agatttcatt aatacataga tgggtgctat ttccgaaact 2880
 ttacttctat ccctgtatc ccttattatc cctctcagtc acatgattgc tgtaattgtc 2940
 gtgcaggaca caaactccct aacggactta aaccataaac aagctcagaa ccataagccg 3000
 acatcactcc ttcttctctc ttctccaacc aatagcatgg acagaccac cctcctatcc 3060
 gaatogaaga cccttattga ctccataccc acctggaagc ccctcaagcc acacacgtca 3120
 tccagcccac ccatcaccac atccctctac tcgacaacgt ccaaagacgg cgagttctgg 3180
 tgtgcccga aatcagccat cccggccaca tacaagcagc cgttgattgc gtgcatactc 3240
 ggcgagccca caatgggagc cacgcattcg gaccatgaag caaagtacat tcacgagatc 3300
 acgggtgttt cagtgtcgca gattgagaag ttcgacgatg gatggaagta cgatctcggt 3360
 gcggattacg acttcggtgg gttgttatct aaacgaagat tctatgagac gcagcatgtg 3420
 tttcggttcg aggattgtgc gtacgtcatg agtgtgcctt ttgatggacc caaggaggaa 3480
 ggttacgtgg ttgggacgta cagatccatt gaaaggttga gctggggtaa agacggggac 3540
 gtggagtgga ccatggcgac gacgtcggat cctggtgggt ttatcccga atggataact 3600
 cgattgagca tcctgggagc aatcgcaaaa gatgtgcta gtgtattaaa ctacatacag 3660
 aaataaaaac gtgtcttgat tcattggttt ggttcttgtt gggttccgag ccaatatttc 3720
 acatcatctc cttaaattctc caagaatccc aacgtagcgt agtccagcac gccctctgag 3780
 atcttattta atatcgactt ctcaaccacc ggtggaatcc cgttcagacc attgttacct 3840
 gtagtgtgtt tgctcttgtt cttgatgaca atgatgtatt tgtcacgata cctgaaataa 3900

taaaacatcc agtcattgag cttattactc gtgaacttat gaaagaactc attcaagccg	3960
ttccccaaaa acccagaatt gaagatcttg ctcaactggt catgcaagta gtagatcgcc	4020
atgatctgat actttaccaa gctatcctct ccaagttctc ccacgtacgg caagtacggc	4080
aacgagctct ggaagctttg ttgtttgggg tcata	4115

<210> 86
 <211> 3948
 <212> DNA
 <213> *Candida tropicalis*

<400> 86	
gacctgtgac gcttcgggtg tottgccacc agtctccaag ttgaccgacg cccaagtcac	60
gtaccacttt atttccggtt acacttccaa gatggctggt actgaagaag gtgtcacgga	120
accacaagct actttctccg cttgtttcgg tcaaccattc ttggtgttgc acccaatgaa	180
gtacgctcaa caattgtctg acaagatctc gcaacacaag gctaacgcct ggttgttgaa	240
caccggttgg gttggttctt ctgctgctag aggtggttaag agatgctcat tgaagtacac	300
cagagccatt ttggacgcta tccactctgg tgaattgtcc aaggttgaat acgaaacttt	360
cccagtcttc aacttgaatg tcccaacctc ctgtccaggt gtcccaagtg aaatcttgaa	420
cccaaccaag gcctggaccg gaagggtgtg actccttcaa caaggaaatc aagtctttgg	480
ctggttaagtt tgctgaaaac ttcaagacct atgctgacca agctaccgct gaagtgagag	540
ctgcaggctc agaagcttaa agatatttat tcattattta gtttgcctat ttatttctca	600
ttaccoatca tcattcaaca ctatatataa agttacttcg gatatcattg taatcgtgcg	660
tgctgcaatt ggatgatttg gaactgcgct tgaaacggat tcatgcacga agcggagata	720
aaagattacg taatttatct cctgagacaa ttttagccgt gttcacacgc ccttctttgt	780
tctgagcgaa ggataaataa ttagacttcc acagctcatt ctaatttccg tcacgcgaat	840
attgaagggg ggtacatgtg gccgctgaat gtgggggcag taaacgcagt ctctcctctc	900
ccaggaatag tgcaacggag gaaggataac ggatagaaag cggaatgcga ggaaaatttt	960
gaacgcgcaa gaaaagcaat atccgggcta ccaggttttg agccagggaa cacactccta	1020
tttctgctca atgactgaac atagaaaaaa caccaagacg caatgaaacg cacatggaca	1080
tttagacctc ccacatgtg atagtttgtc ttaacagaaa agtataataa gaacccatgc	1140
cgctcccttt ctttcgcgcg ttcaactttt ttttttttat cttacacaca tcacgacctat	1200
gactgtacac gatattatcg ccacatactt caccaaattg tacgtgatag taccactcgc	1260
tttgattgct tatagagtcc tcgactactt ctatggcaga tacttgatgt acaagcttgg	1320

tgctaaacca	tttttccaga	aacagacaga	cggtgtttc	ggattcaaag	ctccgcttga	1380
attgttgaag	aagaagagcg	acggtaccct	catagactic	acactccagc	gtatccacga	1440
tctcgatcgt	cccgatatcc	caactttcac	attcccggtc	ttttccatca	accttgtcaa	1500
tacccttgag	ccggagaaca	tcaaggccat	cttggccact	cagttcaacg	atttctcctt	1560
gggtaccaga	cactcgcact	ttgtccttt	gttgggtgat	ggtatcttta	cgttggatgg	1620
cgccggctgg	aagcacagca	gatctatgtt	gagaccacag	tttgccagag	aacagatttc	1680
ccacgtcaag	ttgttgagc	cacacgttca	ggtgttcttc	aaacacgtca	gaaaggcaca	1740
gggcaagact	tttgacatcc	aggaattgtt	tttcagattg	accgtcgact	ccgccaccga	1800
gtttttgttt	ggtgaatccg	ttgagtcctt	gagagatgaa	tctatcggca	tgtccatcaa	1860
tgcgcttgac	tttgacggca	aggctggctt	tgctgatgct	tttaactatt	cgcagaatta	1920
tttggtctcg	agagcggtta	tgcaacaatt	gtactgggtg	ttgaacggga	aaaagttaa	1980
ggagtgaac	gctaaagtgc	acaagtttgc	tgactactac	gtcaacaagg	ctttggactt	2040
gacgcctgaa	caattggaaa	agcaggatgg	ttatgtgttt	ttgtacgaat	tggtaagca	2100
aaccagagac	aagcaagtgt	tgagagacca	attgttgaac	atcatgggtg	ctggtagaga	2160
caccaccgcc	ggtttgttgt	cgtttgtttt	ctttgaattg	gccagaaacc	cagaagttac	2220
caacaagttg	agagaagaaa	ttgaggacaa	gtttggactc	ggtgagaatg	ctagtgttga	2280
agacatttcc	tttgagtcgt	tgaagtctcg	tgaatacttg	aaggctgttc	tcaacgaaac	2340
cttgagattg	tacccatccg	tgccacagaa	tttcagagtt	gccaccaaga	acactaccct	2400
cccaagaggt	ggtggttaagg	acgggttgtc	tcctgttttg	gtgagaaagg	gtcagaccgt	2460
tatttacggt	gtctacgcag	cccacagaaa	cccagctgtt	tacggtaagg	acgctcttga	2520
gttttagacca	gagagatggt	ttgagccaga	gacaaagaag	cttggctggg	ccttcctccc	2580
attcaacggt	ggtccaagaa	tctgtttggg	acagcagttt	gccttgacag	aagcttcgta	2640
tgtcactgtc	aggttgctcc	aggagtttgc	acacttgtct	atggaccag	acaccgaata	2700
tccacctaag	aaaatgtcgc	atttgacca	gtcgttttc	gacggtgcc	atattgagat	2760
gtattagagg	gtcatgtgtt	attttgattg	tttagtttgt	aattactgat	taggttaatt	2820
catggattgt	tatttattga	taggggtttg	cgcgtgttgc	attcacttgg	gatcgttcca	2880
ggttgatgtt	tccttccatc	ctgtcgagtc	aaaaggagtt	ttgttttgta	actccggacg	2940
atgttttaaa	tagaaggctg	atctccatgt	gattgttttg	actgttactg	tgattatgta	3000
atctgcggac	gttatacaag	catgtgattg	tggttttgca	gccttttgca	cgacaaatga	3060
tcgtcagacg	attacgtaat	ctttgttaga	ggggtaaaaa	aaaacaaaat	ggcagccaga	3120

atttcaaaca ttctgcaaac aatgcaaaaa atgggaaact ccaacagaca aaaaaaaaaa 3180
 ctccgcagca ctccgaaccc acagaacaat ggggcgccag aattattgac tattgtgact 3240
 tttttacgct aacgctcatt gcagtgtagt gcgtcttaca cgggggtattg ctttctacaa 3300
 tgcaagggca cagttgaagg tttgcaccta acgttgcccc gtgtcaactc aatttgacga 3360
 gtaacttcct aagctcgaat tatgcagctc gtgcgtcaac ctatgtgcag gaaagaaaaa 3420
 atccaaaaaa atcgaaaatg cgactttcga ttttgaataa accaaaaaga aaaatgtcgc 3480
 acttttttct cgctctcgct ctctcgaccc aaatcacaac aaatcctcgc gcgcagtatt 3540
 tcgacgaaac cacaacaaat aaaaaaaca aattctacac cacttctttt tcttcaccag 3600
 tcaacaaaaa acaacaaatt atacaccatt tcaacgattt ttgctcttat aaatgctata 3660
 taatggttta attcaactca ggtatgttta ttttactgtt ttcagctcaa gtatgttcaa 3720
 atactaacta cttttgatgt ttgtcgcttt tctagaatca aaacaacgcc cacaacacgc 3780
 cgagcttgct gaatagacgg tttgtttact cattagatgg tccagatta cttttcaagc 3840
 caaagtctct cgagttttgt ttgctgtttc cccaattcct aactatgaag gggtttttata 3900
 aggtccaaag accccaaggc atagtttttt tggttccttc ttgtcgtg 3948

<210> 87

<211> 3755

<212> DNA

<213> *Candida tropicalis*

<400> 87

gctcaacaat tgtctgacaa gatctcgcaa cacaaggcta acgcctgggt gttgaacact 60
 ggttgggttg gttcttctgc tgctagaggt ggtaagagat gttcattgaa gtacaccaga 120
 gccatthttg acgctatcca ctctggtgaa ttgtccaagg ttgaatacga gactttccca 180
 gtcttcaact tgaatgtccc aacotcctgc ccaggtgtcc caagtgaat cttgaaccca 240
 accaaggcct ggaccgaagg tgttgactcc ttcaacaagg aaatcaagtc tttggctggt 300
 aagtttgctg aaaacttcaa gacctatgct gaccaagcta ccgctgaagt tagagctgca 360
 ggtccagaag cttaaagata tttattcact atttagtttg cctattttatt tctcatcacc 420
 catcatcatt caacaatata tataaagtta tttcggaact catatatcat tgtaatcgtg 480
 cgtgttgcaa ttgggtaatt tgaaactgta gttggaacgg attcatgcac gatgcggaga 540
 taacacgaga ttatctccta agacaatttt ggctcattc acacgccctt cttctgagct 600
 aaggataaat aattagactt cacaagttca ttaaaatata cgtcacgcga aaactgcaac 660
 aataaggaag gggggggtag acgtagccga tgaatgtggg gtgccagtaa acgcagtctc 720
 tctctcccc ccccccccc cccctcagg aatagtacaa cgggggaagg ataacggata 780

gcaagtggaa tgcgaggaaa attttgaatg cgcaaggaaa gcaatatccg ggctatcagg 840
ttttgagcca ggggacacac tcctcttctg cacaaaaact taacgtagac aaaaaaaaaa 900
aactccacca agacacaatg aatcgcacat ggacatttag acctccccac atgtgaaagc 960
ttctctggcg aaagcaaaaa aagtataata aggacccatg ccttccctct tcctgggccg 1020
tttcaacttt ttctttttct ttgtctatca acacacacac acctcacgac catgactgca 1080
caggatatta tcgccacata catcaccaaa tggtagctga tagtaccact cgctttgatt 1140
gcttataggg tcctcgacta cttttacggc agatacttga tgtacaagct tgggtgctaaa 1200
ccgtttttcc agaaacaaac agacggttat ttcggtattca aagctccact tgaattgtta 1260
aaaaagaaga gtgacggtag cctcatagac ttactctcg agcgtatcca agcgtcaat 1320
cgtccagata tcccaacttt tacattccca atcttttcca tcaaccttat cagcaccctt 1380
gagccggaga acatcaaggc tatcttggcc acccagttca acgatttctc cttgggcacc 1440
agacactcgc actttgctcc tttgttgggc gatggtatct ttaccttga cgggtgccggc 1500
tggaagcaca gcagatctat gttgagacca cagtttgcca gagaacagat ttcccacgtc 1560
aagttgttgg agccacacat gcaggtgttc ttcaagcacg tcagaaaggc acagggcaag 1620
acttttgaca tccaagaatt gtttttcaga ttgaccgtcg actccgccac tgagtttttg 1680
tttggatgaat ccgttgagtc cttgagagat gaatctattg ggatgtccat caatgcactt 1740
gactttgacg gcaaggctgg ctttctgat gcttttaact actcgcagaa ctatttggct 1800
tcgagagcgg ttatgcaaca attgtactgg gtgttgaacg ggaaaaagtt taaggagtgc 1860
aacgctaaag tgcacaagtt tgctgactat tacgtcagca aggctttgga cttgacacct 1920
gaacaattgg aaaagcagga tggttatgtg ttcttgtacg agttgggtcaa gcaaaccaga 1980
gacaggcaag tggtgagaga ccagttgttg aacatcatgg ttgccggtag agacaccacc 2040
gccggtttgt tgcgtttgt tttctttgaa ttggccagaa acccagaggt gaccaacaag 2100
ttgagagaag aaatcgagga caagtttggc cttgggtgaga atgctcgtgt tgaagacatt 2160
tcctttgagt cgttgaagtc atgtgaatac ttgaaggctg ttctcaacga aactttgaga 2220
ttgtacccat ccgtgccaca gaatttcaga gttgccacca aaaacactac ccttccaagg 2280
ggaggtggta aggacgggtt atctcctgtt ttggtcagaa aggtcacaac cgttatgtac 2340
ggtgtctacg ctgccacag aaaccagct gtctacggta aggacgccct tgagttttaga 2400
ccagagaggt ggtttgagcc agagacaaag aagcttggct gggccttcct tcattcaac 2460
ggtggtccaa gaatttgctt gggacagcag ttgctcttga cagaagcttc gtatgtcact 2520
gtcagattgc tccaagagtt tggacacttg tctatggacc ccaacaccga atatccacct 2580

aggaaaatgt cgcatttgac catgtccctt ttcgacggtg ccaacattga gatgtattag 2640
 aggatcatgt gttatTTTTg attggttttag tctgtttgta gctattgatt aggttaattc 2700
 acggattggt atttattgat aggggggtgcg tgtgtgtgtg tgtgttgcat tcacatggga 2760
 tcgttccagg ttgttgtttc cttccatcct gttgagtcaa aaggagtttt gttttgtaac 2820
 tccggacgat gtcttagata gaaggtcgat ctccatgtga ttgtttgact gctactctga 2880
 ttatgtaatc tgtaaagcct agacgttatg caagcatgtg attgtgggtt ttgcaacctg 2940
 ttgacacgac aaatgatcga cagtcgatta cgtaatccat attattttaga ggggtaataa 3000
 aaaataaatg gcagccagaa tttcaaacat ttgcaaaca atgcaaaaaga tgagaaactc 3060
 caacagaaaa aataaaaaaa ctccgcagca ctccgaacca acaaaacaat ggggggcgcc 3120
 agaattattg actattgtga ctttttttta ttttttccgt taactttcat tgcagtgaag 3180
 tgtgttacac ggggtggtga tgggtgttgg ttctacaatg caagggcaca gttgaagggt 3240
 tccacataac gttgcaccat atcaactcaa tttatcctca ttcattgtgat aaaagaagag 3300
 ccaaaaggta attggcagac cccccaaggg gaacacggag tagaaagcaa tggaaacacg 3360
 cccatgacag tgccatttag cccacaacac atctagtatt cttttttttt tttgtgcgca 3420
 ggtgcacacc tggacttttag ttattgcccc ataaagttaa caatctcacc tttggctctc 3480
 ccagtgtctc cgctccaga tgctcgtttt acacctcoga gctaacgaca acacaacacc 3540
 catgagggga atgggcaaag ttaaacactt ttggtttcaa tgattcctat ttgctactct 3600
 cttgttttgt gttttgattt gcaccatgtg aaataaacga caattatata taccttttgc 3660
 tctgtcctcc aatgtctctt tttgctgcca ttttgccttt tgctttttgc ttttgcactc 3720
 tctccactc ccacaatcag tgcagcaaca cacaa 3755

<210> 88

<211> 3900

<212> DNA

<213> *Candida tropicalis*

<400> 88

gacatcataa tgaccgggtt atttcgcctt caggttgctt atttgagccg taaagtgcag 60
 tagaaacttt gccttggtt caaactctag tataatggtg ataactggtt gcaactctgc 120
 cataggcattg aaaataggcc gttatagtac tatatttaat aagcgttaga gtataggatg 180
 catatgaccg gtttttctat atttttaaga taatctctag taaattttgt attctcagta 240
 ggatttcac ccaatttcgca accaattctg gcgaaaaaat gattctttta cgtcaaaagc 300
 tgaatagtgc agtttaaagc acctaaaatc acatatacag cctctagata cgacagagaa 360

gctctttatg atctgaagaa gcattagaat agctactatg agccactatt ggtgtatata 420
ttagggattg gtgcaattaa gtacgtacta ataaacagaa gaaaatactt aaccaatttc 480
tggtgtatac ttagtggtga gggacctttt ctgaacattc ggggtcaaact tttttttgga 540
gtgcgacatc gattttttcgt ttgtgtaata atagtgaacc tttgtgtaat aaatcttcat 600
gcaagacttg cataattcga gcttgggagt tcacgccaat ttgacctcgt tcatgtgata 660
aaagaaaagc caaaaggtaa ttagcagacg caatgggaac atggagtgga aagcaatgga 720
agcacgcca ggacggagta atttagtcca cactacatct gggggttttt tttttgtgcg 780
caagtacaca cctggacttt agtttttgcc ccataaagtt aacaatctaa cctttggctc 840
tccaactctc tcgccccca aatattcgtt tttacacct caagctagcg acagcacaac 900
accattaga ggaatggggc aaagttaaac acttttggt tcaatgattc ctattcgcta 960
ctacattctt ctcttgtttt gtgctttgaa ttgcaccatg tgaaataaac gacaattata 1020
tatacctttt catccctcct cctatatctc tttttgctac attttgtttt ttacgtttct 1080
tgcttttgca ctctccact cccacaaaga aaaaaaact acactatgtc gtcttctcca 1140
tcgtttgcc aagaggttct cgctaccact agtccttaca tcgagtactt tcttgacaac 1200
tacaccagat ggtactactt catacctttg gtgcttcttt cgttgaactt tataagtttg 1260
ctccacacaa ggtacttgga acgcagggtc cagccaagc cactcggtaa ctttgtcagg 1320
gaccctacgt ttggtatcgc tactccgttg cttttgatct acttgaagtc gaaaggtacg 1380
gtcatgaagt ttgcttgggg cctctggaac aacaagtaca tcgtcagaga cccaaagtac 1440
aagacaactg ggctcaggat tgttggcctc ccattgattg aaaccatgga ccagagaaac 1500
atcaaggctg ttttggtctac tcagttcaat gatttctctt tgggaaccag acacgatttc 1560
ttgtactcct tgttgggtga cggtattttc accttgagc gtgctggctg gaaacatagt 1620
agaactatgt tgagaccaca gtttgctaga gaacaggttt ctcacgtcaa gttgttgag 1680
ccacacgttc aggtgttctt caagcacgtt agaaagcacc gcggtcaaac gttcgacatc 1740
caagaattgt tcttcagggt gaccgtcgac tcgcccaccg agttcttggt tggtagtct 1800
gctgaatcct tgagggacga atctattgga ttgacccaa ccaccaagga tttcgatggc 1860
agaagagatt tcgctgacgc tttcaactat tcgcagactt accaggccta cagatttttg 1920
ttgcaacaaa tgtactggat cttgaatggc tcggaattca gaaagtcgat tgctgtcgtg 1980
cacaagtttg ctgaccacta tgtgcaaaag gctttggagt tgaccgacga tgacttgacg 2040
aaacaagacg gctatgtgtt cttgtacgag ttggctaagc aaaccagaga cccaaaggtc 2100
ttgagagacc agttattgaa cattttggtt gccggtagag acacgaccgc cggtttgttg 2160

tcatttggtt tctacgagtt gtcaagaaac cctgaggtgt ttgctaagtt gagagaggag 2220
 gtggaaaaca gatttggact cggatgaaga gctcgtgttg aagagatctc gtttgagtcc 2280
 ttgaagtctt gtgagtactt gaaggctgtc atcaatgaaa ccttgagatt gtacccatcg 2340
 gttccacaca acttttagagt tgctaccaga aacactaccc tccaagagg tgggtggtgaa 2400
 gatggatact cgccaattgt cgtcaagaag ggtcaagttg tcatgtacac tgttattgct 2460
 acccacagag acccaagtat ctacggtgcc gacgctgacg tcttcagacc agaaagatgg 2520
 tttgaaccag aaactagaaa gttgggctgg gcatacgttc cattcaatgg tggccaaga 2580
 atctgtttgg gtcaacagtt tgccttgacc gaagcttcat acgtcactgt cagattgctc 2640
 caggagtttg cacacttgtc tatggacca gacaccgaat atccacaaa attgcagaac 2700
 accttgacct tgctgctctt tgatggtgct gatgtagaa tgtactaagg ttgcttttcc 2760
 ttgctaattt tcttctgtat agcttggtga tttaaattga atcggaatt gatttttctg 2820
 ataccaataa ccgtagtgcg atttgaccaa aaccgttcaa actttttgtt ctctcgttga 2880
 cgtgctcgct catcagcact gtttgaagac gaaagagaaa attttttgta aacaacactg 2940
 tccaaattta cccaacgtga accattatgc aaatgagcgg ccctttcaac tggctgctgg 3000
 aagcattcgg ggatatctac aacgccotta agtttgaaac agacattgat ttagacacca 3060
 tagatttcag cggcatcaag aatgaccttg cccacatttt gacgaccca acaccactgg 3120
 aagaatcacg ccagaaacta ggcgatggat ccaagcctgt gaccttgccc aatggagacg 3180
 aagtggagtt gaaccaagcg ttctagaag ttaccacatt attgtcgaat gagtttgact 3240
 tggaccaatt gaacgaggca gagttgttat actacgtgg cgacatatcc tacaagaagg 3300
 gcacatcaat cgcagacagt gccagattgt cttattattt gagagcaaac tacatcttga 3360
 acatacttgg gtatttgatt tcgaagcagc gattggattt gatagtcacg gacaacgacg 3420
 cgttgtttga tagtattttg aaaagttttg aaaagatcta caagttgata agcgtgttga 3480
 acgatatgat tgacaagcaa aaggtgacaa gcgacatcaa cagtctagca ttcattcaatt 3540
 gcatcaacta ctcgagaggt caactattct ccgcacacga acttttggga ctggttttgt 3600
 ttggatttgt cgacatctat ttcaaccagt ttggcacatt agacaactac aagaaggat 3660
 tggcattgat actgaagaac atcagcgatg aagacatctt gatcatacac ttctcccat 3720
 cgacactaca attgtttaag ctggtgttgg acaagaaaaga cgacgctgca gttgaacagt 3780
 tctacaagta catcacttca acagtgtcac gagactacaa ctccaacatc ggctccacag 3840
 ccaaagatga tatcgatttg tccaaaacca aactcagtgg ctttgaggtg ttgacgagtt 3900

<210> 89

<211> 3668
 <212> DNA
 <213> *Candida tropicalis*

<400> 89
 cctgcagaat tcgcggccgc gtcgacagag tagcagttat gcaagcatgt gattgtgggt 60
 tttgcaacct gtttgacga caaatgatcg acagtcgatt acgtaatcca tattatttag 120
 aggggtaata aaaaataaat ggcagccaga atttcaaaca ttttgaaac aatgcaaaag 180
 atgagaaact ccaacagaaa aaataaaaaa actccgcagc actccgaacc aacaaaacaa 240
 tggggggcgc cagaattatt gactattgtg actttttttt attttttcg ttaactttca 300
 ttgcagtga gtgtgttaca cggggtggtg atggtgttgg tttctacaat gcaagggcac 360
 agttgaaggt ttccacataa cgttgacca tatcaactca atttatctc attcatgtga 420
 taaaagaaga gccaaaaggt aattggcaga cccccaagg ggaacacgga gtagaaagca 480
 atggaaacac gcccatgaca gtgccattta gccacaaca catctagtat tctttttttt 540
 ttttgtgcgc aggtgcacac ctggacttta gttattgcc cataaagtta acaatctcac 600
 ctttggtct cccagtgtct ccgcctccag atgctcgttt tacaccctcg agctaacgac 660
 aacacaacac ccatgagggg aatgggcaaa gttaaacact tttggtttca atgattccta 720
 tttgtactc tcttgttttg tgttttgatt tgcacatgt gaaataaacg acaattatat 780
 ataccttttc gtctgtcctc caatgtctct ttttctgcc attttgcctt ttgctttttg 840
 cttttgcact ctctccact cccacaatca gtgcagcaac acacaaagaa gaaaaataaa 900
 aaaacctaca ctatgtcgtc ttctccatcg tttgtcagg aggttctcgc taccactagt 960
 ccttacatcg agtactttct tgacaactac accagatggg actacttcat ccctttgggtg 1020
 cttctttcgt tgaacttcat cagcttgctc cacacaaagt acttggaacg caggttccac 1080
 gccagccgc tcggtaacgt cgtgttggt cctacgtttg gtatcgctac tccgttgatc 1140
 ttgatctact taaagtcgaa aggtacagtc atgaagtttg cctggagctt ctggaacaac 1200
 aagtacattg tcaaagaccc aaagtacaag accactggcc ttagaattgt cggcctccca 1260
 ttgattgaaa ccatagaccc agagaacatc aaagctgtgt tggctactca gttcaacgat 1320
 ttctccttgg gaactagaca cgatttcttg tactccttgt tgggcgatgg tatttttacc 1380
 ttggacggtg ctggctggaa acacagtaga actatgttga gaccacagtt tgctagagaa 1440
 caggtttccc acgtcaagtt gttggaacca cagttcagg tgttcttcaa gcacgttaga 1500
 aaacaccgcg gtcagacttt tgacatccaa gaattgttct tcagattgac cgtcgactcc 1560
 gccaccgagt tcttgttttg tgagtctgct gaatccttga gagacgactc tgttggtttg 1620
 accccaacca ccaaggattt cgaaggcaga ggagatttcg ctgacgcttt caactactcg 1680

cagacttacc aggcctacag atttttgttg caacaaatgt actggatttt gaatggcgcg 1740
gaattcagaa agtcgattgc catcgtgcac aagtttgctg accactatgt gcaaaaggct 1800
ttggagttga cgcacgatga cttgcagaaa caagacggct atgtgttctt gtacgagttg 1860
gctaagcaaa ctagagaccc aaaggtcttg agagaccagt tgttgaacat tttggttgcc 1920
ggtagagaca cgaccgccgg tttgttgctg tttgtgttct acgagttgtc gagaaaccct 1980
gaagtgtttg ccaagttgag agaggagggtg gaaaacagat ttggactcgg cgaagaggct 2040
cgtgttgaag agatctcttt tgagtccttg aagtcctgtg agtacttgaa ggctgtcatc 2100
aatgaagcct tgagattgta cccatctgtt ccacacaact tcagagttgc caccagaaac 2160
actacccttc caagaggcgg tggtaaagac ggatgctcgc caattgttgt caagaagggt 2220
caagttgtca tgtacactgt cattgggtacc cacagagacc caagtatcta cggtgccgac 2280
gccgacgtct tcagaccaga aagatgggtc gagccagaaa ctagaaagtt gggctgggca 2340
tatgttccat tcaatgggtg tccaagaatc tgtttgggtc agcagtttgc cttgactgaa 2400
gcttcatacg tcaactgtcag attgctccaa gagtttgaa acttgctcct ggatccaaac 2460
gctgagtacc caccaaaatt gcagaacacc ttgacctgt cactctttga tgggtctgac 2520
gttagaatgt tctaagggtg cttatccttg ctagtgttat ttatagtttg tgtattttaa 2580
ttgaatcggc gattgatttt totggtacta ataactgtag tgggttttga ccaaaaccgt 2640
tcaaactttt tttttttttt tottccccct acctcgttg ctgctcatc agcactgttt 2700
gaaaacgaaa aaagaaaatt ttttgtaaac aacattgcc aaacttacc aacgtgaacc 2760
attataacca aatgagcggc gctttcaact ggtcactgga ggcattcggg gatattctaca 2820
acacccttaa gtttgaggaa gacattgatt tagacaccat agatttcagc ggcattcaaga 2880
atgacctgtt ccacattttg acaaccccaa caccactgga agaatcgcg cagaaaactag 2940
gcgatggatc caagcctgtg gccttgccca atggagacga agtggagttg aaccaagcgt 3000
tcctagaagt taccacatta ttgtcgaacg agtttgactt ggaccaattg aacgcggccg 3060
agttgttata ctacgcggc gacatatcct acaagaaggg cacatcaatt gccgacagtg 3120
ccagattgtc ttactatttg agagcaaact acatcttgaa catacttggg tactttattt 3180
cgaagcagcg attggatgtg atagtcaccg acaacaacgc gttgtttgat aatattttga 3240
aaagttttga aaagatctac aagttgataa gcgcgttgaa cgatatgatt gacaagcaaa 3300
aggtgacaag cgacatcaac agtctagcat ttatcaactg catcaactac tcgaggggtc 3360
aactattctc cgcacacgaa cttttgggac tggttttgtt tggattgggt gacaactatt 3420
tcaaccagtt tggctcatta gacaactaca agaaagtatt ggcattgata ctgaagaaca 3480

tcagtgatga agatatcttg atcgtacgct tcctcccatc gacactacaa ttgtttaagc	3540
tggtgttgga taagaaagac gacgccactg ttgaccagtt ctacaagtac atcacctcaa	3600
cagtgtcgca agactacaac tccaacatcg gagccacagc caaagatgat atcgatttgt	3660
ccaaagcc	3668

<210> 90
 <211> 3826
 <212> DNA
 <213> *Candida tropicalis*

<400> 90	
tggagtcgcc agacttgctc acttttgact cccttcgaaa ctcaaagtac gttcaggcgg	60
tgctcaacga aacgctccgt atctacccgg gggtaccacg aaacatgaag acagctacgt	120
gcaacacgac gttgccacgc ggaggaggca aagacggcaa ggaacctatc ttggtgcaga	180
agggacagtc cgttgggttg attactattg ccacgcagac ggaccagag tattttgggg	240
ccgacgctgg tgagttaag ccggagagat ggtttgattc aagcatgaag aacttggggg	300
gtaaatactt gccgttcaat gctgggccac ggacttgctt ggggcagcag tacactttga	360
ttgaagcgag ctacttgcta gtccggttg cccagacctt ccgggcaata gatttgcagc	420
caggatcggc gtacccacca agaaagaagt cgttgatcaa catgagtgtc gccgacgggg	480
tgtttgtaaa gctttataag gatgtaacgg tagatggata gttgtgtagg aggagcggag	540
ataaattaga tttgattttg tgtaagggtt tggatgtcaa cctactccgc acttcatgca	600
gtgtgtgtga cacaagggtg tactacgtgt gcgtgtgcgc caagagacag cccaaggggg	660
tggtagtgtg tgttgcgga agtgcagtgt acacaacgcg tgggttcttg ccaatggtgg	720
actaagtgca ggtaagcagc gacctgaaac attcctcaac gcttaagaca ctggtggtag	780
agatgcggac caggctattc ttgtcgtgct acccggcgca tggaaaatca actgcgggaa	840
gaataaattt atccgtagaa tccacagagc ggataaattt gccacactcc atcatcaacc	900
acgccgccac taactacatc actcccctat tttctctctc tctctttgtc ttactcgcgt	960
ccggtttcct tagccacaga tacacacca ctgcaaacag cagcaacaat tataaagata	1020
cgccaggccc accttctttc tttttcttca cttttttgac tgcaactttc tacaatccac	1080
cacagccacc accacagccg ctatgattga acaactccta gaatattggt atgtcgttgt	1140
gccagtgttg tacatcatca aacaactcct tgcatacaca aagactcgcg tcttgatgaa	1200
aaagtgggt gctgctccag tcacaaacaa gttgtacgac aacgctttcg gtatcgtcaa	1260
tggatggaag gctctccagt tcaagaaaga gggcagggct caagagtaca acgattacaa	1320

gtttgaccac tccaagaacc caagcgtggg cacctacgtc agtattcttt tcggcaccag 1380
gatcgtcgtg accaaagatc cagagaatat caaagctatt ttggcaaccc agtttggtga 1440
tttttctttg ggcaagaggc acactctttt taagcctttg ttaggtgatg ggatcttcac 1500
attggacggc gaaggctgga agcacagcag agccatgttg agaccacagt ttgccagaga 1560
acaagttgct catgtgacgt cgttggaaac acacttccag ttgttgaaga agcatattct 1620
taagcacaag ggtgaatact ttgatatcca ggaattgttc tttagattta ccgttgattc 1680
ggccacggag ttcttatttg gtgagtcctg gcactcctta aaggacgaat ctattggtat 1740
caaccaagac gatatagatt ttgctggtag aaaggacttt gctgagtcgt tcaacaaagc 1800
ccaggaatac ttggctatta gaaccttggg gcagacgttc tactggtttg tcaacaacaa 1860
ggagttttaga gactgtacca agctgggtgca caagttcacc aactactatg ttcagaaagc 1920
tttggtatgct agcccagaag agcttgaaaa gcaaagtggg tatgtgttct tgtacgagct 1980
tgtcaagcag acaagagacc ccaatgtgtt gcgtgaccag tctttgaaca tcttgttggc 2040
cggaagagac accactgctg ggttggtgtc gtttgctgtc tttgagttgg ccagacaccc 2100
agagatctgg gccaaagtga gagaggaaat tgaacaacag tttggtcttg gagaagactc 2160
tcgtgttgaa gagattacct ttgagagctt gaagagatgt gagtacttga aagcgttcct 2220
taatgaaacc ttgcgtattt acccaagtgt cccaagaaac ttcagaatcg ccaccaagaa 2280
cacgacattg ccaaggggcg gtggttcaga cgttacctcg ccaatcttga tccaaaaggg 2340
agaagctgtg tcgtatggtg tcaactctac tcatttggaac cctgtctatt acggccctga 2400
tgctgctgag ttcagaccag agagatggtt tgagccatca accaaaaagc tcggctgggc 2460
ttaacttgcca ttcaacggtg gtccaagaat ctgtttgggt cagcagtttg ccttgacgga 2520
agctggctat gtgttggtta gattggtgca agagttctcc cacgttaggc tggaccaga 2580
cgaggtgtac ccgccaaga gggtgaccaa cttgaccatg tgtttgagcagg atggtgctat 2640
tgtcaagttt gactagcggc gtggtgaatg cgtttgattt tgtagtttct gtttgagta 2700
atgagataac tattcagata aggcgagtggt atgtacgttt tgtaagagtt tccttacaac 2760
cttggtgggg tggtgaggt tgaggttgca tcttggggag attacacctt ttgcagctct 2820
ccgtatacac ttgtactctt tgtaacctct atcaatcatg tggggggggg ggttcattgt 2880
ttggccatgg tgggtgcatgt taaatcogcc aactaccaa tctcacatga aactcaagca 2940
cactaaaaaa aaaaaagatg ttgggggaaa actttggttt ccttcttag taattaaaca 3000
ctctcactct cactctcact ctctccactc agacaaacca accacctggg ctgcagacaa 3060
ccagaaaaaa aaagaacaaa atccagatag aaaaacaaag ggctggacaa ccataaataa 3120

acaatctagg gtctactcca ttttccaactg tttcttcttc ttcagactta gctaacaaac	3180
aactcacttc accatggatt acgcaggcat cacgcgtggc tccatcagag gcgaggcctt	3240
gaagaaactc gcagaattga ccatccagaa ccagccatcc agcttgaaag aaatcaacac	3300
cggcatccag aaggacgact ttgccaagtt gttgtctgcc accccgaaaa tccccaccaa	3360
gcacaagttg aacggcaacc acgaattgtc tgaggtcgcc attgccaaaa aggagtacga	3420
ggtgttgatt gcottgagcg acgccacaaa agaccaatc aaagtgacct cccagatcaa	3480
gatcttgatt gacaagttca aggtgtactt gtttgagttg cctgaccaga agttctccta	3540
ctccatcgtg tccaactccg tcaacatcgc cccctggacc ttgctcgggg agaagttgac	3600
cacgggcttg atcaacttgg ccttcagaa caacaagcag cacttggaag aggtcattga	3660
catcttcaac gagttcatcg acaagttctt tggcaacacg gagccgcaat tgaccaactt	3720
cttgacctg tgcggtgtgt tggacgggtt gattgacct gccaaacttct tgagcgtgtc	3780
ctcgcggaac ttcaagatct tcttgaactt ggactcgtat gtggac	3826

<210> 91
 <211> 3910
 <212> DNA
 <213> Candida tropicalis

<400> 91	
ttacaatcat ggagctcgct aggaaccag atgtctggga gaagctccgc gaagaggtca	60
acacgaactt tggcatggag tcgccagact tgctcacttt tgactctctt agaagctcaa	120
agtacgttca ggcgggtgtc aacgaaacgc ttcgtatcta cccgggggtg ccacgaaaca	180
tgaagacagc tacgtgcaac acgacgttgc cgcgtggagg aggcaaagac ggtaaggaac	240
ctattttggg gcagaagggc cagtcogttg ggttgattac tattgccacg cagacggacc	300
cagagtattt tggggcagat gctggtgagt tcaaaccgga gagatggttt gattcaagca	360
tgaagaactt ggggtgtaag tacttgccgt tcaatgctgg gcccggact tgtttggggc	420
agcagtacac tttgattgaa gcgagctatt tgctagtcag gttggcgag acctaccggg	480
taatcgattt gctgccaggg tcggcgtaac caccaagaaa gaagtcggtt atcaatatga	540
gtgctgccga tgggggtggt gtaaagtttc acaaggatct agatggatat gtaaggtgtg	600
taggaggagc ggagataaat tagatttgat tttgtgtaag gtttagcacg tcaagctact	660
ccgcactttg tgtgtaggga gcacatactc cgtctgcgcc tgtgccaaga gacggcccag	720
gggtagtgtg tgggtgtgga agtgcattgt acacaatacc ctggttctgg ccaattgggg	780
atttagtgta ggtaagctgc gacctgaaac actcctcaac gcttgagaca ctggtgggta	840
gagatgcggg ccaggaggct attcttgctg tgctaccggt gcacggaaaa tcgattgagg	900

atgagtgcgt	ttgattttgt	agtttctgtt	tgcagtaatg	agataactat	tcagataagg	2760
cgggtggatg	tacgttttgt	aagagtttcc	ttacaaccct	ggtgggtgtg	tgaggttgca	2820
tcttagggag	agatagcacc	ttttgcagct	ctccgtatac	agttttactc	tttgtaacct	2880
atgccaatca	tgtggggatt	cattgtttgc	ccatgggtgt	gcatgcaaaa	tcccccaac	2940
taccaatct	cacatgaaac	tcaagcacac	tagaaaaaaa	agatgttgcg	tgggttcttt	3000
tgatgttggg	gaaaactttc	gtttcctttc	tcagtaatta	aacgttctca	ctcagacaaa	3060
ccacctgggc	tgcagacaac	cagaaaaaac	aaaatccaga	tagaagaaga	aagggctgga	3120
caaccataaa	taaacaacct	agggccact	ccatctttca	cttcttcttc	ttcagactta	3180
tctaacaaac	gactcacttc	accatggatt	acgcaggtat	cacgcgtggg	tccatcagag	3240
gcgaagcctt	gaagaaactc	gccgagttga	ccatccagaa	ccagccatcc	agcttgaaag	3300
aatcaacac	cggcatccag	aaggacgact	ttgccaaagt	gttgtcttcc	accccgaaaa	3360
tccacaccaa	gcacaagttg	aatggcaacc	acgaattgtc	cgaagtcgcc	attgccaaaa	3420
aggagtacga	ggtgttgatt	gccttgagcg	acgccacgaa	agaaccaatc	aaagtcacct	3480
cccagatcaa	gatcttgatt	gacaagttca	aggtgtactt	gtttgagttg	cccgaccaga	3540
agttctccta	ctccatcggt	tccaactccg	ttaacattgc	cccctggacc	ttgctcggtg	3600
agaagttgac	cacgggcttg	atcaacttgg	cgttccagaa	caacaagcag	cacttgagcg	3660
aagtcatcga	catcttcaac	gagttcatcg	acaagttctt	tggcaacaca	gagccgcaat	3720
tgaccaactt	cttgaccttg	tccggtgtgt	tggacggggt	gattgaccat	gccaaacttct	3780
tgagcgtgtc	ctccaggacc	ttcaagatct	tottgaactt	ggactcgttt	gtggacaact	3840
cggacttctt	gaacgacgtg	gagaactact	ccgacttttt	gtacgacgag	ccgaacgagt	3900
accagaactt						3910

<210> 92
 <211> 3150
 <212> DNA
 <213> *Candida tropicalis*

<400>	92	
gaattctttg	gatctaattc	cagctgatct
tgctaatect	tatcaacgta	gttgtgatca
		60
ttgtttgtct	gaattatata	caccagtggg
agaatatggt	ctaatttgca	cgtccactg
		120
gcattgtgtg	tttgtggggg	ggggggggtg
cacacatttt	tagtgccatt	ctttgttgat
		180
taccctccc	ccctatcatt	cattcccaca
ggattagttt	tttcctcact	ggaattcgct
		240
gtccacctgt	caaccccccc	cccccccccc
ccactgccc	taccctgccc	tgcctgcac
		300

gtcctgtggtt ttgtgctgtg tctttccac gctataaaag ccotggcgtc cggccaaggt	360
ttttccaccc agccaaaaaa acagtctaaa aaatttggtt gatccttttt ggttgcaagg	420
ttttccacca ccacttccac cacctcaact attcgaacaa aagatgctcg atcagatctt	480
acattactgg tacattgtct tgccattgtt ggccattatc aaccagatcg tggctcatgt	540
caggaccaat tatttgatga agaaattggg tgctaagcca ttcacacacg tccaacgtga	600
cgggtggttg ggcttcaaat tcggccgtga attcctcaaa gcaaaaagtg ctgggagact	660
ggttgattta atcatctccc gtttccacga taatgaggac actttctcca gctatgcttt	720
tggcaaccat gtggtgttca ccagggaccc cgagaatatc aaggcgcttt tggcaacca	780
gtttggtgat ttttcattgg gcagcagggg caagttcttc aaaccattat tggggtacgg	840
tatcttcaca ttggacgccg aaggctggaa gcacagcaga gccatgttga gaccacagtt	900
tgccagagaa caagttgctc atgtgacgtc gttggaacca cacttccagt tgttgaagaa	960
gcatactctt aaacacaagg gtgagtactt tgatatccag gaattgttct ttagatttac	1020
tgctgactcg gccacggagt tcttatttgg tgagtcctgt cactccttaa aggacgagg	1080
aattggctac gacacgaaag acatgtctga agaaagacgc agatttgccg acgcgttcaa	1140
caagtcgcaa gtctacgtgg ccaccagagt tgctttacag aacttgtact ggttggtcaa	1200
caacaaagag ttcaaggagt gcaatgacat tgtccacaag tttaccaact actatgttca	1260
gaaagccttg gatgctaccc cagaggaact tgaaaagcaa ggccgggtatg tgttcttgta	1320
tgagcttgct aagcagacga gagaccccaa ggtgttgctg gaccagtctt tgaacatctt	1380
gttggcagga agagacacca ctgctgggtt gttgtccttt gctgtgtttg agttggccag	1440
aaaccacac atctgggcca agttgagaga ggaaattgaa cagcagtttg gtcttgga	1500
agactctcgt gttgaagaga ttaccttga gagcttgaag agatgtgagt acttgaaggc	1560
cgtgttgaac gaaacttga gattacacc aagtgtccca agaaacgcaa gatttgcat	1620
taaagacacg actttacaa gaggcgggtg cccaacgac aaggatccta tcttgatcag	1680
gaaggatgag gtggtgcagt actccatctc ggcaactcag acaaatcctg cttattatgg	1740
cgccgatgct gctgatttta gaccgaaaag atggtttgaa ccatcaacta gaaacttggg	1800
atgggctttc ttgccattca acggtggtcc aagaatctgt ttgggacaac agtttgcttt	1860
gactgaagcc gggtacgttt tgggttagact tgttcaggag tttccaaact tgtcacaaga	1920
ccccgaaacc aagtaccac cacctagatt ggcacacttg acgatgtgct tgtttgacgg	1980
tgcacacgtc aagatgtcat aggtttcccc atacaagtag ttcagtaatt atacactgtt	2040
tttactttct cttcatacca aatggacaaa agttttaagc atgcctaaca acgtgaccgg	2100

acaattgtgt cgactagta tgtaacaatt gtaaaaatag tgtacactaa tttgtggtgg 2160
 ccggagataa attacagttt ggttttgtgt aaactcgcgg atatctctgg cagtttctct 2220
 tctccgcagc agctttgcc aagggtttgct ctggggccaa caaattcaaa agggggagaa 2280
 acttaacacc ccttatctct ccactctagg ttgtagctct tgtggggatg caattgtcgt 2340
 acgtttttta tgttttgtct agactttgat gattacgttg gatttcttat gtctgaggcg 2400
 tgcttgaaag aagtgtcaaa atgtgacagg cgacgtatt cgacatgaac gcgaaagggt 2460
 tatttgcatc aatacgaggg gctgactcta gtctaggatg gcagtcctag gttgcaaaca 2520
 tgttgcacca tatccctcct ggagttggtc gacctgcct acgccaccct cagcgatcgg 2580
 cactttccgt tgttcaatat ttctccttc cattgttcca ggggttatca acaacgttgc 2640
 cggcctcctc cccaaattac aagaaaaata aattgtcgca cggcaccgat ctgtcaaaga 2700
 tacagataaa ccttaaactc gcaaaaacaa gaccctccc catagcctag aagcaccagc 2760
 aagatgatgg agcaactcct ccagtactgg tacatcgac tctctgtatg gttcatcctt 2820
 cgctacttgg cttcccacgc acgagcgcgc tacttgcgcc acaagctcgg cgcggcgcca 2880
 ttcacgcaca ccagtacga cggctggtat gggttcaagt ttgggcggga gtttctcaag 2940
 gcgaagaaga tcgggcggca gacggacttg gtgcatgcgc ggttccgtgg cggcatggac 3000
 accttctoga gctacacttt cggcatccat atcatcctta cccgggaccc ggagaacatc 3060
 aaggcggctc tggcgacgca gttcgatgac ttctcgctcg gtggcaggat caggttcttg 3120
 aagccgttgt tggggtatgg gatattcacg 3150

<210> 93
 <211> 3579
 <212> DNA
 <213> *Candida tropicalis*

<400> 93
 aaaaccgata caagaagaag acagtcaaca agaacgttaa tgtcaaccag gcgccaagaa 60
 gacggtttgg cggacttggga agaattgtggc atttgcccat gatgtttatg ttctggagag 120
 gtttttcaag gaatcgatc cctccgccac cacaagaacc accagttaac gagatccata 180
 ttcacaaccc acgcaagggt gacaatgctc aacaacaaca gcaacaacaa caacccccac 240
 aagaacagtg gaataatgcc agtcaacaaa gagtggtgac agacgaggga gaaaacgcaa 300
 gcaacagtgg ttctgatgca agatcagcta caccgcttca tcaggaaaag caggagctcc 360
 caccaccata tgccatcac gagcaacacc agcaggttag tgtatagtag tctgtagtta 420
 agtcaatgca atgtaccaat aagactatcc cttcttaca ccaagttttc tgccgcgcct 480
 gtctggcaac agatgctggc cgacacactt tcaactgagt ttggtctaga attcttgcac 540

atgcacgaca aggaaactct tacaaaagaca acacttgtgc totgatgcc a ttgatcttg 600
ctaagcctta tcaacgtaat tgagatcatt gtttgtctga attatacaca ccagtggag 660
aatctggtct aatctgcacg cctcatgggc attgtgtgtt ttgggggggg ggggggggg 720
gcacacattt ttagtgcgaa tgtttgtttg ctggttcccc ctccccctc cccctatca 780
tgcccacagg attagttttt tcctcactgg aattcgctgt ccacctgtca acccctcac 840
tgccctgccc tgccctgcac gccctgtgtt ttgtgctgtg gcactccac gctataaaag 900
ccctggcgta cggccaagg ttttcctcac agccaaaaa aaatttgggt gatcctttt 960
ggctgcaagg tttttacca ccaccaccac caccacctca actattcaa caaaggatgc 1020
tcgaccagat cttccattac tggtagattg tcttgccatt gttggtcatt atcaagcaga 1080
tcgtggctca tgccaggacc aattatttga tgaagaagt gggcgctaag ccattcacac 1140
atgtccaact agacgggtgg tttggcttca aatttggccg tgaattcctc aaagctaaaa 1200
gtgctgggag gcaggttgat ttaatcatct cccgtttcca cgataatgag gacactttct 1260
ccagctatgc ttttggcaac catgtggtgt tcaccagga ccccgagaat atcaaggcgc 1320
ttttggcaac ccagtttggg gatttttcat tgggaagcag ggtcaaattc ttcaaaccat 1380
tgttggggta cggatatctc accttgagc gcgaaggctg gaagcacagc agagccatgt 1440
tgagaccaca gtttgccaga gagcaagttg ctcatgtgac gtcgttgga ccacatttcc 1500
agttgttgaa gaagcatatt cttaagcaca aggggtgaata ctttgatata caggaattgt 1560
tcttttagatt taccgttgat tcagcgacgg agttcttatt tggtagtcc gtgcactcct 1620
taagggacga ggaaattggc tacgatacga aggacatggc tgaagaaaga cgcaaatttg 1680
ccgacgcgtt caacaagtcg caagtctatt tgtccaccag agttgcttta cagacattgt 1740
actggttggg caacaacaaa gagttcaagg agtgcaacga cattgtccac aagttcacca 1800
actactatgt tcagaaagcc ttggatgcta cccagagga acttgaaaa caaggcgggt 1860
atgtgttctt gtacgagctt gccaaagcaga cgaaagaccc caatgtgttg cgtgaccagt 1920
ctttgaacat cttgttggct ggaagggaca cactgctgg gttgttgtcc tttgctgtgt 1980
ttgagttggc caggaacca cacatctggg ccaagttgag agaggaaatt gaatcacact 2040
ttgggctggg tgaggactct cgtgttgaag agattacctt tgagagcttg aagagatgtg 2100
agtacttgaa agcgtgttg aacgaaacgt tgagattaca ccaagtgtc ccaagaaacg 2160
caagatttgc gattaaagac acgactttac caagaggcgg tggcccaac ggcaaggatc 2220
ctatcttgat cagaaagaat gaggtggtgc aatactccat ctgggcaact cagacaaatc 2280
ctgcttatta tggcgccgat gctgctgatt ttagaccgga aagatggttt gagccatcaa 2340

ctagaaactt	gggatgggct	tacttgccat	tcaacgggtg	tccaagaatc	tgcttgggac	2400
aacagtttgc	tttgaccgaa	gccggttacg	ttttggttag	acttgttcag	gaattcccta	2460
gcttgtcaca	ggaccccgaa	actgagtacc	caccacctag	attggcacac	ttgacgatgt	2520
gcttgtttga	cggggcatac	gtcaagatgc	aataggtttt	ggtttgactt	tgtttccata	2580
tgcaagtagt	tcagtaatta	cacactaatt	tgtggtggcc	ggcgataaat	taccgtttgg	2640
ttttgtgtaa	aaattcggac	atctctgggt	gtttcccttc	tccgcagcag	ctttgccacg	2700
ggtttgctct	gcggccaaca	aattcgaaag	gggggggggg	gggggagaaa	gttaacaccc	2760
cctgttccca	ccgtaggctg	tagctcttgt	ggggggatgt	aattgtcgta	cgttttcatg	2820
tttgcccag	actttgatga	ttacgtaggc	tttcttatgt	ctaaggcgtg	cttgacacaa	2880
gtgtcaaaaag	gtgacaggcg	acgttattcg	acatgaacgc	aaaagggtaa	tttgcacga	2940
tacgaggggt	tgccctctgg	ctaagaagga	ccccccaggt	tgcaaacaatg	ttgcactgca	3000
tcccactcag	agttgggtga	ccacgcctac	gottaccctc	agcgatcggc	aotttccgtt	3060
gctcaatatt	tctctcccc	ctgcttcccc	ccattgttcc	agggattatc	aacaacgttg	3120
ccggtctcct	ctcccccccc	tccccccagt	tatgtacaag	aaaattaaat	tgtcgcacgg	3180
caccgatagc	tcaaagatac	agagaaacct	taatccctcc	catagcctag	aagcatcaaa	3240
aagatgattg	agcaactcct	ccagtactgg	tacattgcac	tccctgtatg	gttcattctc	3300
cgctacgtgg	cttcccacgc	acgaaccatc	tacttgcgcc	acaagctcgg	cgcggcgcgg	3360
ttcacgcaca	cccagtagca	cggatggtat	gggttcaagt	ttgggcggga	gtttctcaag	3420
gcgaagaaga	ttggaaggca	gacggacttg	gtgcatgcgc	ggttccgtgg	agggggcatg	3480
gatactttct	cgagctatac	tttcggcatc	catatcattc	ttactcggga	cccggagaac	3540
atcaaggcgg	tcttggcgac	gcagttcgat	gacttttcg			3579

<210> 94
 <211> 3348
 <212> DNA
 <213> *Candida tropicalis*

<400>	94					
gatgtggtgc	ttgatttctc	gagacacatc	cttgtgaggt	gcatgaatc	tgtacctgtc	60
tgtaagcaca	gggaactgct	tcaacacctt	attgcatatt	ctgtctattg	caagcgtgtg	120
ctgcaacgat	atctgccaag	gtatatagca	gaaogtgctg	atggttcctc	cggatcatatt	180
ctgttggtag	ttctgcagg	aaatttggat	gtcaggtagt	ggaggaggt	ttgtatcgg	240
tgtgttttct	tcttctctct	tctctgattc	aaactccacg	tctccttcgg	gttctgtgtc	300

tgtgtctgag	tcgtactgtt	ggattaagtc	catcgcatgt	gtgaaaaaaa	gtagcgctta	360
tttagacaac	cagttcgttg	ggcgggtatc	agaaatagtc	tgttgtgcac	gaccatgagt	420
atgcaacttg	acgagacgtc	gtaggaatc	cacagaatga	tagcaggaag	cttactacgt	480
gagagattct	gcttagagga	tggtctcttc	ttgttgattc	cattaggtgg	gtatcatctc	540
cgggtggtgac	aacttgacac	aagcagttcc	gagaaccacc	cacaacaatc	accattccag	600
ctatcacttc	tacatgtcaa	cctacgatgt	atctcatcac	catctagttt	cttggcaatc	660
gtttatttgt	tatgggtcaa	catccaatac	aactccacca	atgaagaaga	aaaacggaaa	720
gcagaatacc	agaatgacag	tgtgagttcc	tgaccattgc	taatctatgg	ctatatctag	780
tttgctatcg	tgggatgtga	tctgtgtcgt	cttcatttgc	gtttgtgttt	atttcgggta	840
tgaatattgt	tatactaaat	acttgatgca	caaacatggc	gctcgagaaa	tcgagaatgt	900
gatcaacgat	gggttctttg	ggttccgctt	acctttgcta	ctcatgcgag	ccagcaatga	960
gggccgactt	atcgagttca	gtgtcaagag	attcgagtcg	gcgccacatc	cacagaacaa	1020
gacattggtc	aaccgggcat	tgagcgttcc	tgtgatactc	accaaggacc	cagtgaatat	1080
caaagcgatg	ctatcgaccc	agtttgatga	cttttccctt	gggttgagac	tacaccagtt	1140
tgcgccgttg	ttggggaaaag	gcatctttac	tttggaaggc	ccagagtgga	agcagagccg	1200
atctatgttg	cgtccgcaat	ttgccaaaaga	tcgggtttct	catatcctgg	atctagaacc	1260
gcattttgtg	ttgcttcgga	agcacattga	tggccacaat	ggagactact	tcgacatcca	1320
ggagctctac	ttccggttct	cgatggatgt	ggcgacgggg	tttttgtttg	gcgagtctgt	1380
ggggtcgttg	aaagacgaag	atgcgaggtt	cctggaagca	ttcaatgagt	cgcagaagta	1440
tttggaact	agggcaacgt	tgacagagtt	gtactttctt	tgtgacgggt	ttaggtttcg	1500
ccagtacaac	aaggttgtgc	gaaagtctctg	cagccagtgt	gtccacaagg	cgttagatgt	1560
tgcaccggaa	gacaccagcg	agtacgtgtt	tctccgcgag	ttggtcaaac	aaactcgaga	1620
tcccgttggt	ttacaagacc	aagcgttgaa	cgtcttgctt	gctggacgcg	acaccaccgc	1680
gtcgttatta	tcgtttgcaa	catttgagct	agcccggaat	gaccacatgt	ggaggaagct	1740
acgagaggag	gttatcctga	cgatgggacc	gtccagtgat	gaaataaccg	tggccggggt	1800
gaagagttgc	cgttacctca	aagcaatcct	aaacgaaact	cttcgactat	acccaagtgt	1860
gcctaggaac	gcgagatttg	ctacgaggaa	tacgacgctt	cctcgtggcg	gaggtccaga	1920
tggatcgttt	ccgattttga	taagaaaggg	ccagccagtg	gggtatttca	tttgtgctac	1980
aaacttgaat	gagaaggtat	atgggaatga	tagccatgtg	tttcgaccgg	agagatgggc	2040
tgcgttagag	ggcaagagtt	tgggctggtc	gtatcttcca	ttcaacggcg	gcccgagaag	2100

ctgccttggt cagcagtttg caatccttga agcttcgtat gttttggctc gattgacaca 2160
 gtgctacacg acgatacagc ttagaactac cgagtaccca ccaaagaaac tcgttcatct 2220
 cacgatgagt cttctcaacg ggggtgtacat ccgaactaga acttgattat gtgtttatgg 2280
 ttaatcgggg caaagcactg caagtcattg atgtttgtgg aagcccagca ttggtgttcc 2340
 ggagcatcaa taaccaatgt cttgaagggt ttgattttct tgaccttctt cttoctgagc 2400
 ttctttccgt caaacttgta cagaatggcc atcatttcag gaacaaccac gtacgacggc 2460
 cggtagcgca tctggagtat ctgcgcgtcg ttcaagtagc acgaaaacag caacgacgtc 2520
 accatctgct tcccaatctt gacacccaca gatacccctg cggcttcatg gatcaaaaac 2580
 gtcggcaacc ccgcgtatat gtccatgtaa ttctccatgg ccacctccat caacacactg 2640
 atggagcgac tgacgggtgcc accactgccc tcggttgagt caaggcagta tgatgccggg 2700
 atccagtact ccaatgggaa cctctgcacg gtgtcgctgc agtttttgag gcgtatttcg 2760
 atccatgata gttctttggt gctgtagtat aacgagctct tgggtgtcctt gaaatggaac 2820
 aggttggaatg tgttggtgag tttgtctgcg tgcttggttt gcaagtcttc gatcgagcgt 2880
 agtgagtaga cagttggcgg ggggtggtgc tcgggcttta ttctgtgttt gtgtttcctt 2940
 cttagtcttg gaatgacgct gttatcgacg gttcgtagta taagtagcgc caatatgaga 3000
 atgtatatcc gcatcaccca agactcttca gcctgttaca acgactgagg ctgttggccg 3060
 tgtgaccaat tgggtttcttt ggtgacctag attggtcccg cagggaaagc aagggtgct 3120
 aggggggcat accaaacaag gtcgtgtaat cagtatctat ggtgctacca tgtgtgtggt 3180
 tggggggaaa ttcccgcat tttgtgtaac gaaagttcta gaaagttctc gtgggttctg 3240
 agaatctgct ggaaccatcc acccgcat ttcgttgcaa agtggaaga gcaatcaacc 3300
 caccctgctt tgcccaatca gccattcccc tgggaatata aattcaac 3348

<210> 95
 <211> 523
 <212> PRT
 <213> CANDIDATROPICALIS

<400> 95

Met Ala Thr Gln Glu Ile Ile Asp Ser Val Leu Pro Tyr Leu Thr Lys
 1 5 10 15

Trp Tyr Thr Val Ile Thr Ala Ala Val Leu Val Phe Leu Ile Ser Thr
 20 25 30

Asn Ile Lys Asn Tyr Val Lys Ala Lys Lys Leu Lys Cys Val Asp Pro
 35 40 45

Pro Tyr Leu Lys Asp Ala Gly Leu Thr Gly Ile Leu Ser Leu Ile Ala
 50 55 60

Ala Ile Lys Ala Lys Asn Asp Gly Arg Leu Ala Asn Phe Ala Asp Glu
 65 70 75 80

Val Phe Asp Glu Tyr Pro Asn His Thr Phe Tyr Leu Ser Val Ala Gly
 85 90 95

Ala Leu Lys Ile Val Met Thr Val Asp Pro Glu Asn Ile Lys Ala Val
 100 105 110

Leu Ala Thr Gln Phe Thr Asp Phe Ser Leu Gly Thr Arg His Ala His
 115 120 125

Phe Ala Pro Leu Leu Gly Asp Gly Ile Phe Thr Leu Asp Gly Glu Gly
 130 135 140

Trp Lys His Ser Arg Ala Met Leu Arg Pro Gln Phe Ala Arg Asp Gln
 145 150 155 160

Ile Gly His Val Lys Ala Leu Glu Pro His Ile Gln Ile Met Ala Lys
 165 170 175

Gln Ile Lys Leu Asn Gln Gly Lys Thr Phe Asp Ile Gln Glu Leu Phe
 180 185 190

Phe Arg Phe Thr Val Asp Thr Ala Thr Glu Phe Leu Phe Gly Glu Ser
 195 200 205

Val His Ser Leu Tyr Asp Glu Lys Leu Gly Ile Pro Thr Pro Asn Glu
 210 215 220

Ile Pro Gly Arg Glu Asn Phe Ala Ala Ala Phe Asn Val Ser Gln His
 225 230 235 240

Tyr Leu Ala Thr Arg Ser Tyr Ser Gln Thr Phe Tyr Phe Leu Thr Asn
 245 250 255

Pro Lys Glu Phe Arg Asp Cys Asn Ala Lys Val His His Leu Ala Lys
 260 265 270

Tyr Phe Val Asn Lys Ala Leu Asn Phe Thr Pro Glu Glu Leu Glu Glu
 275 280 285

Lys Ser Lys Ser Gly Tyr Val Phe Leu Tyr Glu Leu Val Lys Gln Thr
290 295 300

Arg Asp Pro Lys Val Leu Gln Asp Gln Leu Leu Asn Ile Met Val Ala
305 310 315 320

Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Ala Leu Phe Glu Leu
325 330 335

Ala Arg His Pro Glu Met Trp Ser Lys Leu Arg Glu Glu Ile Glu Val
340 345 350

Asn Phe Gly Val Gly Glu Asp Ser Arg Val Glu Glu Ile Thr Phe Glu
355 360 365

Ala Leu Lys Arg Cys Glu Tyr Leu Lys Ala Ile Leu Asn Glu Thr Leu
370 375 380

Arg Met Tyr Pro Ser Val Pro Val Asn Phe Arg Thr Ala Thr Arg Asp
385 390 395 400

Thr Thr Leu Pro Arg Gly Gly Gly Ala Asn Gly Thr Asp Pro Ile Tyr
405 410 415

Ile Pro Lys Gly Ser Thr Val Ala Tyr Val Val Tyr Lys Thr His Arg
420 425 430

Leu Glu Glu Tyr Tyr Gly Lys Asp Ala Asn Asp Phe Arg Pro Glu Arg
435 440 445

Trp Phe Glu Pro Ser Thr Lys Lys Leu Gly Trp Ala Tyr Val Pro Phe
450 455 460

Asn Gly Gly Pro Arg Val Cys Leu Gly Gln Gln Phe Ala Leu Thr Glu
465 470 475 480

Ala Ser Tyr Val Ile Thr Arg Leu Ala Gln Met Phe Glu Thr Val Ser
485 490 495

Ser Asp Pro Gly Leu Glu Tyr Pro Pro Pro Lys Cys Ile His Leu Thr
500 505 510

Met Ser His Asn Asp Gly Val Phe Val Lys Met
515 520

<210> 96
 <211> 522
 <212> PRT
 <213> CANDIDATROPICALIS

<400> 96

Met Thr Val His Asp Ile Ile Ala Thr Tyr Phe Thr Lys Trp Tyr Val
 1 5 10 15

Ile Val Pro Leu Ala Leu Ile Ala Tyr Arg Val Leu Asp Tyr Phe Tyr
 20 25 30

Gly Arg Tyr Leu Met Tyr Lys Leu Gly Ala Lys Pro Phe Phe Gln Lys
 35 40 45

Gln Thr Asp Gly Cys Phe Gly Phe Lys Ala Pro Leu Glu Leu Leu Lys
 50 55 60

Lys Lys Ser Asp Gly Thr Leu Ile Asp Phe Thr Leu Gln Arg Ile His
 65 70 75 80

Asp Leu Asp Arg Pro Asp Ile Pro Thr Phe Thr Phe Pro Val Phe Ser
 85 90 95

Ile Asn Leu Val Asn Thr Leu Glu Pro Glu Asn Ile Lys Ala Ile Leu
 100 105 110

Ala Thr Gln Phe Asn Asp Phe Ser Leu Gly Thr Arg His Ser His Phe
 115 120 125

Ala Pro Leu Leu Gly Asp Gly Ile Phe Thr Leu Asp Gly Ala Gly Trp
 130 135 140

Lys His Ser Arg Ser Met Leu Arg Pro Gln Phe Ala Arg Glu Gln Ile
 145 150 155 160

Ser His Val Lys Leu Leu Glu Pro His Val Gln Val Phe Phe Lys His
 165 170 175

Val Arg Lys Ala Gln Gly Lys Thr Phe Asp Ile Gln Glu Leu Phe Phe
 180 185 190

Arg Leu Thr Val Asp Ser Ala Thr Glu Phe Leu Phe Gly Glu Ser Val
 195 200 205

Glu Ser Leu Arg Asp Glu Ser Ile Gly Met Ser Ile Asn Ala Leu Asp
210 215 220

Phe Asp Gly Lys Ala Gly Phe Ala Asp Ala Phe Asn Tyr Ser Gln Asn
225 230 235 240

Tyr Leu Ala Ser Arg Ala Val Met Gln Gln Leu Tyr Trp Val Leu Asn
245 250 255

Gly Lys Lys Phe Lys Glu Cys Asn Ala Lys Val His Lys Phe Ala Asp
260 265 270

Tyr Tyr Val Asn Lys Ala Leu Asp Leu Thr Pro Glu Gln Leu Glu Lys
275 280 285

Gln Asp Gly Tyr Val Phe Leu Tyr Glu Leu Val Lys Gln Thr Arg Asp
290 295 300

Lys Gln Val Leu Arg Asp Gln Leu Leu Asn Ile Met Val Ala Gly Arg
305 310 315 320

Asp Thr Thr Ala Gly Leu Leu Ser Phe Val Phe Phe Glu Leu Ala Arg
325 330 335

Asn Pro Glu Val Thr Asn Lys Leu Arg Glu Glu Ile Glu Asp Lys Phe
340 345 350

Gly Leu Gly Glu Asn Ala Ser Val Glu Asp Ile Ser Phe Glu Ser Leu
355 360 365

Lys Ser Cys Glu Tyr Leu Lys Ala Val Leu Asn Glu Thr Leu Arg Leu
370 375 380

Tyr Pro Ser Val Pro Gln Asn Phe Arg Val Ala Thr Lys Asn Thr Thr
385 390 395 400

Leu Pro Arg Gly Gly Gly Lys Asp Gly Leu Ser Pro Val Leu Val Arg
405 410 415

Lys Gly Gln Thr Val Ile Tyr Gly Val Tyr Ala Ala His Arg Asn Pro
420 425 430

Ala Val Tyr Gly Lys Asp Ala Leu Glu Phe Arg Pro Glu Arg Trp Phe
435 440 445

Glu Pro Glu Thr Lys Lys Leu Gly Trp Ala Phe Leu Pro Phe Asn Gly
450 455 460

Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe Ala Leu Thr Glu Ala Ser
465 470 475 480

Tyr Val Thr Val Arg Leu Leu Gln Glu Phe Ala His Leu Ser Met Asp
485 490 495

Pro Asp Thr Glu Tyr Pro Pro Lys Lys Met Ser His Leu Thr Met Ser
500 505 510

Leu Phe Asp Gly Ala Asn Ile Glu Met Tyr
515 520

<210> 97

<211> 522

<212> PRT

<213> CANDIDATROPICALIS

<400> 97

Met Thr Ala Gln Asp Ile Ile Ala Thr Tyr Ile Thr Lys Trp Tyr Val
1 5 10 15

Ile Val Pro Leu Ala Leu Ile Ala Tyr Arg Val Leu Asp Tyr Phe Tyr
20 25 30

Gly Arg Tyr Leu Met Tyr Lys Leu Gly Ala Lys Pro Phe Phe Gln Lys
35 40 45

Gln Thr Asp Gly Tyr Phe Gly Phe Lys Ala Pro Leu Glu Leu Leu Lys
50 55 60

Lys Lys Ser Asp Gly Thr Leu Ile Asp Phe Thr Leu Glu Arg Ile Gln
65 70 75 80

Ala Leu Asn Arg Pro Asp Ile Pro Thr Phe Thr Phe Pro Ile Phe Ser
85 90 95

Ile Asn Leu Ile Ser Thr Leu Glu Pro Glu Asn Ile Lys Ala Ile Leu
100 105 110

Ala Thr Gln Phe Asn Asp Phe Ser Leu Gly Thr Arg His Ser His Phe
115 120 125

Ala Pro Leu Leu Gly Asp Gly Ile Phe Thr Leu Asp Gly Ala Gly Trp
130 135 140

Lys His Ser Arg Ser Met Leu Arg Pro Gln Phe Ala Arg Glu Gln Ile
145 150 155 160

Ser His Val Lys Leu Leu Glu Pro His Met Gln Val Phe Phe Lys His
165 170 175

Val Arg Lys Ala Gln Gly Lys Thr Phe Asp Ile Gln Glu Leu Phe Phe
180 185 190

Arg Leu Thr Val Asp Ser Ala Thr Glu Phe Leu Phe Gly Glu Ser Val
195 200 205

Glu Ser Leu Arg Asp Glu Ser Ile Gly Met Ser Ile Asn Ala Leu Asp
210 215 220

Phe Asp Gly Lys Ala Gly Phe Ala Asp Ala Phe Asn Tyr Ser Gln Asn
225 230 235 240

Tyr Leu Ala Ser Arg Ala Val Met Gln Gln Leu Tyr Trp Val Leu Asn
245 250 255

Gly Lys Lys Phe Lys Glu Cys Asn Ala Lys Val His Lys Phe Ala Asp
260 265 270

Tyr Tyr Val Ser Lys Ala Leu Asp Leu Thr Pro Glu Gln Leu Glu Lys
275 280 285

Gln Asp Gly Tyr Val Phe Leu Tyr Glu Leu Val Lys Gln Thr Arg Asp
290 295 300

Arg Gln Val Leu Arg Asp Gln Leu Leu Asn Ile Met Val Ala Gly Arg
305 310 315 320

Asp Thr Thr Ala Gly Leu Leu Ser Phe Val Phe Phe Glu Leu Ala Arg
325 330 335

Asn Pro Glu Val Thr Asn Lys Leu Arg Glu Glu Ile Glu Asp Lys Phe
340 345 350

Gly Leu Gly Glu Asn Ala Arg Val Glu Asp Ile Ser Phe Glu Ser Leu
355 360 365

Lys Ser Cys Glu Tyr Leu Lys Ala Val Leu Asn Glu Thr Leu Arg Leu
370 375 380

Tyr Pro Ser Val Pro Gln Asn Phe Arg Val Ala Thr Lys Asn Thr Thr
385 390 395 400

Leu Pro Arg Gly Gly Gly Lys Asp Gly Leu Ser Pro Val Leu Val Arg
405 410 415

Lys Gly Gln Thr Val Met Tyr Gly Val Tyr Ala Ala His Arg Asn Pro
420 425 430

Ala Val Tyr Gly Lys Asp Ala Leu Glu Phe Arg Pro Glu Arg Trp Phe
435 440 445

Glu Pro Glu Thr Lys Lys Leu Gly Trp Ala Phe Leu Pro Phe Asn Gly
450 455 460

Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe Ala Leu Thr Glu Ala Ser
465 470 475 480

Tyr Val Thr Val Arg Leu Leu Gln Glu Phe Gly His Leu Ser Met Asp
485 490 495

Pro Asn Thr Glu Tyr Pro Pro Arg Lys Met Ser His Leu Thr Met Ser
500 505 510

Leu Phe Asp Gly Ala Asn Ile Glu Met Tyr
515 520

<210> 98
<211> 540
<212> PRT
<213> CANDIDATROPICALIS

<400> 98

Met Ser Ser Ser Pro Ser Phe Ala Gln Glu Val Leu Ala Thr Thr Ser
1 5 10 15

Pro Tyr Ile Glu Tyr Phe Leu Asp Asn Tyr Thr Arg Trp Tyr Tyr Phe
20 25 30

Ile Pro Leu Val Leu Leu Ser Leu Asn Phe Ile Ser Leu Leu His Thr
35 40 45

Arg Tyr Leu Glu Arg Arg Phe His Ala Lys Pro Leu Gly Asn Phe Val
50 55 60

Arg Asp Pro Thr Phe Gly Ile Ala Thr Pro Leu Leu Leu Ile Tyr Leu
65 70 75 80

Lys Ser Lys Gly Thr Val Met Lys Phe Ala Trp Gly Leu Trp Asn Asn
85 90 95

Lys Tyr Ile Val Arg Asp Pro Lys Tyr Lys Thr Thr Gly Leu Arg Ile
100 105 110

Val Gly Leu Pro Leu Ile Glu Thr Met Asp Pro Glu Asn Ile Lys Ala
115 120 125

Val Leu Ala Thr Gln Phe Asn Asp Phe Ser Leu Gly Thr Arg His Asp
130 135 140

Phe Leu Tyr Ser Leu Leu Gly Asp Gly Ile Phe Thr Leu Asp Gly Ala
145 150 155 160

Gly Trp Lys His Ser Arg Thr Met Leu Arg Pro Gln Phe Ala Arg Glu
165 170 175

Gln Val Ser His Val Lys Leu Leu Glu Pro His Val Gln Val Phe Phe
180 185 190

Lys His Val Arg Lys His Arg Gly Gln Thr Phe Asp Ile Gln Glu Leu
195 200 205

Phe Phe Arg Leu Thr Val Asp Ser Ala Thr Glu Phe Leu Phe Gly Glu
210 215 220

Ser Ala Glu Ser Leu Arg Asp Glu Ser Ile Gly Leu Thr Pro Thr Thr
225 230 235 240

Lys Asp Phe Asp Gly Arg Arg Asp Phe Ala Asp Ala Phe Asn Tyr Ser
245 250 255

Gln Thr Tyr Gln Ala Tyr Arg Phe Leu Leu Gln Gln Met Tyr Trp Ile
260 265 270

Leu Asn Gly Ser Glu Phe Arg Lys Ser Ile Ala Val Val His Lys Phe
275 280 285

Ala Asp His Tyr Val Gln Lys Ala Leu Glu Leu Thr Asp Asp Asp Leu
290 295 300

Gln Lys Gln Asp Gly Tyr Val Phe Leu Tyr Glu Leu Ala Lys Gln Thr
305 310 315 320

Arg Asp Pro Lys Val Leu Arg Asp Gln Leu Leu Asn Ile Leu Val Ala
325 330 335

Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Val Phe Tyr Glu Leu
340 345 350

Ser Arg Asn Pro Glu Val Phe Ala Lys Leu Arg Glu Glu Val Glu Asn
355 360 365

Arg Phe Gly Leu Gly Glu Glu Ala Arg Val Glu Glu Ile Ser Phe Glu
370 375 380

Ser Leu Lys Ser Cys Glu Tyr Leu Lys Ala Val Ile Asn Glu Thr Leu
385 390 395 400

Arg Leu Tyr Pro Ser Val Pro His Asn Phe Arg Val Ala Thr Arg Asn
405 410 415

Thr Thr Leu Pro Arg Gly Gly Gly Glu Asp Gly Tyr Ser Pro Ile Val
420 425 430

Val Lys Lys Gly Gln Val Val Met Tyr Thr Val Ile Ala Thr His Arg
435 440 445

Asp Pro Ser Ile Tyr Gly Ala Asp Ala Asp Val Phe Arg Pro Glu Arg
450 455 460

Trp Phe Glu Pro Glu Thr Arg Lys Leu Gly Trp Ala Tyr Val Pro Phe
465 470 475 480

Asn Gly Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe Ala Leu Thr Glu
485 490 495

Ala Ser Tyr Val Thr Val Arg Leu Leu Gln Glu Phe Ala His Leu Ser
500 505 510

Met Asp Pro Asp Thr Glu Tyr Pro Pro Lys Leu Gln Asn Thr Leu Thr
515 520 525

Leu Ser Leu Phe Asp Gly Ala Asp Val Arg Met Tyr
530 535 540

<210> 99

<211> 540
 <212> PRT
 <213> CANDIDATROPICALIS

<400> 99

Met Ser Ser Ser Pro Ser Phe Ala Gln Glu Val Leu Ala Thr Thr Ser
 1 5 10 15

Pro Tyr Ile Glu Tyr Phe Leu Asp Asn Tyr Thr Arg Trp Tyr Tyr Phe
 20 25 30

Ile Pro Leu Val Leu Leu Ser Leu Asn Phe Ile Ser Leu Leu His Thr
 35 40 45

Lys Tyr Leu Glu Arg Arg Phe His Ala Lys Pro Leu Gly Asn Val Val
 50 55 60

Leu Asp Pro Thr Phe Gly Ile Ala Thr Pro Leu Ile Leu Ile Tyr Leu
 65 70 75 80

Lys Ser Lys Gly Thr Val Met Lys Phe Ala Trp Ser Phe Trp Asn Asn
 85 90 95

Lys Tyr Ile Val Lys Asp Pro Lys Tyr Lys Thr Thr Gly Leu Arg Ile
 100 105 110

Val Gly Leu Pro Leu Ile Glu Thr Ile Asp Pro Glu Asn Ile Lys Ala
 115 120 125

Val Leu Ala Thr Gln Phe Asn Asp Phe Ser Leu Gly Thr Arg His Asp
 130 135 140

Phe Leu Tyr Ser Leu Leu Gly Asp Gly Ile Phe Thr Leu Asp Gly Ala
 145 150 155 160

Gly Trp Lys His Ser Arg Thr Met Leu Arg Pro Gln Phe Ala Arg Glu
 165 170 175

Gln Val Ser His Val Lys Leu Leu Glu Pro His Val Gln Val Phe Phe
 180 185 190

Lys His Val Arg Lys His Arg Gly Gln Thr Phe Asp Ile Gln Glu Leu
 195 200 205

Phe Phe Arg Leu Thr Val Asp Ser Ala Thr Glu Phe Leu Phe Gly Glu
 210 215 220

Ser Ala Glu Ser Leu Arg Asp Asp Ser Val Gly Leu Thr Pro Thr Thr
225 230 235 240

Lys Asp Phe Glu Gly Arg Gly Asp Phe Ala Asp Ala Phe Asn Tyr Ser
245 250 255

Gln Thr Tyr Gln Ala Tyr Arg Phe Leu Leu Gln Gln Met Tyr Trp Ile
260 265 270

Leu Asn Gly Ala Glu Phe Arg Lys Ser Ile Ala Ile Val His Lys Phe
275 280 285

Ala Asp His Tyr Val Gln Lys Ala Leu Glu Leu Thr Asp Asp Asp Leu
290 295 300

Gln Lys Gln Asp Gly Tyr Val Phe Leu Tyr Glu Leu Ala Lys Gln Thr
305 310 315 320

Arg Asp Pro Lys Val Leu Arg Asp Gln Leu Leu Asn Ile Leu Val Ala
325 330 335

Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Val Phe Tyr Glu Leu
340 345 350

Ser Arg Asn Pro Glu Val Phe Ala Lys Leu Arg Glu Glu Val Glu Asn
355 360 365

Arg Phe Gly Leu Gly Glu Glu Ala Arg Val Glu Glu Ile Ser Phe Glu
370 375 380

Ser Leu Lys Ser Cys Glu Tyr Leu Lys Ala Val Ile Asn Glu Ala Leu
385 390 395 400

Arg Leu Tyr Pro Ser Val Pro His Asn Phe Arg Val Ala Thr Arg Asn
405 410 415

Thr Thr Leu Pro Arg Gly Gly Gly Lys Asp Gly Cys Ser Pro Ile Val
420 425 430

Val Lys Lys Gly Gln Val Val Met Tyr Thr Val Ile Gly Thr His Arg
435 440 445

Asp Pro Ser Ile Tyr Gly Ala Asp Ala Asp Val Phe Arg Pro Glu Arg
450 455 460

Trp Phe Glu Pro Glu Thr Arg Lys Leu Gly Trp Ala Tyr Val Pro Phe
465 470 475 480

Asn Gly Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe Ala Leu Thr Glu
485 490 495

Ala Ser Tyr Val Thr Val Arg Leu Leu Gln Glu Phe Gly Asn Leu Ser
500 505 510

Leu Asp Pro Asn Ala Glu Tyr Pro Pro Lys Leu Gln Asn Thr Leu Thr
515 520 525

Leu Ser Leu Phe Asp Gly Ala Asp Val Arg Met Phe
530 535 540

<210> 100

<211> 517

<212> PRT

<213> CANDIDATROPICALIS

<400> 100

Met Ile Glu Gln Leu Leu Glu Tyr Trp Tyr Val Val Val Pro Val Leu
1 5 10 15

Tyr Ile Ile Lys Gln Leu Leu Ala Tyr Thr Lys Thr Arg Val Leu Met
20 25 30

Lys Lys Leu Gly Ala Ala Pro Val Thr Asn Lys Leu Tyr Asp Asn Ala
35 40 45

Phe Gly Ile Val Asn Gly Trp Lys Ala Leu Gln Phe Lys Lys Glu Gly
50 55 60

Arg Ala Gln Glu Tyr Asn Asp Tyr Lys Phe Asp His Ser Lys Asn Pro
65 70 75 80

Ser Val Gly Thr Tyr Val Ser Ile Leu Phe Gly Thr Arg Ile Val Val
85 90 95

Thr Lys Asp Pro Glu Asn Ile Lys Ala Ile Leu Ala Thr Gln Phe Gly
100 105 110

Asp Phe Ser Leu Gly Lys Arg His Thr Leu Phe Lys Pro Leu Leu Gly
115 120 125

Asp Gly Ile Phe Thr Leu Asp Gly Glu Gly Trp Lys His Ser Arg Ala
 130 135 140

Met Leu Arg Pro Gln Phe Ala Arg Glu Gln Val Ala His Val Thr Ser
 145 150 155 160

Leu Glu Pro His Phe Gln Leu Leu Lys Lys His Ile Leu Lys His Lys
 165 170 175

Gly Glu Tyr Phe Asp Ile Gln Glu Leu Phe Phe Arg Phe Thr Val Asp
 180 185 190

Ser Ala Thr Glu Phe Leu Phe Gly Glu Ser Val His Ser Leu Lys Asp
 195 200 205

Glu Ser Ile Gly Ile Asn Gln Asp Asp Ile Asp Phe Ala Gly Arg Lys
 210 215 220

Asp Phe Ala Glu Ser Phe Asn Lys Ala Gln Glu Tyr Leu Ala Ile Arg
 225 230 235 240

Thr Leu Val Gln Thr Phe Tyr Trp Leu Val Asn Asn Lys Glu Phe Arg
 245 250 255

Asp Cys Thr Lys Leu Val His Lys Phe Thr Asn Tyr Tyr Val Gln Lys
 260 265 270

Ala Leu Asp Ala Ser Pro Glu Glu Leu Glu Lys Gln Ser Gly Tyr Val
 275 280 285

Phe Leu Tyr Glu Leu Val Lys Gln Thr Arg Asp Pro Asn Val Leu Arg
 290 295 300

Asp Gln Ser Leu Asn Ile Leu Leu Ala Gly Arg Asp Thr Thr Ala Gly
 305 310 315 320

Leu Leu Ser Phe Ala Val Phe Glu Leu Ala Arg His Pro Glu Ile Trp
 325 330 335

Ala Lys Leu Arg Glu Glu Ile Glu Gln Gln Phe Gly Leu Gly Glu Asp
 340 345 350

Ser Arg Val Glu Glu Ile Thr Phe Glu Ser Leu Lys Arg Cys Glu Tyr
 355 360 365

Leu Lys Ala Phe Leu Asn Glu Thr Leu Arg Ile Tyr Pro Ser Val Pro
370 375 380

Arg Asn Phe Arg Ile Ala Thr Lys Asn Thr Thr Leu Pro Arg Gly Gly
385 390 395 400

Gly Ser Asp Gly Thr Ser Pro Ile Leu Ile Gln Lys Gly Glu Ala Val
405 410 415

Ser Tyr Gly Ile Asn Ser Thr His Leu Asp Pro Val Tyr Tyr Gly Pro
420 425 430

Asp Ala Ala Glu Phe Arg Pro Glu Arg Trp Phe Glu Pro Ser Thr Lys
435 440 445

Lys Leu Gly Trp Ala Tyr Leu Pro Phe Asn Gly Gly Pro Arg Ile Cys
450 455 460

Leu Gly Gln Gln Phe Ala Leu Thr Glu Ala Gly Tyr Val Leu Val Arg
465 470 475 480

Leu Val Gln Glu Phe Ser His Val Arg Leu Asp Pro Asp Glu Val Tyr
485 490 495

Pro Pro Lys Arg Leu Thr Asn Leu Thr Met Cys Leu Gln Asp Gly Ala
500 505 510

Ile Val Lys Phe Asp
515

<210> 101

<211> 517

<212> PRT

<213> CANDIDATROPICALIS

<400> 101

Met Ile Glu Gln Ile Leu Glu Tyr Trp Tyr Ile Val Val Pro Val Leu
1 5 10 15

Tyr Ile Ile Lys Gln Leu Ile Ala Tyr Ser Lys Thr Arg Val Leu Met
20 25 30

Lys Gln Leu Gly Ala Ala Pro Ile Thr Asn Gln Leu Tyr Asp Asn Val
35 40 45

Phe Gly Ile Val Asn Gly Trp Lys Ala Leu Gln Phe Lys Lys Glu Gly
50 55 60

Arg Ala Gln Glu Tyr Asn Asp His Lys Phe Asp Ser Ser Lys Asn Pro
 65 70 75 80
 Ser Val Gly Thr Tyr Val Ser Ile Leu Phe Gly Thr Lys Ile Val Val
 85 90 95
 Thr Lys Asp Pro Glu Asn Ile Lys Ala Ile Leu Ala Thr Gln Phe Gly
 100 105 110
 Asp Phe Ser Leu Gly Lys Arg His Ala Leu Phe Lys Pro Leu Leu Gly
 115 120 125
 Asp Gly Ile Phe Thr Leu Asp Gly Glu Gly Trp Lys His Ser Arg Ser
 130 135 140
 Met Leu Arg Pro Gln Phe Ala Arg Glu Gln Val Ala His Val Thr Ser
 145 150 155 160
 Leu Glu Pro His Phe Gln Leu Leu Lys Lys His Ile Leu Lys His Lys
 165 170 175
 Gly Glu Tyr Phe Asp Ile Gln Glu Leu Phe Phe Arg Phe Thr Val Asp
 180 185 190
 Ser Ala Thr Glu Phe Leu Phe Gly Glu Ser Val His Ser Leu Lys Asp
 195 200 205
 Glu Thr Ile Gly Ile Asn Gln Asp Asp Ile Asp Phe Ala Gly Arg Lys
 210 215 220
 Asp Phe Ala Glu Ser Phe Asn Lys Ala Gln Glu Tyr Leu Ser Ile Arg
 225 230 235 240
 Ile Leu Val Gln Thr Phe Tyr Trp Leu Ile Asn Asn Lys Glu Phe Arg
 245 250 255
 Asp Cys Thr Lys Leu Val His Lys Phe Thr Asn Tyr Tyr Val Gln Lys
 260 265 270
 Ala Leu Asp Ala Thr Pro Glu Glu Leu Glu Lys Gln Gly Gly Tyr Val
 275 280 285
 Phe Leu Tyr Glu Leu Val Lys Gln Thr Arg Asp Pro Lys Val Leu Arg
 290 295 300

Asp Gln Ser Leu Asn Ile Leu Leu Ala Gly Arg Asp Thr Thr Ala Gly
305 310 315 320

Leu Leu Ser Phe Ala Val Phe Glu Leu Ala Arg Asn Pro His Ile Trp
325 330 335

Ala Lys Leu Arg Glu Glu Ile Glu Gln Gln Phe Gly Leu Gly Glu Asp
340 345 350

Ser Arg Val Glu Glu Ile Thr Phe Glu Ser Leu Lys Arg Cys Glu Tyr
355 360 365

Leu Lys Ala Phe Leu Asn Glu Thr Leu Arg Val Tyr Pro Ser Val Pro
370 375 380

Arg Asn Phe Arg Ile Ala Thr Lys Asn Thr Thr Leu Pro Arg Gly Gly
385 390 395 400

Gly Pro Asp Gly Thr Gln Pro Ile Leu Ile Gln Lys Gly Glu Gly Val
405 410 415

Ser Tyr Gly Ile Asn Ser Thr His Leu Asp Pro Val Tyr Tyr Gly Pro
420 425 430

Asp Ala Ala Glu Phe Arg Pro Glu Arg Trp Phe Glu Pro Ser Thr Arg
435 440 445

Lys Leu Gly Trp Ala Tyr Leu Pro Phe Asn Gly Gly Pro Arg Ile Cys
450 455 460

Leu Gly Gln Gln Phe Ala Leu Thr Glu Ala Gly Tyr Val Leu Val Arg
465 470 475 480

Leu Val Gln Glu Phe Ser His Ile Arg Leu Asp Pro Asp Glu Val Tyr
485 490 495

Pro Pro Lys Arg Leu Thr Asn Leu Thr Met Cys Leu Gln Asp Gly Ala
500 505 510

Ile Val Lys Phe Asp
515

<210> 102

<211> 512

<212> PRT

<213> CANDIDATROPICALIS

<400> 102

Met Leu Asp Gln Ile Leu His Tyr Trp Tyr Ile Val Leu Pro Leu Leu
1 5 10 15

Ala Ile Ile Asn Gln Ile Val Ala His Val Arg Thr Asn Tyr Leu Met
20 25 30

Lys Lys Leu Gly Ala Lys Pro Phe Thr His Val Gln Arg Asp Gly Trp
35 40 45

Leu Gly Phe Lys Phe Gly Arg Glu Phe Leu Lys Ala Lys Ser Ala Gly
50 55 60

Arg Leu Val Asp Leu Ile Ile Ser Arg Phe His Asp Asn Glu Asp Thr
65 70 75 80

Phe Ser Ser Tyr Ala Phe Gly Asn His Val Val Phe Thr Arg Asp Pro
85 90 95

Glu Asn Ile Lys Ala Leu Leu Ala Thr Gln Phe Gly Asp Phe Ser Leu
100 105 110

Gly Ser Arg Val Lys Phe Phe Lys Pro Leu Leu Gly Tyr Gly Ile Phe
115 120 125

Thr Leu Asp Ala Glu Gly Trp Lys His Ser Arg Ala Met Leu Arg Pro
130 135 140

Gln Phe Ala Arg Glu Gln Val Ala His Val Thr Ser Leu Glu Pro His
145 150 155 160

Phe Gln Leu Leu Lys Lys His Ile Leu Lys His Lys Gly Glu Tyr Phe
165 170 175

Asp Ile Gln Glu Leu Phe Phe Arg Phe Thr Val Asp Ser Ala Thr Glu
180 185 190

Phe Leu Phe Gly Glu Ser Val His Ser Leu Lys Asp Glu Glu Ile Gly
195 200 205

Tyr Asp Thr Lys Asp Met Ser Glu Glu Arg Arg Arg Phe Ala Asp Ala
210 215 220

Phe Asn Lys Ser Gln Val Tyr Val Ala Thr Arg Val Ala Leu Gln Asn
225 230 235 240

Leu Tyr Trp Leu Val Asn Asn Lys Glu Phe Lys Glu Cys Asn Asp Ile
245 250 255

Val His Lys Phe Thr Asn Tyr Tyr Val Gln Lys Ala Leu Asp Ala Thr
260 265 270

Pro Glu Glu Leu Glu Lys Gln Gly Gly Tyr Val Phe Leu Tyr Glu Leu
275 280 285

Val Lys Gln Thr Arg Asp Pro Lys Val Leu Arg Asp Gln Ser Leu Asn
290 295 300

Ile Leu Leu Ala Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Ala
305 310 315 320

Val Phe Glu Leu Ala Arg Asn Pro His Ile Trp Ala Lys Leu Arg Glu
325 330 335

Glu Ile Glu Gln Gln Phe Gly Leu Gly Glu Asp Ser Arg Val Glu Glu
340 345 350

Ile Thr Phe^c Glu Ser Leu Lys Arg Cys Glu Tyr Leu Lys Ala Val Leu
355 360 365

Asn Glu Thr Leu Arg Leu His Pro Ser Val Pro Arg Asn Ala Arg Phe
370 375 380

Ala Ile Lys Asp Thr Thr Leu Pro Arg Gly Gly Gly Pro Asn Gly Lys
385 390 395 400

Asp Pro Ile Leu Ile Arg Lys Asp Glu Val Val Gln Tyr Ser Ile Ser
405 410 415

Ala Thr Gln Thr Asn Pro Ala Tyr Tyr Gly Ala Asp Ala Ala Asp Phe
420 425 430

Arg Pro Glu Arg Trp Phe Glu Pro Ser Thr Arg Asn Leu Gly Trp Ala
435 440 445

Phe Leu Pro Phe Asn Gly Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe
450 455 460

Ala Leu Thr Glu Ala Gly Tyr Val Leu Val Arg Leu Val Gln Glu Phe
465 470 475 480

Pro Asn Leu Ser Gln Asp Pro Glu Thr Lys Tyr Pro Pro Pro Arg Leu
485 490 495

Ala His Leu Thr Met Cys Leu Phe Asp Gly Ala His Val Lys Met Ser
500 505 510

<210> 103

<211> 512

<212> PRT

<213> CANDIDATROPICALIS

<400> 103

Met Leu Asp Gln Ile Phe His Tyr Trp Tyr Ile Val Leu Pro Leu Leu
1 5 10 15

Val Ile Ile Lys Gln Ile Val Ala His Ala Arg Thr Asn Tyr Leu Met
20 25 30

Lys Lys Leu Gly Ala Lys Pro Phe Thr His Val Gln Leu Asp Gly Trp
35 40 45

Phe Gly Phe Lys Phe Gly Arg Glu Phe Leu Lys Ala Lys Ser Ala Gly
50 55 60

Arg Gln Val Asp Leu Ile Ile Ser Arg Phe His Asp Asn Glu Asp Thr
65 70 75 80

Phe Ser Ser Tyr Ala Phe Gly Asn His Val Val Phe Thr Arg Asp Pro
85 90 95

Glu Asn Ile Lys Ala Leu Leu Ala Thr Gln Phe Gly Asp Phe Ser Leu
100 105 110

Gly Ser Arg Val Lys Phe Phe Lys Pro Leu Leu Gly Tyr Gly Ile Phe
115 120 125

Thr Leu Asp Gly Glu Gly Trp Lys His Ser Arg Ala Met Leu Arg Pro
130 135 140

Gln Phe Ala Arg Glu Gln Val Ala His Val Thr Ser Leu Glu Pro His
145 150 155 160

Phe Gln Leu Leu Lys Lys His Ile Leu Lys His Lys Gly Glu Tyr Phe
165 170 175

Asp Ile Gln Glu Leu Phe Phe Arg Phe Thr Val Asp Ser Ala Thr Glu
180 185 190

Phe Leu Phe Gly Glu Ser Val His Ser Leu Arg Asp Glu Glu Ile Gly
195 200 205

Tyr Asp Thr Lys Asp Met Ala Glu Glu Arg Arg Lys Phe Ala Asp Ala
210 215 220

Phe Asn Lys Ser Gln Val Tyr Leu Ser Thr Arg Val Ala Leu Gln Thr
225 230 235 240

Leu Tyr Trp Leu Val Asn Asn Lys Glu Phe Lys Glu Cys Asn Asp Ile
245 250 255

Val His Lys Phe Thr Asn Tyr Tyr Val Gln Lys Ala Leu Asp Ala Thr
260 265 270

Pro Glu Glu Leu Glu Lys Gln Gly Gly Tyr Val Phe Leu Tyr Glu Leu
275 280 285

Ala Lys Gln Thr Lys Asp Pro Asn Val Leu Arg Asp Gln Ser Leu Asn
290 295 300

Ile Leu Leu Ala Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Ala
305 310 315 320

Val Phe Glu Leu Ala Arg Asn Pro His Ile Trp Ala Lys Leu Arg Glu
325 330 335

Glu Ile Glu Ser His Phe Gly Leu Gly Glu Asp Ser Arg Val Glu Glu
340 345 350

Ile Thr Phe Glu Ser Leu Lys Arg Cys Glu Tyr Leu Lys Ala Val Leu
355 360 365

Asn Glu Thr Leu Arg Leu His Pro Ser Val Pro Arg Asn Ala Arg Phe
370 375 380

Ala Ile Lys Asp Thr Thr Leu Pro Arg Gly Gly Gly Pro Asn Gly Lys
385 390 395 400

Asp Pro Ile Leu Ile Arg Lys Asn Glu Val Val Gln Tyr Ser Ile Ser
405 410 415

Ala Thr Gln Thr Asn Pro Ala Tyr Tyr Gly Ala Asp Ala Ala Asp Phe
420 425 430

Arg Pro Glu Arg Trp Phe Glu Pro Ser Thr Arg Asn Leu Gly Trp Ala
435 440 445

Tyr Leu Pro Phe Asn Gly Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe
450 455 460

Ala Leu Thr Glu Ala Gly Tyr Val Leu Val Arg Leu Val Gln Glu Phe
465 470 475 480

Pro Ser Leu Ser Gln Asp Pro Glu Thr Glu Tyr Pro Pro Pro Arg Leu
485 490 495

Ala His Leu Thr Met Cys Leu Phe Asp Gly Ala Tyr Val Lys Met Gln
500 505 510

<210> 104

<211> 499

<212> PRT

<213> CANDIDATROPICALIS

<400> 104

Met Ala Ile Ser Ser Leu Leu Ser Trp Asp Val Ile Cys Val Val Phe
1 5 10 15

Ile Cys Val Cys Val Tyr Phe Gly Tyr Glu Tyr Cys Tyr Thr Lys Tyr
20 25 30

Leu Met His Lys His Gly Ala Arg Glu Ile Glu Asn Val Ile Asn Asp
35 40 45

Gly Phe Phe Gly Phe Arg Leu Pro Leu Leu Leu Met Arg Ala Ser Asn
50 55 60

Glu Gly Arg Leu Ile Glu Phe Ser Val Lys Arg Phe Glu Ser Ala Pro
65 70 75 80

His Pro Gln Asn Lys Thr Leu Val Asn Arg Ala Leu Ser Val Pro Val
85 90 95

Ile Leu Thr Lys Asp Pro Val Asn Ile Lys Ala Met Leu Ser Thr Gln
100 105 110

Phe Asp Asp Phe Ser Leu Gly Leu Arg Leu His Gln Phe Ala Pro Leu
115 120 125

Leu Gly Lys Gly Ile Phe Thr Leu Asp Gly Pro Glu Trp Lys Gln Ser
130 135 140

Arg Ser Met Leu Arg Pro Gln Phe Ala Lys Asp Arg Val Ser His Ile
145 150 155 160

Leu Asp Leu Glu Pro His Phe Val Leu Leu Arg Lys His Ile Asp Gly
165 170 175

His Asn Gly Asp Tyr Phe Asp Ile Gln Glu Leu Tyr Phe Arg Phe Ser
180 185 190

Met Asp Val Ala Thr Gly Phe Leu Phe Gly Glu Ser Val Gly Ser Leu
195 200 205

Lys Asp Glu Asp Ala Arg Phe Leu Glu Ala Phe Asn Glu Ser Gln Lys
210 215 220

Tyr Leu Ala Thr Arg Ala Thr Leu His Glu Leu Tyr Phe Leu Cys Asp
225 230 235 240

Gly Phe Arg Phe Arg Gln Tyr Asn Lys Val Val Arg Lys Phe Cys Ser
245 250 255

Gln Cys Val His Lys Ala Leu Asp Val Ala Pro Glu Asp Thr Ser Glu
260 265 270

Tyr Val Phe Leu Arg Glu Leu Val Lys His Thr Arg Asp Pro Val Val
275 280 285

Leu Gln Asp Gln Ala Leu Asn Val Leu Leu Ala Gly Arg Asp Thr Thr
290 295 300

Ala Ser Leu Leu Ser Phe Ala Thr Phe Glu Leu Ala Arg Asn Asp His
305 310 315 320

Met Trp Arg Lys Leu Arg Glu Glu Val Ile Leu Thr Met Gly Pro Ser
325 330 335

Ser Asp Glu Ile Thr Val Ala Gly Leu Lys Ser Cys Arg Tyr Leu Lys
340 345 350

Ala Ile Leu Asn Glu Thr Leu Arg Leu Tyr Pro Ser Val Pro Arg Asn
355 360 365

Ala Arg Phe Ala Thr Arg Asn Thr Thr Leu Pro Arg Gly Gly Gly Pro
370 375 380

Asp Gly Ser Phe Pro Ile Leu Ile Arg Lys Gly Gln Pro Val Gly Tyr
385 390 395 400

Phe Ile Cys Ala Thr His Leu Asn Glu Lys Val Tyr Gly Asn Asp Ser
405 410 415

His Val Phe Arg Pro Glu Arg Trp Ala Ala Leu Glu Gly Lys Ser Leu
420 425 430

Gly Trp Ser Tyr Leu Pro Phe Asn Gly Gly Pro Arg Ser Cys Leu Gly
435 440 445

Gln Gln Phe Ala Ile Leu Glu Ala Ser Tyr Val Leu Ala Arg Leu Thr
450 455 460

Gln Cys Tyr Thr Thr Ile Gln Leu Arg Thr Thr Glu Tyr Pro Pro Lys
465 470 475 480

Lys Leu Val His Leu Thr Met Ser Leu Leu Asn Gly Val Tyr Ile Arg
485 490 495

Thr Arg Thr

<210> 105

<211> 1712

<212> DNA

<213> Candida tropicalis

<400> 105

ggtaccgagc tcacgagttt tgggattttc gagtttggat tgtttccttt gttgattgaa 60
ttgacgaaac cagaggtttt caagacagat aagattgggt ttatcaaaac gcagtttgaa 120
atattccagt tggttttccaa gatattctga agaagattga cgatttgaaa tttgaagaag 180
tggaagaagat ctggtttggga ttgttgaga atttcaagaa tctcaagatt tactctaacg 240
acgggtacaa cgagaattgt attgaattga tcaagaacat gatcttgggtg ttacagaaca 300
tcaagttctt ggaccagact gagaatgcca cagatatata aggcgtcatg tgataaaatg 360
gatgagattt atcccacaat tgaagaaaga gtttatggaa agtgggtcaac cagaagctaa 420

acaggaagaa gcaaacgaag aggtgaaaca agaagaagaa ggtaaataag tattttgtat 480
 tatataacaa acaaagtaag gaatacagat ttatacaata aattggcata ctagtcacgt 540
 gagatatctc atccattccc caactcccaa gaaaaaaaaa aagtgaaaaa aaaaatcaaa 600
 cccaaagatc aacctcccca tcatcatcgt catcaaacc ccagctcaat tcgcaatggt 660
 tagcacaaaa acatacacag aaagggcatc agcacacccc tccaagggtg cccaacgttt 720
 attccgctta atggagtcca aaaagaccaa cctctgcgcc tcgatcgacg tgaccacaac 780
 cgccgagttc ctttcgctca tcgacaagct cggccccac atctgtctcg tgaagacgca 840
 catcgatatc atctcagact tcagctacga gggcacgatt gagccgttgc ttgtgcttgc 900
 agagcgccac ggggttcttga tattcgagga caggaagttt gctgatatcg gaaacacggt 960
 gatgttgcag tacacctcgg gggataaccg gatcgcgccg tggagtgaca tcacgaacgc 1020
 gcacggagtg actgggaagg gcgtcgttga agggttgaaa cgcggtgcgg agggggtaga 1080
 aaaggaaagg ggcgtgttga tgttgccgga gttgtcgagt aaaggctcgt tggcgcatgg 1140
 tgaatatacc cgtgagacga tcgagattgc gaagagtgat cgggagttcg tgattgggtt 1200
 catcgcgag cgggacatgg ggggtagaga agaagggttt gattggatca tcatgacgcc 1260
 tgggtgtggg ttggatgata aaggcgatgc gttgggccag cagtatagga ctggtgatga 1320
 ggtggttctg actggtaccg atgtgattat tgtcgggaga gggttgtttg gaaaaggaag 1380
 agaccctgag gtggagggaa agagatacag ggatgctgga tggaaggcat acttgaagag 1440
 aactggtcag ttagaataaa tattgtaata aataggtcta tatacataca ctaagcttct 1500
 aggacgtcat tgtagtcttc gaagttgtct gctagtttag ttctcatgat ttcgaaaacc 1560
 aataacgcaa tggatgtagc agggatggtg gttagtgcgt tcctgacaaa cccagagtac 1620
 gccgcctcaa accacgtcac attcgccctt tgcctcatcc gcatcacttg cttgaaggta 1680
 tccacgtacg agttgtaata caccttgaag aa 1712

<210> 106
 <211> 267
 <212> PRT
 <213> CANDIDATROPICALIS
 <400> 106

Met Val Ser Thr Lys Thr Tyr Thr Glu Arg Ala Ser Ala His Pro Ser
 1 5 10 15

Lys Val Ala Gln Arg Leu Phe Arg Leu Met Glu Ser Lys Lys Thr Asn
 20 25 30

Leu Cys Ala Ser Ile Asp Val Thr Thr Thr Ala Glu Phe Leu Ser Leu
 35 40 45

Ile Asp Lys Leu Gly Pro His Ile Cys Leu Val Lys Thr His Ile Asp
50 55 60

Ile Ile Ser Asp Phe Ser Tyr Glu Gly Thr Ile Glu Pro Leu Leu Val
65 70 75 80

Leu Ala Glu Arg His Gly Phe Leu Ile Phe Glu Asp Arg Lys Phe Ala
85 90 95

Asp Ile Gly Asn Thr Val Met Leu Gln Tyr Thr Ser Gly Val Tyr Arg
100 105 110

Ile Ala Ala Trp Ser Asp Ile Thr Asn Ala His Gly Val Thr Gly Lys
115 120 125

Gly Val Val Glu Gly Leu Lys Arg Gly Ala Glu Gly Val Glu Lys Glu
130 135 140

Arg Gly Val Leu Met Leu Ala Glu Leu Ser Ser Lys Gly Ser Leu Ala
145 150 155 160

His Gly Glu Tyr Thr Arg Glu Thr Ile Glu Ile Ala Lys Ser Asp Arg
165 170 175

Glu Phe Val Ile Gly Phe Ile Ala Gln Arg Asp Met Gly Gly Arg Glu
180 185 190

Glu Gly Phe Asp Trp Ile Ile Met Thr Pro Gly Val Gly Leu Asp Asp
195 200 205

Lys Gly Asp Ala Leu Gly Gln Gln Tyr Arg Thr Val Asp Glu Val Val
210 215 220

Leu Thr Gly Thr Asp Val Ile Ile Val Gly Arg Gly Leu Phe Gly Lys
225 230 235 240

Gly Arg Asp Pro Glu Val Glu Gly Lys Arg Tyr Arg Asp Ala Gly Trp
245 250 255

Lys Ala Tyr Leu Lys Arg Thr Gly Gln Leu Glu
260 265

<210> 107
<211> 473

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 107
gtcaaagcaa attggtggcc caagcagact cttggaccac cgttgaatgg aacataagcc 60
cagcccaact tcttagtaga tgggtcaaac catctttctg gtctgaagtc gttagcgtcc 120
ttaccgtagt attcttccaa acggtgggtc ttgtagacaa cgtaagcaac agtggagcct 180
ttaggaatgt agattgggtc ggtaccgtta gcaaccaccac ctcttggcaa agtgggtgtct 240
ctggtggcgg ttctaaagtt gacaggaaca gatgggtaca tacgcaaggt ttcgttaagg 300
atagccttca agtattcaca tctcttcaag gcttcgaaag taatttcttc aacgcgggag 360
tcttcaccaa caccaaagtt aacttcgatt tcttctctca acttggacca catctctggg 420
tgtctagcca attcaaaca agcaaaggac aacaaacccg cgggtggtgtc tct 473

<210> 108
<211> 540
<212> DNA
<213> *Candida tropicalis*

<400> 108
tactaacttg ttgaggatct tataaccata cagcaacacg gtcacaacat gtagtagttt 60
gttgaggaac gtatgtgttt ctgagcgcag aactactttt tcaaccacg acgaggtcag 120
tgtttgttca acatgctgtt gcgaaagcca tagcagttac ctaccttccg agagggtcaag 180
ttctttctcc cgtcccagat tctcatgttg ctaatgttca aactggtgag gttcttgggt 240
tcgcaccctg ggatgcagtc ataagaaaag ccgtggctct agcagcactg gtttctaggt 300
ctcttatagt ttcgataaaa ccgttgggtc aaaccactaa aaagaaaccc gttctccgtg 360
tgagaaaaat tcggaacaaa tccactaccc tagaagtgtg acctgccgt tccgaccttc 420
gtgtcgtctc ggtacaactc tgggtgtcaa cggctctctg ttcaacgagt aactgcagc 480
aaccttggtg tgaagggtcaa caacttcttc gtataagaat tcgtgttccc acttatgaaa 540

<210> 109
<211> 29
<212> DNA
<213> Bacteriophage T7

<400> 109
ggatccta atcagactcact atagggagg 29

<210> 110
<211> 523

<212> PRT
 <213> CANDIDATROPICALIS

<400> 110

Met Ala Thr Gln Glu Ile Ile Asp Ser Val Leu Pro Tyr Leu Thr Lys
 1 5 10 15

Trp Tyr Thr Val Ile Thr Ala Ala Val Leu Val Phe Leu Ile Ser Thr
 20 25 30

Asn Ile Lys Asn Tyr Val Lys Ala Lys Lys Leu Lys Cys Val Asp Pro
 35 40 45

Pro Tyr Leu Lys Asp Ala Gly Leu Thr Gly Ile Ser Ser Leu Ile Ala
 50 55 60

Ala Ile Lys Ala Lys Asn Asp Gly Arg Leu Ala Asn Phe Ala Asp Glu
 65 70 75 80

Val Phe Asp Glu Tyr Pro Asn His Thr Phe Tyr Leu Ser Val Ala Gly
 85 90 95

Ala Leu Lys Ile Val Met Thr Val Asp Pro Glu Asn Ile Lys Ala Val
 100 105 110

Leu Ala Thr Gln Phe Thr Asp Phe Ser Leu Gly Thr Arg His Ala His
 115 120 125

Phe Ala Pro Leu Leu Gly Asp Gly Ile Phe Thr Leu Asp Gly Glu Gly
 130 135 140

Trp Lys His Ser Arg Ala Met Leu Arg Pro Gln Phe Ala Arg Asp Gln
 145 150 155 160

Ile Gly His Val Lys Ala Leu Glu Pro His Ile Gln Ile Met Ala Lys
 165 170 175

Gln Ile Lys Leu Asn Gln Gly Lys Thr Phe Asp Ile Gln Glu Leu Phe
 180 185 190

Phe Arg Phe Thr Val Asp Thr Ala Thr Glu Phe Leu Phe Gly Glu Ser
 195 200 205

Val His Ser Leu Tyr Asp Glu Lys Leu Gly Ile Pro Thr Pro Asn Glu
 210 215 220

Ile Pro Gly Arg Glu Asn Phe Ala Ala Ala Phe Asn Val Ser Gln His
225 230 235 240

Tyr Leu Ala Thr Arg Ser Tyr Ser Gln Thr Phe Tyr Phe Leu Thr Asn
245 250 255

Pro Lys Glu Phe Arg Asp Cys Asn Ala Lys Val His His Leu Ala Lys
260 265 270

Tyr Phe Val Asn Lys Ala Leu Asn Phe Thr Pro Glu Glu Leu Glu Glu
275 280 285

Lys Ser Lys Ser Gly Tyr Val Phe Leu Tyr Glu Leu Val Lys Gln Thr
290 295 300

Arg Asp Pro Lys Val Leu Gln Asp Gln Leu Leu Asn Ile Met Val Ala
305 310 315 320

Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Ala Leu Phe Glu Leu
325 330 335

Ala Arg His Pro Glu Met Trp Ser Lys Leu Arg Glu Glu Ile Glu Val
340 345 350

Asn Phe Gly Val Gly Glu Asp Ser Arg Val Glu Glu Ile Thr Phe Glu
355 360 365

Ala Leu Lys Arg Cys Glu Tyr Leu Lys Ala Ile Leu Asn Glu Thr Leu
370 375 380

Arg Met Tyr Pro Ser Val Pro Val Asn Phe Arg Thr Ala Thr Arg Asp
385 390 395 400

Thr Thr Leu Pro Arg Gly Gly Gly Ala Asn Gly Thr Asp Pro Ile Tyr
405 410 415

Ile Pro Lys Gly Ser Thr Val Ala Tyr Val Val Tyr Lys Thr His Arg
420 425 430

Leu Glu Glu Tyr Tyr Gly Lys Asp Ala Asn Asp Phe Arg Pro Glu Arg
435 440 445

Trp Phe Glu Pro Ser Thr Lys Lys Leu Gly Trp Ala Tyr Val Pro Phe
450 455 460

Asn Gly Gly Pro Arg Val Cys Leu Gly Gln Gln Phe Ala Leu Thr Glu
465 470 475 480

Ala Ser Tyr Val Ile Thr Arg Leu Ala Gln Met Phe Glu Thr Val Ser
485 490 495

Ser Asp Pro Gly Leu Glu Tyr Pro Pro Pro Lys Cys Ile His Leu Thr
500 505 510

Met Ser His Asn Asp Gly Val Phe Val Lys Met
515 520

<210> 111

<211> 540

<212> PRT

<213> CANDIDATROPICALIS

<400> 111

Met Ser Ser Ser Pro Ser Phe Ala Gln Glu Val Leu Ala Thr Thr Ser
1 5 10 15

Pro Tyr Ile Glu Tyr Phe Leu Asp Asn Tyr Thr Arg Trp Tyr Tyr Phe
20 25 30

Ile Pro Leu Val Leu Leu Ser Leu Asn Phe Ile Ser Leu Leu His Thr
35 40 45

Lys Tyr Leu Glu Arg Arg Phe His Ala Lys Pro Leu Gly Asn Val Val
50 55 60

Leu Asp Pro Thr Phe Gly Ile Ala Thr Pro Leu Ile Leu Ile Tyr Leu
65 70 75 80

Lys Ser Lys Gly Thr Val Met Lys Phe Ala Trp Ser Phe Trp Asn Asn
85 90 95

Lys Tyr Ile Val Lys Asp Pro Lys Tyr Lys Thr Thr Gly Leu Arg Ile
100 105 110

Val Gly Leu Pro Leu Ile Glu Thr Ile Asp Pro Glu Asn Ile Lys Ala
115 120 125

Val Leu Ala Thr Gln Phe Asn Asp Phe Ser Leu Gly Thr Arg His Asp
130 135 140

Phe Leu Tyr Ser Leu Leu Gly Asp Gly Ile Phe Thr Leu Asp Gly Ala
145 150 155 160

Gly Trp Lys His Ser Arg Thr Met Leu Arg Pro Gln Phe Ala Arg Glu
165 170 175

Gln Val Ser His Val Lys Leu Leu Glu Pro His Val Gln Val Phe Phe
180 185 190

Lys His Val Arg Lys His Arg Gly Gln Thr Phe Asp Ile Gln Glu Leu
195 200 205

Phe Phe Arg Leu Thr Val Asp Ser Ala Thr Glu Phe Leu Phe Gly Glu
210 215 220

Ser Ala Glu Ser Leu Arg Asp Asp Ser Val Gly Leu Thr Pro Thr Thr
225 230 235 240

Lys Asp Phe Glu Gly Arg Gly Asp Phe Ala Asp Ala Phe Asn Tyr Ser
245 250 255

Gln Thr Tyr Gln Ala Tyr Arg Phe Leu Leu Gln Gln Met Tyr Trp Ile
260 265 270

Leu Asn Gly Ala Glu Phe Arg Lys Ser Ile Ala Ile Val His Lys Phe
275 280 285

Ala Asp His Tyr Val Gln Lys Ala Leu Glu Leu Thr Asp Asp Asp Leu
290 295 300

Gln Lys Gln Asp Gly Tyr Val Phe Leu Tyr Glu Leu Ala Lys Gln Thr
305 310 315 320

Arg Asp Pro Lys Val Leu Arg Asp Gln Leu Leu Asn Ile Leu Val Ala
325 330 335

Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Val Phe Tyr Glu Leu
340 345 350

Ser Arg Asn Pro Glu Val Phe Ala Lys Leu Arg Glu Glu Val Glu Asn
355 360 365

Arg Phe Gly Leu Gly Glu Glu Ala Arg Val Glu Glu Ile Ser Phe Glu
370 375 380

Ser Leu Lys Ser Cys Glu Tyr Leu Lys Ala Val Ile Asn Glu Ala Leu
385 390 395 400

Arg Leu Tyr Pro Ser Val Pro His Asn Phe Arg Val Ala Thr Arg Asn
405 410 415

Thr Thr Leu Pro Arg Gly Gly Gly Lys Asp Gly Cys Ser Pro Ile Val
420 425 430

Val Lys Lys Gly Gln Val Val Met Tyr Thr Val Ile Gly Thr His Arg
435 440 445

Asp Pro Ser Ile Tyr Gly Ala Asp Ala Asp Val Phe Arg Pro Glu Arg
450 455 460

Trp Phe Glu Pro Glu Thr Arg Lys Leu Gly Trp Ala Tyr Val Pro Phe
465 470 475 480

Asn Gly Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe Ala Leu Thr Glu
485 490 495

Ala Ser Tyr Val Thr Val Arg Leu Leu Gln Glu Phe Gly Asn Leu Ser
500 505 510

Ser Asp Pro Asn Ala Glu Tyr Pro Pro Lys Leu Gln Asn Thr Leu Thr
515 520 525

Leu Ser Leu Phe Asp Gly Ala Asp Val Arg Met Phe
530 535 540

<210> 112

<211> 517

<212> PRT

<213> CANDIDATROPICALIS

<400> 112

Met Ile Glu Gln Leu Leu Glu Tyr Trp Tyr Val Val Val Pro Val Leu
1 5 10 15

Tyr Ile Ile Lys Gln Leu Leu Ala Tyr Thr Lys Thr Arg Val Leu Met
20 25 30

Lys Lys Leu Gly Ala Ala Pro Val Thr Asn Lys Leu Tyr Asp Asn Ala
35 40 45

Phe Gly Ile Val Asn Gly Trp Lys Ala Leu Gln Phe Lys Lys Glu Gly
50 55 60

Arg Ala Gln Glu Tyr Asn Asp Tyr Lys Phe Asp His Ser Lys Asn Pro
65 70 75 80

Ser Val Gly Thr Tyr Val Ser Ile Leu Phe Gly Thr Arg Ile Val Val
85 90 95

Thr Lys Asp Pro Glu Asn Ile Lys Ala Ile Leu Ala Thr Gln Phe Gly
100 105 110

Asp Phe Ser Leu Gly Lys Arg His Thr Leu Phe Lys Pro Leu Leu Gly
115 120 125

Asp Gly Ile Phe Thr Leu Asp Gly Glu Gly Trp Lys His Ser Arg Ala
130 135 140

Met Leu Arg Pro Gln Phe Ala Arg Glu Gln Val Ala His Val Thr Ser
145 150 155 160

Leu Glu Pro His Phe Gln Leu Leu Lys Lys His Ile Leu Lys His Lys
165 170 175

Gly Glu Tyr Phe Asp Ile Gln Glu Leu Phe Phe Arg Phe Thr Val Asp
180 185 190

Ser Ala Thr Glu Phe Leu Phe Gly Glu Ser Val His Ser Leu Lys Asp
195 200 205

Glu Ser Ile Gly Ile Asn Gln Asp Asp Ile Asp Phe Ala Gly Arg Lys
210 215 220

Asp Phe Ala Glu Ser Phe Asn Lys Ala Gln Glu Tyr Leu Ala Ile Arg
225 230 235 240

Thr Leu Val Gln Thr Phe Tyr Trp Leu Val Asn Asn Lys Glu Phe Arg
245 250 255

Asp Cys Thr Lys Ser Val His Lys Phe Thr Asn Tyr Tyr Val Gln Lys
260 265 270

Ala Leu Asp Ala Ser Pro Glu Glu Leu Glu Lys Gln Ser Gly Tyr Val
275 280 285

Phe Leu Tyr Glu Leu Val Lys Gln Thr Arg Asp Pro Asn Val Leu Arg
290 295 300

Asp Gln Ser Leu Asn Ile Leu Leu Ala Gly Arg Asp Thr Thr Ala Gly
305 310 315 320

Leu Leu Ser Phe Ala Val Phe Glu Leu Ala Arg His Pro Glu Ile Trp
325 330 335

Ala Lys Leu Arg Glu Glu Ile Glu Gln Gln Phe Gly Leu Gly Glu Asp
340 345 350

Ser Arg Val Glu Glu Ile Thr Phe Glu Ser Leu Lys Arg Cys Glu Tyr
355 360 365

Leu Lys Ala Phe Leu Asn Glu Thr Leu Arg Ile Tyr Pro Ser Val Pro
370 375 380

Arg Asn Phe Arg Ile Ala Thr Lys Asn Thr Thr Leu Pro Arg Gly Gly
385 390 395 400

Gly Ser Asp Gly Thr Ser Pro Ile Leu Ile Gln Lys Gly Glu Ala Val
405 410 415

Ser Tyr Gly Ile Asn Ser Thr His Leu Asp Pro Val Tyr Tyr Gly Pro
420 425 430

Asp Ala Ala Glu Phe Arg Pro Glu Arg Trp Phe Glu Pro Ser Thr Lys
435 440 445

Lys Leu Gly Trp Ala Tyr Leu Pro Phe Asn Gly Gly Pro Arg Ile Cys
450 455 460

Leu Gly Gln Gln Phe Ala Leu Thr Glu Ala Gly Tyr Val Leu Val Arg
465 470 475 480

Leu Val Gln Glu Phe Ser His Val Arg Ser Asp Pro Asp Glu Val Tyr
485 490 495

Pro Pro Lys Arg Leu Thr Asn Leu Thr Met Cys Leu Gln Asp Gly Ala
500 505 510

Ile Val Lys Phe Asp
515

<210> 113

<211> 517

<212> PRT

<213> CANDIDATROPICALIS

<400> 113

Met Ile Glu Gln Ile Leu Glu Tyr Trp Tyr Ile Val Val Pro Val Leu
1 5 10 15

Tyr Ile Ile Lys Gln Leu Ile Ala Tyr Ser Lys Thr Arg Val Leu Met
20 25 30

Lys Gln Leu Gly Ala Ala Pro Ile Thr Asn Gln Leu Tyr Asp Asn Val
35 40 45

Phe Gly Ile Val Asn Gly Trp Lys Ala Leu Gln Phe Lys Lys Glu Gly
50 55 60

Arg Ala Gln Glu Tyr Asn Asp His Lys Phe Asp Ser Ser Lys Asn Pro
65 70 75 80

Ser Val Gly Thr Tyr Val Ser Ile Leu Phe Gly Thr Lys Ile Val Val
85 90 95

Thr Lys Asp Pro Glu Asn Ile Lys Ala Ile Leu Ala Thr Gln Phe Gly
100 105 110

Asp Phe Ser Leu Gly Lys Arg His Ala Leu Phe Lys Pro Leu Leu Gly
115 120 125

Asp Gly Ile Phe Thr Leu Asp Gly Glu Gly Trp Lys His Ser Arg Ser
130 135 140

Met Leu Arg Pro Gln Phe Ala Arg Glu Gln Val Ala His Val Thr Ser
145 150 155 160

Leu Glu Pro His Phe Gln Leu Leu Lys Lys His Ile Leu Lys His Lys
165 170 175

Gly Glu Tyr Phe Asp Ile Gln Glu Leu Phe Phe Arg Phe Thr Val Asp
180 185 190

Ser Ala Thr Glu Phe Leu Phe Gly Glu Ser Val His Ser Leu Lys Asp
195 200 205

Glu Thr Ile Gly Ile Asn Gln Asp Asp Ile Asp Phe Ala Gly Arg Lys
210 215 220

Asp Phe Ala Glu Ser Phe Asn Lys Ala Gln Glu Tyr Leu Ser Ile Arg
225 230 235 240

Ile Leu Val Gln Thr Phe Tyr Trp Leu Ile Asn Asn Lys Glu Phe Arg
 245 250 255

Asp Cys Thr Lys Ser Val His Lys Phe Thr Asn Tyr Tyr Val Gln Lys
 260 265 270

Ala Leu Asp Ala Thr Pro Glu Glu Leu Glu Lys Gln Gly Gly Tyr Val
 275 280 285

Phe Leu Tyr Glu Leu Val Lys Gln Thr Arg Asp Pro Lys Val Leu Arg
 290 295 300

Asp Gln Ser Leu Asn Ile Leu Leu Ala Gly Arg Asp Thr Thr Ala Gly
 305 310 315 320

Leu Leu Ser Phe Ala Val Phe Glu Leu Ala Arg Asn Pro His Ile Trp
 325 330 335

Ala Lys Leu Arg Glu Glu Ile Glu Gln Gln Phe Gly Leu Gly Glu Asp
 340 345 350

Ser Arg Val Glu Glu Ile Thr Phe Glu Ser Leu Lys Arg Cys Glu Tyr
 355 360 365

Leu Lys Ala Phe Leu Asn Glu Thr Leu Arg Val Tyr Pro Ser Val Pro
 370 375 380

Arg Asn Phe Arg Ile Ala Thr Lys Asn Thr Thr Leu Pro Arg Gly Gly
 385 390 395 400

Gly Pro Asp Gly Thr Gln Pro Ile Leu Ile Gln Lys Gly Glu Gly Val
 405 410 415

Ser Tyr Gly Ile Asn Ser Thr His Leu Asp Pro Val Tyr Tyr Gly Pro
 420 425 430

Asp Ala Ala Glu Phe Arg Pro Glu Arg Trp Phe Glu Pro Ser Thr Arg
 435 440 445

Lys Leu Gly Trp Ala Tyr Leu Pro Phe Asn Gly Gly Pro Arg Ile Cys
 450 455 460

Leu Gly Gln Gln Phe Ala Leu Thr Glu Ala Gly Tyr Val Leu Val Arg
 465 470 475 480

Leu Val Gln Glu Phe Ser His Ile Arg Ser Asp Pro Asp Glu Val Tyr
 485 490 495

Pro Pro Lys Arg Leu Thr Asn Leu Thr Met Cys Leu Gln Asp Gly Ala
 500 505 510

Ile Val Lys Phe Asp
 515

<210> 114
 <211> 512
 <212> PRT
 <213> CANDIDATROPICALIS

<400> 114

Met Leu Asp Gln Ile Leu His Tyr Trp Tyr Ile Val Leu Pro Leu Leu
 1 5 10 15

Ala Ile Ile Asn Gln Ile Val Ala His Val Arg Thr Asn Tyr Leu Met
 20 25 30

Lys Lys Leu Gly Ala Lys Pro Phe Thr His Val Gln Arg Asp Gly Trp
 35 40 45

Leu Gly Phe Lys Phe Gly Arg Glu Phe Leu Lys Ala Lys Ser Ala Gly
 50 55 60

Arg Ser Val Asp Leu Ile Ile Ser Arg Phe His Asp Asn Glu Asp Thr
 65 70 75 80

Phe Ser Ser Tyr Ala Phe Gly Asn His Val Val Phe Thr Arg Asp Pro
 85 90 95

Glu Asn Ile Lys Ala Leu Leu Ala Thr Gln Phe Gly Asp Phe Ser Leu
 100 105 110

Gly Ser Arg Val Lys Phe Phe Lys Pro Leu Leu Gly Tyr Gly Ile Phe
 115 120 125

Thr Leu Asp Ala Glu Gly Trp Lys His Ser Arg Ala Met Leu Arg Pro
 130 135 140

Gln Phe Ala Arg Glu Gln Val Ala His Val Thr Ser Leu Glu Pro His
 145 150 155 160

Phe Gln Leu Leu Lys Lys His Ile Leu Lys His Lys Gly Glu Tyr Phe
 165 170 175

Asp Ile Gln Glu Leu Phe Phe Arg Phe Thr Val Asp Ser Ala Thr Glu
 180 185 190

Phe Leu Phe Gly Glu Ser Val His Ser Leu Lys Asp Glu Glu Ile Gly
 195 200 205

Tyr Asp Thr Lys Asp Met Ser Glu Glu Arg Arg Arg Phe Ala Asp Ala
 210 215 220

Phe Asn Lys Ser Gln Val Tyr Val Ala Thr Arg Val Ala Leu Gln Asn
 225 230 235 240

Leu Tyr Trp Leu Val Asn Asn Lys Glu Phe Lys Glu Cys Asn Asp Ile
 245 250 255

Val His Lys Phe Thr Asn Tyr Tyr Val Gln Lys Ala Leu Asp Ala Thr
 260 265 270

Pro Glu Glu Leu Glu Lys Gln Gly Gly Tyr Val Phe Leu Tyr Glu Leu
 275 280 285

Val Lys Gln Thr Arg Asp Pro Lys Val Leu Arg Asp Gln Ser Leu Asn
 290 295 300

Ile Leu Leu Ala Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Ala
 305 310 315 320

Val Phe Glu Leu Ala Arg Asn Pro His Ile Trp Ala Lys Leu Arg Glu
 325 330 335

Glu Ile Glu Gln Gln Phe Gly Leu Gly Glu Asp Ser Arg Val Glu Glu
 340 345 350

Ile Thr Phe Glu Ser Leu Lys Arg Cys Glu Tyr Leu Lys Ala Val Leu
 355 360 365

Asn Glu Thr Leu Arg Leu His Pro Ser Val Pro Arg Asn Ala Arg Phe
 370 375 380

Ala Ile Lys Asp Thr Thr Leu Pro Arg Gly Gly Gly Pro Asn Gly Lys
 385 390 395 400

Asp Pro Ile Leu Ile Arg Lys Asp Glu Val Val Gln Tyr Ser Ile Ser
405 410 415

Ala Thr Gln Thr Asn Pro Ala Tyr Tyr Gly Ala Asp Ala Ala Asp Phe
420 425 430

Arg Pro Glu Arg Trp Phe Glu Pro Ser Thr Arg Asn Leu Gly Trp Ala
435 440 445

Phe Leu Pro Phe Asn Gly Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe
450 455 460

Ala Leu Thr Glu Ala Gly Tyr Val Leu Val Arg Leu Val Gln Glu Phe
465 470 475 480

Pro Asn Leu Ser Gln Asp Pro Glu Thr Lys Tyr Pro Pro Pro Arg Leu
485 490 495

Ala His Leu Thr Met Cys Leu Phe Asp Gly Ala His Val Lys Met Ser
500 505 510

<210> 115
<211> 512
<212> PRT
<213> CANDIDATROPICALIS
<400> 115

Met Leu Asp Gln Ile Phe His Tyr Trp Tyr Ile Val Leu Pro Leu Leu
1 5 10 15

Val Ile Ile Lys Gln Ile Val Ala His Ala Arg Thr Asn Tyr Leu Met
20 25 30

Lys Lys Leu Gly Ala Lys Pro Phe Thr His Val Gln Leu Asp Gly Trp
35 40 45

Phe Gly Phe Lys Phe Gly Arg Glu Phe Leu Lys Ala Lys Ser Ala Gly
50 55 60

Arg Gln Val Asp Leu Ile Ile Ser Arg Phe His Asp Asn Glu Asp Thr
65 70 75 80

Phe Ser Ser Tyr Ala Phe Gly Asn His Val Val Phe Thr Arg Asp Pro
85 90 95

Glu Asn Ile Lys Ala Leu Leu Ala Thr Gln Phe Gly Asp Phe Ser Leu
100 105 110

Gly Ser Arg Val Lys Phe Phe Lys Pro Leu Leu Gly Tyr Gly Ile Phe
 115 120 125

Thr Leu Asp Gly Glu Gly Trp Lys His Ser Arg Ala Met Leu Arg Pro
 130 135 140

Gln Phe Ala Arg Glu Gln Val Ala His Val Thr Ser Leu Glu Pro His
 145 150 155 160

Phe Gln Leu Leu Lys Lys His Ile Leu Lys His Lys Gly Glu Tyr Phe
 165 170 175

Asp Ile Gln Glu Leu Phe Phe Arg Phe Thr Val Asp Ser Ala Thr Glu
 180 185 190

Phe Leu Phe Gly Glu Ser Val His Ser Leu Arg Asp Glu Glu Ile Gly
 195 200 205

Tyr Asp Thr Lys Asp Met Ala Glu Glu Arg Arg Lys Phe Ala Asp Ala
 210 215 220

Phe Asn Lys Ser Gln Val Tyr Leu Ser Thr Arg Val Ala Leu Gln Thr
 225 230 235 240

Leu Tyr Trp Leu Val Asn Asn Lys Glu Phe Lys Glu Cys Asn Asp Ile
 245 250 255

Val His Lys Phe Thr Asn Tyr Tyr Val Gln Lys Ala Leu Asp Ala Thr
 260 265 270

Pro Glu Glu Leu Glu Lys Gln Gly Gly Tyr Val Phe Leu Tyr Glu Leu
 275 280 285

Ala Lys Gln Thr Lys Asp Pro Asn Val Leu Arg Asp Gln Ser Leu Asn
 290 295 300

Ile Leu Leu Ala Gly Arg Asp Thr Thr Ala Gly Leu Leu Ser Phe Ala
 305 310 315 320

Val Phe Glu Leu Ala Arg Asn Pro His Ile Trp Ala Lys Leu Arg Glu
 325 330 335

Glu Ile Glu Ser His Phe Gly Ser Gly Glu Asp Ser Arg Val Glu Glu
 340 345 350

Ile Thr Phe Glu Ser Leu Lys Arg Cys Glu Tyr Leu Lys Ala Val Leu
355 360 365

Asn Glu Thr Leu Arg Leu His Pro Ser Val Pro Arg Asn Ala Arg Phe
370 375 380

Ala Ile Lys Asp Thr Thr Leu Pro Arg Gly Gly Gly Pro Asn Gly Lys
385 390 395 400

Asp Pro Ile Leu Ile Arg Lys Asn Glu Val Val Gln Tyr Ser Ile Ser
405 410 415

Ala Thr Gln Thr Asn Pro Ala Tyr Tyr Gly Ala Asp Ala Ala Asp Phe
420 425 430

Arg Pro Glu Arg Trp Phe Glu Pro Ser Thr Arg Asn Leu Gly Trp Ala
435 440 445

Tyr Leu Pro Phe Asn Gly Gly Pro Arg Ile Cys Leu Gly Gln Gln Phe
450 455 460

Ala Leu Thr Glu Ala Gly Tyr Val Leu Val Arg Leu Val Gln Glu Phe
465 470 475 480

Pro Ser Leu Ser Gln Asp Pro Glu Thr Glu Tyr Pro Pro Pro Arg Leu
485 490 495

Ala His Leu Thr Met Cys Leu Phe Asp Gly Ala Tyr Val Lys Met Gln
500 505 510

<210> 116

<211> 499

<212> PRT

<213> CANDIDATROPICALIS

<400> 116

Met Ala Ile Ser Ser Leu Leu Ser Trp Asp Val Ile Cys Val Val Phe
1 5 10 15

Ile Cys Val Cys Val Tyr Phe Gly Tyr Glu Tyr Cys Tyr Thr Lys Tyr
20 25 30

Leu Met His Lys His Gly Ala Arg Glu Ile Glu Asn Val Ile Asn Asp
35 40 45

Gly Phe Phe Gly Phe Arg Leu Pro Leu Leu Leu Met Arg Ala Ser Asn
 50 55 60

Glu Gly Arg Leu Ile Glu Phe Ser Val Lys Arg Phe Glu Ser Ala Pro
 65 70 75 80

His Pro Gln Asn Lys Thr Leu Val Asn Arg Ala Leu Ser Val Pro Val
 85 90 95

Ile Leu Thr Lys Asp Pro Val Asn Ile Lys Ala Met Leu Ser Thr Gln
 100 105 110

Phe Asp Asp Phe Ser Leu Gly Leu Arg Leu His Gln Phe Ala Pro Leu
 115 120 125

Leu Gly Lys Gly Ile Phe Thr Leu Asp Gly Pro Glu Trp Lys Gln Ser
 130 135 140

Arg Ser Met Leu Arg Pro Gln Phe Ala Lys Asp Arg Val Ser His Ile
 145 150 155 160

Ser Asp Leu Glu Pro His Phe Val Leu Leu Arg Lys His Ile Asp Gly
 165 170 175

His Asn Gly Asp Tyr Phe Asp Ile Gln Glu Leu Tyr Phe Arg Phe Ser
 180 185 190

Met Asp Val Ala Thr Gly Phe Leu Phe Gly Glu Ser Val Gly Ser Leu
 195 200 205

Lys Asp Glu Asp Ala Arg Phe Ser Glu Ala Phe Asn Glu Ser Gln Lys
 210 215 220

Tyr Leu Ala Thr Arg Ala Thr Leu His Glu Leu Tyr Phe Leu Cys Asp
 225 230 235 240

Gly Phe Arg Phe Arg Gln Tyr Asn Lys Val Val Arg Lys Phe Cys Ser
 245 250 255

Gln Cys Val His Lys Ala Leu Asp Val Ala Pro Glu Asp Thr Ser Glu
 260 265 270

Tyr Val Phe Leu Arg Glu Leu Val Lys His Thr Arg Asp Pro Val Val
 275 280 285

Leu Gln Asp Gln Ala Leu Asn Val Leu Leu Ala Gly Arg Asp Thr Thr
 290 295 300

Ala Ser Leu Leu Ser Phe Ala Thr Phe Glu Leu Ala Arg Asn Asp His
 305 310 315 320

Met Trp Arg Lys Leu Arg Glu Glu Val Ile Ser Thr Met Gly Pro Ser
 325 330 335

Ser Asp Glu Ile Thr Val Ala Gly Leu Lys Ser Cys Arg Tyr Leu Lys
 340 345 350

Ala Ile Leu Asn Glu Thr Leu Arg Leu Tyr Pro Ser Val Pro Arg Asn
 355 360 365

Ala Arg Phe Ala Thr Arg Asn Thr Thr Leu Pro Arg Gly Gly Gly Pro
 370 375 380

Asp Gly Ser Phe Pro Ile Leu Ile Arg Lys Gly Gln Pro Val Gly Tyr
 385 390 395 400

Phe Ile Cys Ala Thr His Leu Asn Glu Lys Val Tyr Gly Asn Asp Ser
 405 410 415

His Val Phe Arg Pro Glu Arg Trp Ala Ala Leu Glu Gly Lys Ser Leu
 420 425 430

Gly Trp Ser Tyr Leu Pro Phe Asn Gly Gly Pro Arg Ser Cys Leu Gly
 435 440 445

Gln Gln Phe Ala Ile Leu Glu Ala Ser Tyr Val Leu Ala Arg Leu Thr
 450 455 460

Gln Cys Tyr Thr Thr Ile Gln Leu Arg Thr Thr Glu Tyr Pro Pro Lys
 465 470 475 480

Lys Leu Val His Leu Thr Met Ser Leu Leu Asn Gly Val Tyr Ile Arg
 485 490 495

Thr Arg Thr

<210> 117
 <211> 679
 <212> PRT
 <213> CANDIDATROPICALIS

<400> 117

Met Ala Leu Asp Lys Leu Asp Leu Tyr Val Ile Ile Thr Leu Val Val
1 5 10 15

Ala Val Ala Ala Tyr Phe Ala Lys Asn Gln Phe Leu Asp Gln Pro Gln
20 25 30

Asp Thr Gly Phe Leu Asn Thr Asp Ser Gly Ser Asn Ser Arg Asp Val
35 40 45

Leu Ser Thr Leu Lys Lys Asn Asn Lys Asn Thr Leu Leu Leu Phe Gly
50 55 60

Ser Gln Thr Gly Thr Ala Glu Asp Tyr Ala Asn Lys Leu Ser Arg Glu
65 70 75 80

Leu His Ser Arg Phe Gly Leu Lys Thr Met Val Ala Asp Phe Ala Asp
85 90 95

Tyr Asp Trp Asp Asn Phe Gly Asp Ile Thr Glu Asp Ile Leu Val Phe
100 105 110

Phe Ile Val Ala Thr Tyr Gly Glu Gly Glu Pro Thr Asp Asn Ala Asp
115 120 125

Glu Phe His Thr Trp Leu Thr Glu Glu Ala Asp Thr Leu Ser Thr Leu
130 135 140

Lys Tyr Thr Val Phe Gly Leu Gly Asn Ser Thr Tyr Glu Phe Phe Asn
145 150 155 160

Ala Ile Gly Arg Lys Phe Asp Arg Leu Leu Ser Glu Lys Gly Gly Asp
165 170 175

Arg Phe Ala Glu Tyr Ala Glu Gly Asp Asp Gly Thr Gly Thr Leu Asp
180 185 190

Glu Asp Phe Met Ala Trp Lys Asp Asn Val Phe Asp Ala Leu Lys Asn
195 200 205

Asp Leu Asn Phe Glu Glu Lys Glu Leu Lys Tyr Glu Pro Asn Val Lys
210 215 220

Leu Thr Glu Arg Asp Asp Leu Ser Ala Ala Asp Ser Gln Val Ser Leu
225 230 235 240

Gly Glu Pro Asn Lys Lys Tyr Ile Asn Ser Glu Gly Ile Asp Leu Thr
 245 250 255

Lys Gly Pro Phe Asp His Thr His Pro Tyr Leu Ala Arg Ile Thr Glu
 260 265 270

Thr Arg Glu Leu Phe Ser Ser Lys Asp Arg His Cys Ile His Val Glu
 275 280 285

Phe Asp Ile Ser Glu Ser Asn Leu Lys Tyr Thr Thr Gly Asp His Leu
 290 295 300

Ala Ile Trp Pro Ser Asn Ser Asp Glu Asn Ile Lys Gln Phe Ala Lys
 305 310 315 320

Cys Phe Gly Leu Glu Asp Lys Leu Asp Thr Val Ile Glu Leu Lys Ala
 325 330 335

Leu Asp Ser Thr Tyr Thr Ile Pro Phe Pro Thr Pro Ile Thr Tyr Gly
 340 345 350

Ala Val Ile Arg His His Leu Glu Ile Ser Gly Pro Val Ser Arg Gln
 355 360 365

Phe Phe Leu Ser Ile Ala Gly Phe Ala Pro Asp Glu Glu Thr Lys Lys
 370 375 380

Ala Phe Thr Arg Leu Gly Gly Asp Lys Gln Glu Phe Ala Ala Lys Val
 385 390 395 400

Thr Arg Arg Lys Phe Asn Ile Ala Asp Ala Leu Leu Tyr Ser Ser Asn
 405 410 415

Asn Ala Pro Trp Ser Asp Val Pro Phe Glu Phe Leu Ile Glu Asn Val
 420 425 430

Pro His Leu Thr Pro Arg Tyr Tyr Ser Ile Ser Ser Ser Ser Leu Ser
 435 440 445

Glu Lys Gln Leu Ile Asn Val Thr Ala Val Val Glu Ala Glu Glu Glu
 450 455 460

Ala Asp Gly Arg Pro Val Thr Gly Val Val Thr Asn Leu Leu Lys Asn
 465 470 475 480

Val Glu Ile Val Gln Asn Lys Thr Gly Glu Lys Pro Leu Val His Tyr
485 490 495

Asp Leu Ser Gly Pro Arg Gly Lys Phe Asn Lys Phe Lys Leu Pro Val
500 505 510

His Val Arg Arg Ser Asn Phe Lys Leu Pro Lys Asn Ser Thr Thr Pro
515 520 525

Val Ile Leu Ile Gly Pro Gly Thr Gly Val Ala Pro Leu Arg Gly Phe
530 535 540

Val Arg Glu Arg Val Gln Gln Val Lys Asn Gly Val Asn Val Gly Lys
545 550 555 560

Thr Leu Leu Phe Tyr Gly Cys Arg Asn Ser Asn Glu Asp Phe Leu Tyr
565 570 575

Lys Gln Glu Trp Ala Glu Tyr Ala Ser Val Leu Gly Glu Asn Phe Glu
580 585 590

Met Phe Asn Ala Phe Ser Arg Gln Asp Pro Ser Lys Lys Val Tyr Val
595 600 605

Gln Asp Lys Ile Leu Glu Asn Ser Gln Leu Val His Glu Leu Leu Thr
610 615 620

Glu Gly Ala Ile Ile Tyr Val Cys Gly Asp Ala Ser Arg Met Ala Arg
625 630 635 640

Asp Val Gln Thr Thr Ile Ser Lys Ile Val Ala Lys Ser Arg Glu Ile
645 650 655

Ser Glu Asp Lys Ala Ala Glu Leu Val Lys Ser Trp Lys Val Gln Asn
660 665 670

Arg Tyr Gln Glu Asp Val Trp
675

<210> 118

<211> 679

<212> PRT

<213> CANDIDATROPICALIS

<400> 118

Met Ala Leu Asp Lys Leu Asp Leu Tyr Val Ile Ile Thr Leu Val Val
 1 5 10 15
 Ala Val Ala Ala Tyr Phe Ala Lys Asn Gln Phe Leu Asp Gln Pro Gln
 20 25 30
 Asp Thr Gly Phe Leu Asn Thr Asp Ser Gly Ser Asn Ser Arg Asp Val
 35 40 45
 Leu Ser Thr Leu Lys Lys Asn Asn Lys Asn Thr Leu Leu Leu Phe Gly
 50 55 60
 Ser Gln Thr Gly Thr Ala Glu Asp Tyr Ala Asn Lys Leu Ser Arg Glu
 65 70 75 80
 Leu His Ser Arg Phe Gly Leu Lys Thr Met Val Ala Asp Phe Ala Asp
 85 90 95
 Tyr Asp Trp Asp Asn Phe Gly Asp Ile Thr Glu Asp Ile Leu Val Phe
 100 105 110
 Phe Ile Val Ala Thr Tyr Gly Glu Gly Glu Pro Thr Asp Asn Ala Asp
 115 120 125
 Glu Phe His Thr Trp Leu Thr Glu Glu Ala Asp Thr Leu Ser Thr Leu
 130 135 140
 Arg Tyr Thr Val Phe Gly Leu Gly Asn Ser Thr Tyr Glu Phe Phe Asn
 145 150 155 160
 Ala Ile Gly Arg Lys Phe Asp Arg Leu Leu Ser Glu Lys Gly Gly Asp
 165 170 175
 Arg Phe Ala Glu Tyr Ala Glu Gly Asp Asp Gly Thr Gly Thr Leu Asp
 180 185 190
 Glu Asp Phe Met Ala Trp Lys Asp Asn Val Phe Asp Ala Leu Lys Asn
 195 200 205
 Asp Leu Asn Phe Glu Glu Lys Glu Leu Lys Tyr Glu Pro Asn Val Lys
 210 215 220
 Leu Thr Glu Arg Asp Asp Leu Ser Ala Ala Asp Ser Gln Val Ser Leu
 225 230 235 240

Gly Glu Pro Asn Lys Lys Tyr Ile Asn Ser Glu Gly Ile Asp Leu Thr
 245 250 255

Lys Gly Pro Phe Asp His Thr His Pro Tyr Leu Ala Arg Ile Thr Glu
 260 265 270

Thr Arg Glu Leu Phe Ser Ser Lys Glu Arg His Cys Ile His Val Glu
 275 280 285

Phe Asp Ile Ser Glu Ser Asn Leu Lys Tyr Thr Thr Gly Asp His Leu
 290 295 300

Ala Ile Trp Pro Ser Asn Ser Asp Glu Asn Ile Lys Gln Phe Ala Lys
 305 310 315 320

Cys Phe Gly Leu Glu Asp Lys Leu Asp Thr Val Ile Glu Leu Lys Ala
 325 330 335

Leu Asp Ser Thr Tyr Thr Ile Pro Phe Pro Thr Pro Ile Thr Tyr Gly
 340 345 350

Ala Val Ile Arg His His Leu Glu Ile Ser Gly Pro Val Ser Arg Gln
 355 360 365

Phe Phe Leu Ser Ile Ala Gly Phe Ala Pro Asp Glu Glu Thr Lys Lys
 370 375 380

Thr Phe Thr Arg Leu Gly Gly Asp Lys Gln Glu Phe Ala Thr Lys Val
 385 390 395 400

Thr Arg Arg Lys Phe Asn Ile Ala Asp Ala Leu Leu Tyr Ser Ser Asn
 405 410 415

Asn Thr Pro Trp Ser Asp Val Pro Phe Glu Phe Leu Ile Glu Asn Ile
 420 425 430

Gln His Leu Thr Pro Arg Tyr Tyr Ser Ile Ser Ser Ser Ser Leu Ser
 435 440 445

Glu Lys Gln Leu Ile Asn Val Thr Ala Val Val Glu Ala Glu Glu Glu
 450 455 460

Ala Asp Gly Arg Pro Val Thr Gly Val Val Thr Asn Leu Leu Lys Asn
 465 470 475 480

Ile Glu Ile Ala Gln Asn Lys Thr Gly Glu Lys Pro Leu Val His Tyr
 485 490 495

Asp Leu Ser Gly Pro Arg Gly Lys Phe Asn Lys Phe Lys Leu Pro Val
 500 505 510

His Val Arg Arg Ser Asn Phe Lys Leu Pro Lys Asn Ser Thr Thr Pro
 515 520 525

Val Ile Leu Ile Gly Pro Gly Thr Gly Val Ala Pro Leu Arg Gly Phe
 530 535 540

Val Arg Glu Arg Val Gln Gln Val Lys Asn Gly Val Asn Val Gly Lys
 545 550 555 560

Thr Leu Leu Phe Tyr Gly Cys Arg Asn Ser Asn Glu Asp Phe Leu Tyr
 565 570 575

Lys Gln Glu Trp Ala Glu Tyr Ala Ser Val Leu Gly Glu Asn Phe Glu
 580 585 590

Met Phe Asn Ala Phe Ser Arg Gln Asp Pro Ser Lys Lys Val Tyr Val
 595 600 605

Gln Asp Lys Ile Leu Glu Asn Ser Gln Leu Val His Glu Leu Leu Thr
 610 615 620

Glu Gly Ala Ile Ile Tyr Val Cys Gly Asp Ala Ser Arg Met Ala Arg
 625 630 635 640

Asp Val Gln Thr Thr Ile Ser Lys Ile Val Ala Lys Ser Arg Glu Ile
 645 650 655

Ser Glu Asp Lys Ala Ala Glu Leu Val Lys Ser Trp Lys Val Gln Asn
 660 665 670

Arg Tyr Gln Glu Asp Val Trp
 675